

Marquette University

e-Publications@Marquette

---

Exercise Science Faculty Research and  
Publications

Exercise Science, Department of

---

1-2010

## Response to task-specific sex differences in muscle fatigue: Is there a common underlying cause?

Sandra K. Hunter

Marquette University, [sandra.hunter@marquette.edu](mailto:sandra.hunter@marquette.edu)

Follow this and additional works at: [https://epublications.marquette.edu/exsci\\_fac](https://epublications.marquette.edu/exsci_fac)



Part of the [Physical Therapy Commons](#)

---

### Recommended Citation

Hunter, Sandra K., "Response to task-specific sex differences in muscle fatigue: Is there a common underlying cause?" (2010). *Exercise Science Faculty Research and Publications*. 13.

[https://epublications.marquette.edu/exsci\\_fac/13](https://epublications.marquette.edu/exsci_fac/13)

Marquette University

e-Publications@Marquette

## ***Exercise Science Faculty Research and Publications/College of Health Sciences***

***This paper is NOT THE PUBLISHED VERSION; but the author's final, peer-reviewed manuscript.*** The published version may be accessed by following the link in the citation below.

*Exercise and Sport Sciences Reviews*, Vol. 38, No. 1 (January 2010): 37. [DOI](#). This article is © The American College of Sports Medicine and permission has been granted for this version to appear in [e-Publications@Marquette](#). The American College of Sports Medicine does not grant permission for this article to be further copied/distributed or hosted elsewhere without the express permission from The American College of Sports Medicine.

# Response

Sandra K. Hunter

Exercise Science Program, Department of Physical Therapy, Marquette University, Milwaukee, WI

Dear Editor-in-Chief:

I thank Mr. Dotan and Dr. Falk for their interest and comments on my article. They propose that sex differences in muscle fatigue are due to differences in the ability of men and women to recruit and use fast-contracting, Type II motor units. It is tempting and more simplistic to suggest one global mechanism to explain the difference in muscle fatigue between men and women for different tasks. Although men may have a larger proportional area of fast twitch fibers in their muscle than women for some muscles (see my previous article), the available evidence suggests that there is little sex difference in motor unit recruitment and rate coding during maximal and submaximal contractions in the absence of fatigue and also during an isometric fatiguing contraction.

Unlike children <sup>(7)</sup>, for example, adult men and women have a similar ability to activate their available motor units during maximal strength contractions before a fatiguing task for various muscle groups within the upper and lower limbs <sup>(1,3,5,8)</sup>. This is demonstrated with experiments that have quantified voluntary activation using a combination of stimulation of the nervous system and voluntary contractions. The reduction in voluntary activation that occurs with fatigue also is similar in men and women for the elbow flexor muscles <sup>(1,8)</sup> but greater

for men than women for muscles in the lower limb<sup>(3,5)</sup>. For leg muscles, but not elbow flexor muscles, the greater fatigue of the men than women is in part due to a greater loss of neural drive to the muscle for the men. Thus, men seem more limited in their ability to recruit fast-contracting motor units with fatigue for some muscles compared with women.

Furthermore, single motor unit experiments that are used to quantify motor unit recruitment and rate coding also show no systematic sex differences that would explain the difference in muscle fatigue between men and women<sup>(4,6)</sup>. For example, motor unit behavior of biceps brachii was examined during submaximal isometric contractions of the elbow flexor muscles sustained until failure in men and women<sup>(4)</sup>. Although the women had a longer time to task failure than the men, the mean motor unit recruitment threshold, mean discharge rate, and the decline in discharge rate during the fatiguing contractions were similar for the men and women<sup>(4)</sup>. In support of these findings, the transient recruitment of motor units that typically increases during a fatiguing contraction (and quantified as the bursting activity of the interference electromyography signal) shows no systematic sex difference in the increase during an isometric fatiguing contraction<sup>(2)</sup>.

Collectively, these studies provide initial evidence that the sex difference in muscle fatigue for various muscle groups cannot be explained by one global mechanism under the umbrella of "neuromuscular activation." Further studies to understand single motor unit behavior of men and women during maximal contractions and various other types of fatiguing tasks would, however, complement the limited number of studies on isometric contractions that are highlighted.

Sandra K. Hunter, Ph.D., FACSM  
Exercise Science Program  
Department of Physical Therapy  
Marquette University  
Milwaukee, WI, United States

## References

1. Hunter SK, Butler JE, Todd G, Gandevia SC, Taylor JL. Supraspinal fatigue does not explain the sex difference in muscle fatigue of maximal contractions. *J Appl Physiol.* 2006; 101:1036-44.
2. Hunter SK, Critchlow A, Enoka RM. Influence of aging on sex differences in muscle fatigability. *J Appl Physiol.* 2004; 97:1723-32.
3. Martin PG, Rattey J. Central fatigue explains sex differences in muscle fatigue and contralateral cross-over effects of maximal contractions. *Pflugers Arch.* 2007; 454:957-69.
4. Mottram CJ, Hunter SK, Rochette L, Anderson MK, Enoka RM. Time to task failure varies with the gain of the feedback signal for women, but not for men. *Exp Brain Res.* 2006; 174:575-87.
5. Russ DW, Kent-Braun JA. Sex differences in human skeletal muscle fatigue are eliminated under ischemic conditions. *J Appl Physiol.* 2003; 94:2414-22.
6. Spiegel KM, Stratton J, Burke JR, Glendinning DS, Enoka RM. The influence of age on the assessment of motor unit activation in a human hand muscle. *Exp Physiol.* 1996; 81:805-19.
7. Streckis V, Skurvydas A, Ratkevicius A. Children are more susceptible to central fatigue than adults. *Muscle Nerve.* 2007; 36:357-63.
8. Yoon T, Schlinder Delap B, Griffith EE, Hunter SK. Mechanisms of fatigue differ after low- and high-force fatiguing contractions in men and women. *Muscle Nerve.* 2007; 36:512-24.