Marquette University

e-Publications@Marquette

Physics Faculty Research and Publications

Physics, Department of

8-2009

Motion of the Zinc Ions in Catalysis by a Dizinc Metallo- β -Lactamase

Robert M. Breece

University of New Mexico

Zhenxin Hu

Miami University - Oxford

Brian Bennett

Marquette University, brian.bennett@marquette.edu

Michael W. Crowder

Miami University - Oxford

David L. Tierney

University of New Mexico

Follow this and additional works at: https://epublications.marquette.edu/physics_fac

Part of the [Physics Commons](#)

Recommended Citation

Breece, Robert M.; Hu, Zhenxin; Bennett, Brian; Crowder, Michael W.; and Tierney, David L., "Motion of the Zinc Ions in Catalysis by a Dizinc Metallo- β -Lactamase" (2009). *Physics Faculty Research and Publications*. 17.

https://epublications.marquette.edu/physics_fac/17

Motion of the Zinc Ions in Catalysis by a Dizinc Metallo- β -Lactamase

Robert M. Breece

*Department of Chemistry and Chemical Biology,
University of New Mexico,
Albuquerque, NM*

Zhenxin Hu

*Department of Chemistry and Biochemistry, Miami University,
Oxford, OH*

Brian Bennett

*Department of Biophysics, Medical College of Wisconsin,
Milwaukee, WI*

Michael W. Crowder

*Department of Chemistry and Biochemistry, Miami University,
Oxford, OH*

David L. Tierney

*Department of Chemistry and Chemical Biology,
University of New Mexico,
Albuquerque, NM*

*Department of Chemistry and Biochemistry, Miami University,
Oxford, OH*

Abstract

We report rapid-freeze-quench X-ray absorption spectroscopy of a dizinc metallo- β -lactamase (M β L) reaction intermediate. The Zn(II) ions in the dinuclear active site of the *S. maltophilia* Class B3 M β L move away from each other, by $\sim 0.3 \text{ \AA}$ after 10 ms of reaction with nitrocefins, from 3.4 to 3.7 $\text{\AA}$. Together with our previous characterization of the resting enzyme and its nitrocefin product complex, where the Zn(II) ion separation relaxes to 3.6 $\text{\AA}$, these data indicate a scissoring motion of the active site that accompanies the ring-opening step. The average Zn(II) coordination number of 4.5 in the resting enzyme appears to be maintained throughout the reaction with nitrocefin. This is the first direct structural information available on early stage dizinc metallo- β -lactamase catalysis.

β -Lactam containing compounds remain the most widely used antibiotics, despite the growing number of bacteria that show resistance to this group of compounds. The most common resistance strategy is secretion of β -lactamases, which hydrolyze the antibiotic's four-membered β -lactam ring, leading to an inactive product. A subset of these enzymes (class B of subclasses A–D) require Zn(II) for activity. These metallo- β -lactamases (M β Ls) are further subdivided into three subclasses, based on sequence homology.^{1,2} L1, a class B3 M β L from *Stenotrophomonas maltophilia* requires two Zn(II) ions for maximal activity.³ The crystal structure of L1 (Protein Data Bank entry pdb 1SML) shows two distinct Zn(II) ions: Zn₁ is tetrahedrally coordinated by three histidines and one solvent molecule that bridges to Zn₂, which is five-coordinate, ligated by two histidines, a monodentate aspartate, one terminally bound water, and the bridging solvent molecule.⁴ The crystallographic Zn₁–Zn₂ separation of 3.45 $\text{\AA}$ is similar to that determined by EXAFS (3.42 $\text{\AA}$) of the same enzyme.⁵

The observation of product inhibition⁶ allows for characterization of product complexes, and a more recent crystal structure of ZnZn-L1, complexed with hydrolyzed moxalactam (pdb 2AIO), showed that the distance between the two Zn(II) ions had increased to 3.68 Å, due to a partial rotation of the bound product, made possible by the ring-opening step.⁷ A similar increase in metal–metal distance was observed in extended X-ray absorption fine structure (EXAFS) studies of L1 complexed with hydrolyzed nitrocefin (3.62 Å), which rotates to a greater degree with ultimate release of the product's auxiliary carboxylate in favor of the ring sulfur, as illustrated in Figure 1. The increased M–M distance, without a change in coordination number (as evidenced by a lack of significant changes in the average first shell Zn–L distance), was taken as evidence that the bridging hydroxide was absent in the L1–nitrocefin product complex. A solvent-derived bridge was observed in the moxalactam-bound crystal structure, where the average Zn(II) coordination number had increased from 4.5 to 5.5.⁷

Figure 1. Summary of the time-dependent structure of resting ZnZn-L1 (left), freeze quenched after 10 ms of reaction with nitrocefin (center), and the resulting product complex (right) derived from EXAFS. The structure of nitrocefin, shown above the resting structure, is truncated in the 10 ms and product structures. The Zn–Zn separations are given below each structure.

Rotation of the product, as observed in both the EXAFS and diffraction studies, makes structural characterization of reaction intermediates that precede this rotation critical to understanding their action. Crystal structures of MβLs complexed with substrate or trapped during catalysis have not been reported. To date, the only direct structural information available on MβL catalysis comes from EPR studies of a mixed-metal ZnCo hybrid of L1⁸ and the di-Co(II)

substituted form of the related Class B1 enzyme BcII from *Bacillus cereus*,⁹ both of which demonstrate a clear interaction of substrate with both metal ions during turnover. However, similar information on the native Zn(II)-containing enzymes is not available. To probe the relative positions of Zn₁ and Zn₂ during catalysis, we examined the EXAFS of 0.5 mM ZnZn-L1 rapid-freeze-quenched after 10 ms of reaction with 2.5 mM nitrocefin at 2 °C.¹⁰ The 10 ms quench time closely mirrors stopped-flow studies of the same system, which show maximum concentration of the nitrocefin-derived reaction intermediate at 10–20 ms of reaction time.⁶ The present studies mark the first structural characterization of a true Zn(II)-containing MβL reaction intermediate.

Comparing the Fourier transformed EXAFS of the 10 ms freeze-quenched sample with the EXAFS of the resting enzyme (Figure 2, top) shows (i) an increase in the amplitude associated with the first shell, without a change in peak position or fwhm; (ii) the loss of a feature at $R + a \approx 3.0$ Å; and (iii) increased magnitude at $R + a \approx 2.2$ and 3.4 Å. This qualitative description is consistent with curve fitting results (Figure S1 and Table S1), which indicate that the average coordination number of Zn is retained (i) and that the Zn–Zn distance increases by nearly 0.3 Å (ii and iii). The larger feature at 2.2 Å in the freeze-quenched FT is attributed to a more ordered Zn–C_{CO2}[−] interaction, compared to the resting enzyme.

Figure 2. Comparison of the Fourier transformed EXAFS of ZnZn-L1, freeze quenched after 10 ms of reaction with nitrocefin (bold line), with resting ZnZn-L1 (top) and the ZnZn-L1 product complex with nitrocefin (center), and the best fit to the data (bottom). The data for ZnZnL1 and the ZnZnL1-nitrocefin product complex are reproduced, for comparison, from Costello, et al.⁵

Table 1. EXAFS-Derived Structural Models for ZnZn-L1 in the Resting State,^a Freeze-Quenched after 10 ms of Reaction with Nitrocefin,^b and Its Product Complex^a with Hydrolyzed Nitrocefin

sample	Zn–L ^c	Zn–His	Zn–Zn (Å)
resting ^a	4 N/O @ 2.02 Å	2.5	3.42
10 ms ^b	4.5 N/O @ 2.03 Å	2.5	3.72
product ^a	4 N/O @ 2.04 Å 0.5 S @ 2.29 Å	2.5	3.62

^aFrom Costello, et al.⁵

^bThis work. See Supporting Information for details.

^cZn–L distances ± 0.01 Å; Zn–Zn distances ± 0.02 Å.

The lack of any apparent change in coordination number has significant mechanistic implications. Without changing the coordination number of either Zn ion, the presence of a bridging reaction intermediate, as implied by a relatively well-ordered Zn–Zn vector, requires the loss of two Zn ligands from the resting structure. The first most likely comes from the loss of one bond to the bridging solvent, which is incorporated into the product as part of the newly formed carboxylate and remains coordinated in the product complex.⁷

In several mechanistic models,¹¹ the bond between Zn₂ and its terminal solvent ligand is lost during the catalytic cycle, though the bond is reformed on product release. Loss of this solvent ligand is required to maintain the Zn coordination number and still form a bond to the anionic nitrogen that results from ring opening. We show these two bonds as dashed in Figure 1, only in that we have no direct evidence for one over the other. However, given the well-established relationship between the protonation state of the ring nitrogen and the blue color of the intermediate¹² (the same color displayed by the present set of samples), we must at present favor loss of the solvent ligand.

Comparison with the product complex shows the intermediate has a similar first shell amplitude, but the main peak in the product complex FT is shifted to higher *R*, due to incorporation of the product sulfur atom into the Zn₂ coordination sphere. With no evidence of a large atom scatterer in the first shell of the intermediate, we must conclude that product rotation, leading to a measurable Zn–S interaction, has not yet occurred at 10 ms and that the apparent

increase in first shell amplitude for the intermediate is therefore due to lower disorder in the first shell of the intermediate. The outer shell scattering of the product complex is of lower amplitude than in the intermediate, also consistent with a more ordered site in the intermediate. This may be reflective of a less planar orientation of the product, relative to substrate and/or intermediate.

Meanwhile, the dramatic increase in Zn–Zn separation to 3.72 Å suggests that, even at 10 ms, the Zn–(OH)–Zn bridge has been lost (this nearly linear arrangement is expected to lead to more dramatic metal–oxygen–metal scattering than is observed). We note that a second, much shallower minimum was observed, with a Zn–Zn separation of 3.37 Å. This distance is more in line with that of the resting enzyme, although it is closer to that observed in similar systems with a bridging bidentate carboxylate,¹³ as opposed to the terminal monodentate carboxylate expected in the MβL active site. In either case, inclusion of two Zn–Zn populations, suggests that such a species represents no more than ~11% of the total enzyme present.

The motion of the Zn ions has potentially large mechanistic implications. The two metals are expected to be anchored to the auxiliary carboxylate of the fused ring (Zn₂) and the lactam carbonyl (Zn₁). The 0.3 Å movement of the metal ions, away from each other, suggests that binding of substrate, and possibly release of the bridging hydroxyl, is accompanied by a scissoring motion that exerts added pressure on the N–C(=O) bond, helping facilitate the ring-opening reaction. This proposal is consistent with prior observations that suggest all mono-Zn MβLs are active^{14,15} and that inclusion of the second metal ion increases the enzyme's catalytic efficiency, without affecting the overall rates of reaction.⁵ Further experiments are planned for the mixed-metal hybrid, which will allow us to assess structural changes at each metal site, independently.

Acknowledgment

This work was supported by the National Institutes of Health (NCRR P20RR-16480 to D.L.T. and AI056231 to B.B.), by the Volwiler Professorship at Miami University (to M.W.C.), and by the National Science Foundation (CHE0809985 to D.L.T.).

Supporting Information

Experimental procedures, including sample preparation and EXAFS data collection and analysis, along with Figure S1 and Table S1, showing detailed EXAFS curve fitting results, are included as Supporting Information. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- ¹Galleni, M., Lamotte-Brasseur, J., Rossolini, G. M., Spencer, J., Dideberg, O. and Frere, J.-M. *Antimicrob. Agents Chemother.* 2001, 45, 660– 663
- ²Bebrone, C. *Biochem. Pharmacol.* 2007, 74, 1686– 1701
- ³Periyannan, G., Shaw, P. J., Sigdel, T. and Crowder, M. W. *Protein Sci.* 2004, 13, 2236– 2243
- ⁴Ullah, J. H., Walsh, T. R., Taylor, I. A., Emery, D. C., Verma, C. S., Gamblin, S. J. and Spenser, J. *J. Mol. Biol.* 1998, 287, 125– 136
- ⁵Costello, A. L., Periyannan, G., Yang, K.-W., Crowder, M. W. and Tierney, D. L. *J. Biol. Inorg. Chem.* 2006, 11, 351– 358
- ⁶McManus-Munoz, S. and Crowder, M. W. *Biochemistry* 1999, 38, 1547– 1553
- ⁷Spencer, J., Read, J., Sessions, R. B., Howell, S., Blackburn, G. M. and Gamblin, S. J. *J. Am. Chem. Soc.* 2005, 127, 14439– 14444
- ⁸Hu, Z., Periyannan, G., Bennett, B. and Crowder, M. W. *J. Am. Chem. Soc.* 2008, 130, 14207– 14216
- ⁹Tioni, M. F., Llarrull, L. I., Poeylout-Palena, A. A., Marti, M. A., Saggi, M., Periyannan, G. R., Mata, E. G., Bennett, B., Murgida, D. H. and Vila, A. J. *J. Am. Chem. Soc.* 2008, 130, 15852– 15863
- ¹⁰Garritty, J. D., Bennett, B. and Crowder, M. W. *Biochemistry* 2005, 44, 1078– 1087
- ¹¹Xu, D., Guo, H. and Cui, Q. *J. Am. Chem. Soc.* 2007, 129, 10814– 10822
- ¹²Kaminskaia, N. V., Spingler, B. and Lippard, S. J. *J. Am. Chem. Soc.* 2001, 123, 6555– 6563
- ¹³Thomas, P. W., Stone, E. M., Costello, A. L., Tierney, D. L. and Fast, W. *Biochemistry* 2005, 44, 7559– 7565
- ¹⁴Cricco, J. A., Orellano, E. G., Rasia, R. M., Ceccarelli, E. A. and Vila, A. J. *Coord. Chem. Rev.* 1999, 190–192, 519– 535
- ¹⁵Llarrull, L. I., Tioni, M. F. and Vila, A. J. *J. Am. Chem. Soc.* 2008, 130, 15842– 15851