

6-1-2012

Glutathione-Mediated Neuroprotection Against Methylmercury Neurotoxicity in Cortical Culture is Dependent on MRP1

Travis Rush
Marquette University

Xiaoqian Liu
Marquette University, xiaoqian.1.liu@marquette.edu

Andrew B. Nowakowski
University of Wisconsin - Milwaukee

David H. Petering
University of Wisconsin - Milwaukee

Doug Lobner
Marquette University, doug.lobner@marquette.edu

Marquette University

e-Publications@Marquette

Biomedical Sciences Faculty Research and Publications/College of Health Sciences

This paper is NOT THE PUBLISHED VERSION; but the author's final, peer-reviewed manuscript. The published version may be accessed by following the link in the citation below.

NeuroToxicology, Vol. 33, No. 3 (June 2012): 476-481. [DOI](#). This article is © Elsevier and permission has been granted for this version to appear in [e-Publications@Marquette](#). Elsevier does not grant permission for this article to be further copied/distributed or hosted elsewhere without the express permission from Elsevier.

Glutathione-mediated Neuroprotection Against Methylmercury Neurotoxicity in Cortical Culture is Dependent on MRP1

Travis Rush

Department of Biomedical Sciences, Marquette University, Milwaukee, WI

XiaoQian Liu

Department of Biomedical Sciences, Marquette University, Milwaukee, WI

Andrew B. Nowakowski

Department of Chemistry and Biochemistry, University of Wisconsin-Milwaukee, Milwaukee, WI

David H. Petering

Department of Chemistry and Biochemistry, University of Wisconsin-Milwaukee, Milwaukee, WI

Doug Lobner

Department of Biomedical Sciences, Marquette University, Milwaukee, WI

Abstract

Methylmercury (MeHg) exposure at high concentrations poses significant neurotoxic threat to humans worldwide. The present study investigated the mechanisms of glutathione-mediated attenuation of MeHg neurotoxicity in primary cortical culture. MeHg (5 μ M) caused depletion of mono- and disulfide glutathione in

neuronal, glial and mixed cultures. Supplementation with exogenous glutathione, specifically glutathione monoethyl ester (GSHME) protected against the MeHg induced neuronal death. MeHg caused increased [reactive oxygen species](#) (ROS) formation measured by dichlorodihydrofluorescein (DCF) fluorescence with an early increase at 30 min and a late increase at 6 h. This [oxidative stress](#) was prevented by the presence of either GSHME or the free radical scavenger, trolox. While trolox was capable of quenching the ROS, it showed no neuroprotection. Exposure to MeHg at subtoxic concentrations (3 μ M) caused an increase in system x_c^- mediated 14 C-cystine uptake that was blocked by the protein synthesis inhibitor, cycloheximide (CHX). Interestingly, blockade of the early ROS burst prevented the functional upregulation of system x_c^- . Inhibition of multidrug resistance protein-1 (MRP1) potentiated MeHg neurotoxicity and increased cellular MeHg. Taken together, these data suggest glutathione offers neuroprotection against MeHg [toxicity](#) in a manner dependent on MRP1-mediated efflux.

Abbreviations

CPG carboxyphenylglycine
CGN cerebellar granular neurons
CHX cycloheximide
DCF dichlorodihydrofluorescein
GSHME glutathione monoethyl ester
ICP-MS inductively-coupled plasma mass spectrometry
LDH lactate dehydrogenase
MeHg methylmercury
MRP1 multi-drug resistance protein 1
NAC *N*-acetyl cysteine
ROS reactive oxygen species

Keywords

Glutathione, Methylmercury, System x_c^- MRP1, Neurotoxicity, Neuroprotection

1. Introduction

[Methylmercury](#) (MeHg) is an ubiquitous neurotoxicant posing significant threat to [humans](#) when ingested at high concentrations. Exposure to MeHg has been implicated as a factor in the development of [neurodegenerative diseases](#) such as [Alzheimer's Disease](#), [Parkinson's Disease](#), and [amyotrophic lateral sclerosis](#) (ALS) ([Hock et al., 1998](#), [Monnet-Tschudi et al., 2006](#), [Praline et al., 2007](#)). Humans are primarily exposed to MeHg through consumption of [contaminated fish and shellfish](#). MeHg is formed by bacterial [methylation](#) of inorganic [mercury](#) in aquatic [sediments](#) ([Clarkson, 1997](#)). MeHg then travels up the [food chain](#) and accumulates in [fish](#) and [shellfish](#), and is subsequently consumed by humans ([Mahaffey et al., 2004](#)). There is clear evidence for neurological deficits following exposure to high levels of mercury. For example, a large release of mercury into Minamata Bay in Japan led to toxic levels of exposure and severe [injury](#) to the local population, including neurological dysfunctions ([Harada, 1995](#), [Tsuda et al., 2009](#)). In a separate incident, methylmercury-contaminated seed grain was sent to a population in [Iraq](#) with its consumption leading to severe neurological injuries ([Myers et al., 2000](#)). Acute exposure to high levels of mercury is known to target cerebellar granular [neurons](#) ([Fonnum and Lock, 2000](#)), however, due to the location of damage in neurodegenerative diseases, there is also interest in the effects of mercury on cortical neurons. While humans are exposed to other forms of mercury, such as [elemental](#) mercury found in [dental amalgam restorations](#), or ethylmercury found in the commercial [preservative](#), [thimerosal](#), MeHg is of specific concern because it is easily absorbed in the gastrointestinal tract, and readily traverses the [blood–brain barrier](#) (BBB) ([Bridges and Zalups, 2005](#), [Bridges and Zalups, 2010](#)).

Proposed mechanisms for MeHg [neurotoxicity](#) are primarily focused on four possibilities: disruption of intracellular [calcium](#) and [zinc ion homeostasis](#), likely involving mitochondrial deficits ([Atchison and Hare,](#)

1994, [Kawanai et al., 2009](#)), redox imbalance by increased production of [reactive oxygen species](#) and/or by decreasing endogenous cellular [antioxidant defenses](#) ([Ali et al., 1992](#), [Yee and Choi, 1996](#), [Aschner, 2000](#), [Franco et al., 2007](#), [Amonpatumrat et al., 2008](#), [Wang et al., 2009](#)), direct interactions with free [protein sulfhydryl groups](#) ([Rooney, 2007](#)), and inhibition of selenoenzymes, especially those involved in redox signaling ([Carvalho et al., 2008](#)). The present study is aimed at determining the effects of MeHg on the [glutathione](#) (GSH) cycling system and [oxidative stress](#).

[Astrocytes](#) are thought to be responsible for de novo synthesis of [glutathione](#) from [glutamate](#), [glycine](#) and the rate limiting substrate, [cysteine](#), which is brought into the astrocytes primarily in its oxidized form, [cystine](#) ([Kranich et al., 1998](#)). Glutathione can be utilized by the cell to reduce [reactive oxygen species](#), for example, [superoxide](#) produced as a [byproduct](#) of mitochondrial [energy production](#) rapidly reacts to form [hydrogen peroxide](#) which is then reduced by [glutathione](#) to form [glutathione-disulfide](#) (GSSG) and water in a reaction catalyzed by [glutathione peroxidase](#). Glutathione may also be utilized as a [xenobiotic](#) detoxicant as has been well characterized involving chemotherapeutics in [cancer](#) treatment ([Salinas and Wong, 1999](#)). That is, glutathione can be directly conjugated to exogenous substrates via a [disulfide bond](#) with the free sulfhydryl groups; these reactions are directed by a class of [enzymes](#) known as [glutathione-S-transferases](#) (GSTs) ([Dringen, 2000](#), [Dringen and Hirrlinger, 2003](#)). GSH, GSSG and the glutathione-conjugates are then exported from the cell in a glutathione-dependent manner via [multi-drug resistance proteins](#) (MRP), specifically [MRP1](#) in the CNS ([Hirrlinger and Dringen, 2005](#), [Minich et al., 2006](#)). These glutathione [molecules](#) can then be broken down in the extracellular space by [glutathione reductase](#), [amino-peptidase N](#), or [gamma-glutamyl transpeptidase](#). This [metabolism](#) produces the substrate [cysteine](#), which can be taken up and utilized by neurons to produce their own glutathione. In this way, neurons are dependent on astrocytes to supply glutathione ([Dringen et al., 1999](#)). Since glutathione serves a dual role as both an [antioxidant](#) and detoxicant, and both of these roles may offer independent protective mechanisms against MeHg, the present study examines the impact of MeHg on these aspects of glutathione action.

2. Materials and methods

2.1. Materials

Timed pregnant Swiss Webster [mice](#) were obtained from Charles [River](#) Laboratories (Wilmington, DE, USA). [Serum](#) was from Atlanta [Biologicals](#) (Atlanta, GA, USA). [NADPH](#) was from Applichem (Darmstadt, Germany). Radiolabeled ^{14}C -cystine was purchased from PerkinElmer (Boston, MA, USA). S-(4)-carboxyphenylglycine (CPG) and MK571 were obtained from Tocris Bioscience (Ellisville, MO, USA). DCF was from Molecular Probes (Eugene, OR, USA). All other chemicals were from Sigma (St. Louis, MO, USA).

2.2. Cortical cell cultures

Mixed cortical [cell cultures](#) containing glial and neuronal cells were prepared from fetal (15–16 day gestation) [mice](#) as previously described ([Lobner, 2000](#)). Dissociated cortical cells were plated on 24-well plates (2.0 cm² [surface area](#) per well) coated with poly-d-lysine and [laminin](#) in Eagles' Minimal Essential Medium (MEM, Earle's [salts](#), supplied glutamine-free) supplemented with 5% (v/v) heat-inactivated [horse](#) serum, 5% (v/v) [fetal bovine serum](#), 2 mM [glutamine](#) and d-glucose (total 21 mM). Neuronal cultures were prepared exactly as above with the addition of 10 μM [cytosine](#) arabinocide 48 h after [plating](#) to inhibit glial replication ([Dugan et al., 1995](#), [Rush et al., 2010](#)). Cultures were plated at approximately 8 hemispheres per plate in a volume of 500 μl per well. Glial cultures were prepared as described for mixed cultures from cortical tissue taken from post-natal day 1–3 mice and plated at approximately 3 hemispheres per plate ([Choi et al., 1987](#), [Schwartz and Wilson, 1992](#), [Rush et al., 2009](#)). Cultures were maintained in humidified 5% CO₂ incubators at 37 °C. Mice were handled in accordance with a protocol approved by our institutional [animal](#) care committee and in compliance with the Public [Health Service](#) Policy on Humane Care and Use of [Laboratory Animals](#). All experiments were performed in media identical to growth media except lacking serum (MS) at a volume of 400 μl per well. All experiments were performed on mixed cortical cultures, except where noted.

2.3. Assay of neuronal death

[Cell death](#) was assessed in cultures by the measurement of [lactate dehydrogenase](#) (LDH), released from damaged or destroyed cells, in the extracellular fluid 24 h after the beginning of the insult. Insults were performed by washing cultures into MS or MS containing the indicated chemicals and incubating overnight. Control [LDH](#) levels were subtracted from insult [LDH](#) values, and results normalized to 100% [neuronal death](#) caused by 500 μ M [NMDA](#). Control experiments have shown previously that the [efflux](#) of LDH occurring from either necrotic or apoptotic cells is proportional to the number of cells damaged or destroyed ([Koh and Choi, 1987](#), [Lobner, 2000](#)). [Trypan blue](#) staining indicated that the cell [death](#) observed was [neuron](#) selective. Cell-free experiments were performed to exclude the possibility of direct inhibition of LDH by MeHg. That is, MeHg at concentrations up to 10 μ M do not alter the quantification of an LDH standard.

2.4. Glutathione assay

Total cellular [glutathione](#) and media [glutathione](#) were assayed using a modified [enzymatic method](#) ([Baker et al., 1990](#), [Lobner et al., 2003](#)). Cultures were washed into MS with or without MeHg (5 μ M) and incubated for 6 h. Media samples of 25 μ l were taken after the [drug](#) application period and assayed as aliquots of supernatant below. Cultures were washed with cold (4 $^{\circ}$ C) [HEPES](#) buffered [saline](#) solution, dissolved in 200 μ l of 1% sulfosalicylic [acid](#), and centrifuged. A 25 μ l aliquot of the supernatant was combined with 150 μ l of 0.1 M phosphate/5 mM [EDTA buffer](#), 10 μ l of 20 mM dithiobis-2-nitrobenzoic [acid](#), 100 μ l of 5 mM [nicotinamide adenine dinucleotide phosphate](#) (NADPH), and 0.2 U of [glutathione reductase](#). Total [glutathione](#) was determined by kinetic analysis of [absorbance](#) changes at 402 nm for 1.5 min, with concentrations determined by comparison to a standard curve. GSSG was measured as above except samples were treated with 2-vinylpyridine and [triethanolamine](#) for 1 h prior to beginning the reaction. Cell-free experiments were performed to exclude the possibility of assay inhibition by the experimental [reagents](#) used. Due to inter-plate culture variance in absolute glutathione measures data in [Table 1](#) are reported as percent respective intra-plate control glutathione measures. Untreated control values across all culture types were typically in the following ranges (represented as mean nmol/well): 1.99–3.33 cellular GSH, 0.24–0.50 media GSH, 0.14–0.74 cellular GSSG, 0.060–0.16 media GSSG.

Table 1. Effect of MeHg (5 μ M) on [glutathione](#) levels in glial, neuronal and mixed cortical cultures after 6 h exposure.

| Culture type | GSH | | GSSG | |
|--------------|----------------|----------------|----------------|----------------|
| | Cellular | Media | Cellular | Media |
| Mixed | 83.4 \pm 4.7 | 22.8 \pm 1.0 | 60.7 \pm 5.7 | 12.6 \pm 2.4 |
| Glia | 44.8 \pm 5.2 | 62.3 \pm 9.8 | 18.4 \pm 2.9 | 29.3 \pm 3.7 |
| Neurons | 59.6 \pm 6.5 | 36.9 \pm 1.8 | 20.6 \pm 6.2 | N/D |

Results are expressed as % untreated control (mean \pm [SEM](#), $n = 8$ –16). All values are significantly different from control. N/D = not detectable.

2.5. Assay for oxidative stress

[Oxidative stress](#) was measured with 5-(and -6)-2',7'-dichlorodihydrofluorescein [diacetate](#) (DCF-DA) using a fluorescent plate [reader](#) following a modification of a previous method ([Wang and Joseph, 1999](#), [Lobner et al., 2007](#)). 10 μ M of the non-fluorescent, lipophilic compound DCF-DA is added to the cultures for 30 min where it is de-esterified and may be oxidized to fluorescent DCF (2',7'-dichlorofluorescein). [Fluorescence](#) is read using a Fluoroskan [Ascent](#) plate reader (Thermo LabSystems) with [excitation](#) and emission filters set to 485 nm and 538 nm, respectively. Background fluorescence (no DCF added) was subtracted and the results normalized to control conditions.

2.6. 14 C-cystine uptake

Radiolabeled [cystine](#) uptake was performed as previously described with modifications ([Liu et al., 2009](#)). Cultures were exposed to MS containing the indicated drug treatments for 24 h, or for 1 h followed by 23 h in

drug-free MS. 24 h after the start of the experiment, cultures were washed into [HEPES](#) buffered saline solution and immediately exposed to ^{14}C -cystine (0.025 $\mu\text{Ci}/\text{mL}$, 200 nM total cystine) for 20 min. Following ^{14}C -cystine exposure, cultures were washed with [ice](#) cold HEPES buffered saline solution and dissolved in 250 μl warm [sodium dodecyl sulfate](#) (0.1%). An aliquot (200 μl) was removed and added to [scintillation](#) fluid for counting. Values were normalized to ^{14}C -cystine uptake in untreated controls of the same experimental plate.

2.7. Determination of MeHg content in cells by ICP-MS

Following 6 h exposure to MS with or without the indicated treatments, cultures were washed 3 \times with cold MS and then dissolved in 100 μl 1% sulfosalicylic acid. Samples were combined with 900 μl 5% HNO_3 containing 500 ppb Au and digested at 70 $^\circ\text{C}$ for 2 h. [Digestions](#) were then centrifuged at 6000 $\times g$ for 5 min and the supernatants were diluted with an additional 1 mL of 5% HNO_3 . [Mercury](#) content was determined using a Micromass Platform ICP-MS controlled by MassLynx software (Waters Corporation, Milford, MA). [Isotopes](#) ^{198}Hg , ^{199}Hg , ^{200}Hg , ^{201}Hg , ^{202}Hg , and ^{204}Hg were recorded and total mercury was quantified by comparing sample responses to those produced by commercial standards treated identically to the samples.

2.8. Statistical analysis

Differences between test groups were examined for statistical significance by means of one-way ANOVA followed by the Student–Neuman Keuls post hoc analysis, with $p < 0.05$ being considered significant.

3. Results

[Methylmercury](#) (0.3–10 μM) exposure induced concentration-dependent [neurotoxicity](#) in mixed neuronal and glial cortical cultures 24 h after onset of insult ([Fig. 1](#)). [Cell death](#) was assayed by release of the cytosolic [enzyme LDH](#) 24 h after the beginning of the insult. Further, [trypan blue](#) staining of these cultures demonstrated that the MeHg-induced [toxicity](#) at the concentrations tested was purely neuronal (data not shown). From these data we chose 5 μM MeHg as a moderate insult for subsequent experiments as it allows bidirectional alterations to MeHg-induced [neurotoxicity](#) (i.e. [neuroprotection](#) or potentiation). For other experiments we used 3 μM MeHg as a low-level insult as no significant [neurotoxicity](#) was observed at this concentration ([Fig. 1](#)).




Fig. 1. Concentration response curve of MeHg (24 h exposure) on [LDH](#) release in primary cortical culture. Results are expressed as mean + [SEM](#) ($n = 8$ –16). *Significant difference from untreated control.

MeHg [toxicity](#) has been shown to cause [glutathione depletion](#) in several cell types ([Kitahara et al., 1993](#), [Kaur et al., 2006](#), [Amonpatumrat et al., 2008](#)). We found that exposure of mixed neural and glial, pure neuronal and pure glial cultures to 5 μM MeHg significantly decreased total GSH and [disulfide glutathione](#) (GSSG) levels in the cells as well as in the extracellular media ([Table 1](#)). This [glutathione](#) depletion occurred following a 6 h MeHg exposure, a time that precedes cell [death](#) as no [LDH](#) activity was detectable at this time (data not shown). We next tested whether [glutathione](#) supplementation or the addition of the [free radical scavenger, trolox](#) could protect the [neurons](#). 24 h co-treatment of MeHg with glutathione

monoethyl [ester](#) (GSHME) but not trolox, was [neuroprotective](#) ([Fig. 2](#)). Since glutathione supplementation was protective, and this effect could be due to its nature as an [antioxidant](#) or a [xenobiotic](#) detoxicant, we next assessed the abilities of GSHME and Trolox to block MeHg-induced [oxidative stress](#).




Fig. 2. MeHg-induced [neuronal death](#) is attenuated by [glutathione](#) monoethyl [ester](#) (GSHME, 100 μM) but not [trolox](#) (100 μM). Results are expressed as mean + [SEM](#) ($n = 16-20$). *Significantly different from MeHg-alone.

Mixed neuronal and glial cortical cultures exhibited a biphasic pattern of [oxidative stress](#) induced by exposure to 5 μM MeHg for 0, 30, 60, 180 or 360 min ([Fig. 3](#)) as measured by [fluorescence](#) of DCF. DCF [fluorescence](#) peaked at 30 min, returned to control levels at 3 h and increased once again by 6 h following onset of MeHg insult. Both 100 μM GSHME and 100 μM trolox were capable of preventing the MeHg-induced rise in DCF fluorescence indicative of their ability to block MeHg-induced oxidative stress. However, trolox was much more effective than GSHME actually decreasing the DCF signal well below control levels.




Fig. 3. Timecourse of MeHg (5 μM) induced ROS formation as detected by DCF [fluorescence](#). Results are expressed as mean + [SEM](#) ($n = 8-16$) after normalizing to average [fluorescence](#) of untreated controls (i.e. DCF only). *Significantly different from untreated controls.

To further examine MeHg induced alterations to glutathione-related systems, we next tested whether low-level MeHg exposure had an effect on cellular uptake of [cystine](#), the rate limiting substrate for GSH synthesis. Acute exposure to MeHg had no effect on ^{14}C -cystine uptake (3 μM MeHg, 20 min exposure, data not shown); however, 24-h exposure to low-level MeHg (3 μM) caused a robust increase in subsequent ^{14}C -cystine uptake that was blocked by co-treatment with the [protein synthesis inhibitor](#), [cycloheximide](#) (CHX, 500 ng/mL) ([Fig. 4A](#)). To determine whether the MeHg-induced increase in [cystine](#) uptake was mediated by cystine/glutamate exchange, we tested the effects of an inhibitor of this system, CPG (300 μM, present only

during the 20 min uptake period). The finding that CPG blocked the MeHg-induced increase in cystine uptake indicates that the increase was entirely mediated by [functional](#) upregulation of system x_c^- . When attempting to determine the effect of trolox on the MeHg-induced increase, we found that overnight exposure to trolox alone induced a large increase in subsequent cystine uptake (data not shown). To avoid this [complication](#) we used an altered protocol shortening the exposure time to the trolox and/or MeHg to 1 h, followed by a 23 h [incubation](#) in identical exposure media except lacking the trolox and MeHg. 60 min exposure to MeHg (3 μ M) produced a similar increase in xCT-mediated cystine uptake 23 h later ([Fig. 4B](#)). Co-treatment with trolox during this period blocked this increase while trolox alone for 60 min had no effect on uptake.




Fig. 4. Subtoxic MeHg exposure causes a robust increase in system x_c^- mediated 14 C-cystine uptake. (A) Cultures were exposed to MeHg (3 μ M), [cycloheximide](#) (CHX, 500 ng/mL), or in combination for 24 h prior to measuring uptake. Carboxyphenylglycine (CPG, 300 μ M) was present only during the 20 min uptake period. (B) Cultures were exposed to MeHg (3 μ M), [trolox](#) (100 μ M) or in combination for 60 min, washed into drug-free media and incubated overnight (23 h) prior to assay for [cystine](#) uptake. Results are expressed as mean + [SEM](#) ($n = 8-16$) after normalizing to untreated control uptake. *Significantly different from control.

It is possible that glutathione [conjugates](#) to MeHg intracellularly and that this conjugate is exported via [MRP1](#). To test whether the export of glutathione is necessary for its protective capabilities, we co-treated cultures with MeHg and an inhibitor of [MRP1](#), MK571. MK571 alone caused no significant [cell toxicity](#); however, by inhibiting MRP1-mediated glutathione export MeHg neurotoxicity was severely potentiated ([Fig. 5A](#)). Supporting the idea that MeHg conjugates to GSH, we found that inhibiting MRP1-mediated glutathione export with MK571 led to a doubling of MeHg accumulation within the cellular [compartment](#) ([Fig. 5B](#)). Supplementation with GSHME during the MeHg treatment had no effect on [mercury](#) accumulation in the cells (data not shown).




Fig. 5. Blockade of [glutathione efflux](#) augments MeHg [toxicity](#) and cellular accumulation of [mercury](#). (A) Cultures were exposed to MeHg (5 μ M), MK571 (10 μ M), or both in combination with or without GSHME (100 μ M) for 24 h at which time samples were taken and [neuronal death](#) quantified by [LDH](#) release. Results are expressed as mean + [SEM](#) ($n = 8-12$). *Significantly different from untreated control, #significantly different from MeHg-only treated. (B) Cultures were exposed to MeHg (5 μ M) in the presence of GSHME (100 μ M) or MK571 (50 μ M) for 6 h. Samples were then taken and subsequently analyzed for mercury content by ICP-MS. Results are expressed as mean + SEM ($n = 3-6$).

4. Discussion

The present study offers several insights into the contributions of alterations in [glutathione](#) cycling and [oxidative stress](#) to methylmercury-induced [neurotoxicity](#). We observed significant [toxicity](#) with overnight exposure of mixed cortical cultures to 5 μ M MeHg. This concentration is similar to those used in other studies demonstrating [cytotoxicity](#), [oxidative stress](#) and mitochondrial deficits in cortical and cerebellar granular [neuron](#) (CGN) cultures ([Gasso et al., 2001](#), [Morken et al., 2005](#), [Kaur et al., 2006](#), [Yin et al., 2007](#)). However, it is important to note that CGNs have been shown to be more sensitive to MeHg [toxicity](#) ([Sarafian and Verity, 1991](#), [Marty and Atchison, 1997](#), [Sakaue et al., 2005](#)). Though not the only factor recognized, the increased sensitivity of CGNs has been posited as attributable to the relatively low [glutathione](#) content of these cells ([Yee and Choi, 1996](#), [Shafer et al., 2002](#), [Kaur et al., 2007](#), [Wang et al., 2009](#)).

Previous studies have shown that MeHg causes a [depletion](#) of cellular [glutathione](#) ([Yee and Choi, 1996](#), [Franco et al., 2007](#), [Amonpatumrat et al., 2008](#), [Wang et al., 2009](#)). Here, we also report that MeHg depletes both total and [disulfide glutathione](#) in cortical culture; total GSH and [disulfide](#) GSSG were decreased in cellular and media samples from glia-enriched, neuron-enriched and mixed cultures. GSH can reduce [free radicals](#) or eliminate exogenous [molecules](#) from the cell. The MeHg-induced loss of glutathione may be due to the formation of ROS and subsequent use of GSH by the cells to reduce the oxidative damage. However, since

reduction of oxidative stress by GSH results in the formation of GSSG, the observed depletion of both GSH and GSSG suggests that the loss of glutathione was not due to oxidative stress. This data is consistent with the [hypothesis](#) that glutathione is acting to detoxify MeHg by direct [conjugation](#) which may render the utilized GSH molecules undetectable by our assay, and perhaps unrecoverable by the cell.

It has been shown that [modulation](#) of cellular [thiols](#) (including GSH) alters sensitivity to MeHg in culture ([Kaur et al., 2006](#)) and that exogenous supply of glutathione precursors such as *N*-acetyl-cysteine (NAC) or [cystine](#) is [neuroprotective](#) ([Fujiyama et al., 1994](#), [Kaur et al., 2006](#), [Kaur et al., 2007](#)). Specifically, NAC is able to boost cellular GSH availability and leads to decreased MeHg-induced [cytotoxicity](#) and ROS formation. Inhibition of glutathione synthesis with buthionine sulfoxamine (BSO) increases sensitivity to MeHg and increases subsequent ROS formation ([Toyama et al., 2011](#)). These studies also demonstrate that modulation of glutathione availability alters cellular ROS formation in response to MeHg insult. Further, this ROS formation is sensitive to [antioxidants](#) ([Gasso et al., 2001](#), [Shanker and Aschner, 2003](#), [Kaur et al., 2010](#)). Importantly, however, in our hands [trolox](#) does not offer the neuroprotective capacity that glutathione supplementation with GSHME provides. This suggests that oxidative stress alone is not primary to the toxicity of MeHg under these conditions. Others have shown that trolox does offer [neuroprotection](#) against early MeHg-induced mitochondrial deficits detected by MTT [metabolism](#) after short, 1 h exposure ([Kaur et al., 2010](#)). Notably, however, these studies did not investigate later timepoints as seen in the present data.

An important action of MeHg is the inhibition of [selenoproteins](#). Among the [selenoproteins](#) that MeHg has been shown to inhibit are [thioredoxin reductase](#) and thioredoxin-glutathione [peroxidase](#) ([Carvalho et al., 2008](#), [Reeves and Hoffmann, 2009](#), [Branco et al., 2011](#)). In fact, the [IC50](#) for the inhibition of [thioredoxin reductase 1](#) is 9 nM ([Carvalho et al., 2008](#)); therefore, at the concentrations of MeHg used in the current study this [enzyme](#), and similar selenoproteins, are likely greatly inhibited. Therefore, it is clear that interaction with selenoproteins could be a mechanism of MeHg induced GSH depletion and oxidative stress. However, while preincubation with [selenomethionine](#) has been shown to attenuate MeHg induced production of [reactive oxygen species](#), it did not prevent depletion of GSH ([Kaur et al., 2009](#)). Although, dietary [selenium](#) is known to be protective against MeHg exposure ([Ralston and Raymond, 2010](#)).

MeHg can also inhibit other enzymes important for the cells response to oxidative stress, such as thioredoxin and [glutathione reductase](#) ([Carvalho et al., 2008](#)). However, the inhibition of [glutathione reductase](#) was found to only occur [in vitro](#) and not in cells treated with MeHg ([Carvalho et al., 2008](#)).

We also examined the function of the cystine–glutamate [antiporter](#) in response to MeHg exposure by radiolabeled ¹⁴C-cystine uptake. As discussed above, [cystine](#) uptake is an important aspect of glutathione synthesis and subsequent cycling. Low-level MeHg insult induced a robust increase in cystine uptake. This [functional](#) upregulation of xCT was blocked by CHX suggesting that the upregulation requires translation of new [protein](#). As well, CPG blocked this increased [radioligand](#) uptake to CPG-treated control levels suggesting the entire MeHg-induced increase was mediated by system x_c^- . Trolox also blocked the MeHg-increased cystine uptake. Considering that upregulation of xCT is likely an endogenous mechanism of [cytoprotection](#) in response to MeHg-insult, the fact that trolox blocks upregulation of xCT may account for the inability of trolox to protect against the neuronal [injury](#). The most likely mechanism for increased system x_c^- activity is upregulation of xCT by [activation](#) of the nrf2-ARE pathway ([Mysona et al., 2009](#), [Wang et al., 2009](#), [Qin et al., 2010](#)). Consistent with this explanation is a recent report demonstrating nrf2-dependent [methylmercury detoxification](#) in hepatocytes ([Toyama et al., 2011](#)) and an earlier report in [astrocytes](#) ([Wang et al., 2009](#)).

The current data indicate that blockade of oxidative stress alone is insufficient to prevent MeHg-induced [neuronal death](#), and that [elimination](#) of MeHg from the cell by conjugation to GSH is necessary. Consistent with this, inhibition of glutathione export via the [MRP1 inhibitor](#), MK571, not only potentiated MeHg-induced [neurotoxicity](#), but this potentiation was associated with increased accumulation of MeHg in the cell. Similar results have been reported in primary [mouse](#) hepatocytes where it was found that upregulation of [MRP1](#) by [isothiocyanates](#) decreased MeHg accumulation and toxicity ([Toyama et al., 2011](#)). Interestingly, and consistent with a previous report using NAC ([Kaur et al., 2006](#)), glutathione supplementation with GSHME did not affect MeHg accumulation in the cell. Trolox has previously been reported to have no effect on MeHg accumulation nor on glutathione availability ([Kaur et al., 2010](#)); these and the present data, support the

assertion that oxidative stress is not primary to MeHg-induced [neurotoxicity](#). Rather, conjugation with MeHg and subsequent export of glutathione from the cytosol is crucial to alleviate or prevent MeHg-induced neurotoxicity and that solely [blocking](#) oxidative stress is not likely to offer significant [neuroprotection](#).

Conflict of interest statement

The authors declare that there are no [conflicts of interest](#).

References

- [Ali et al., 1992](#) S.F. Ali, C.P. LeBel, S.C. Bondy **Reactive oxygen species formation as a biomarker of methylmercury and trimethyltin neurotoxicity** *Neurotoxicology*, 13 (1992), pp. 637-648
- [Amonpatumrat et al., 2008](#)
S. Amonpatumrat, H. Sakurai, P. Wiriyasermkul, N.Khunweeraphong, S. Nagamori, H. Tanaka, *et al.* **I-glutamate enhances methylmercury toxicity by synergistically increasing oxidative stress** *J Pharmacol Sci*, 108 (2008), pp. 280-289
- [Aschner, 2000](#) M. Aschner **Astrocytic swelling, phospholipase A2, glutathione and glutamate: interactions in methylmercury-induced neurotoxicity** *Cell Mol Biol (Noisy-le-grand)*, 46 (2000), pp. 843-854
- [Atchison and Hare, 1994](#) W.D. Atchison, M.F. Hare **Mechanisms of methylmercury-induced neurotoxicity** *FASEB J*, 8 (1994), pp. 622-629
- [Baker et al., 1990](#) M.A. Baker, G.J. Cerniglia, A. Zaman **Microtiter plate assay for the measurement of glutathione and glutathione disulfide in large numbers of biological samples** *Anal Biochem*, 190 (1990), pp. 360-365
- [Branco et al., 2011](#) V. Branco, J. Canario, J. Lu, A. Holmgren, C. Carvalho **Mercury and selenium interaction in vivo: effects on thioredoxin reductase and glutathione peroxidase** *Free Radic Biol Med* (2011) [Epub ahead of print]
- [Bridges and Zalups, 2005](#) C.C. Bridges, R.K. Zalups **Molecular and ionic mimicry and the transport of toxic metals** *Toxicol Appl Pharmacol*, 204 (2005), pp. 274-308
- [Bridges and Zalups, 2010](#) C.C. Bridges, R.K. Zalups **Transport of inorganic mercury and methylmercury in target tissues and organs** *J Toxicol Environ Health B Crit Rev*, 13 (2010), pp. 385-410
- [Carvalho et al., 2008](#) C.M.L. Carvalho, E.-H. Chew, S.I. Hashemy, J. Lu, A. Holmgren **Inhibition of the human thioredoxin system. A molecular mechanism of mercury toxicity** *J Biol Chem*, 283 (2008), pp. 11913-11923
- [Choi et al., 1987](#) D.W. Choi, M. Maulucci-Gedde, A.R. Kriegstein **Glutamate neurotoxicity in cortical cell culture** *J Neurosci*, 7 (1987), pp. 357-368
- [Clarkson, 1997](#) T.W. Clarkson **The toxicology of mercury** *Crit Rev Clin Lab Sci*, 34 (1997), pp. 369-403
- [Dringen, 2000](#) R. Dringen **Metabolism and functions of glutathione in brain** *Prog Neurobiol*, 62 (2000), pp. 649-671
- [Dringen and Hirrlinger, 2003](#) R. Dringen, J. Hirrlinger **Glutathione pathways in the brain** *Biol Chem*, 384 (2003), pp. 505-516
- [Dringen et al., 1999](#) R. Dringen, B. Pfeiffer, B. Hamprecht **Synthesis of the antioxidant glutathione in neurons: supply by astrocytes of CysGly as precursor for neuronal glutathione** *J Neurosci*, 19 (1999), pp. 562-569
- [Dugan et al., 1995](#) L.L. Dugan, V.M. Bruno, S.M. Amagasu, R.G. Giffard **Glia modulate the response of murine cortical neurons to excitotoxicity: glia exacerbate AMPA neurotoxicity** *J Neurosci*, 15 (1995), pp. 4545-4555
- [Fonnum and Lock, 2000](#) F. Fonnum, E.A. Lock **Cerebellum as a target for toxic substances** *Toxicol Lett*, 15 (2000), pp. 9-16
- [Franco et al., 2007](#) J.L. Franco, H.C. Braga, J. Stringari, F.C. Missau, T. Posser, B.G. Mendes, *et al.* **Mercurial-induced hydrogen peroxide generation in mouse brain mitochondria: protective effects of quercetin** *Chem Res Toxicol*, 20 (2007), pp. 1919-1926
- [Fujiyama et al., 1994](#) J. Fujiyama, K. Hirayama, A. Yasutake **Mechanism of methylmercury efflux from cultured astrocytes** *Biochem Pharmacol*, 47 (1994), pp. 1525-1530

- [Gasso et al., 2001](#) S. Gasso, R.M. Cristofol, G. Selema, R. Rosa, E. Rodriguez-Farre, C.Sanfeliu **Antioxidant compounds and Ca(2+) pathway blockers differentially protect against methylmercury and mercuric chloride neurotoxicity** *J Neurosci Res*, 66 (2001), pp. 135-145
- [Harada, 1995](#) M. Harada **Minamata disease: methylmercury poisoning in Japan caused by environmental pollution** *Crit Rev Toxicol*, 25 (1995), pp. 1-24
- [Hirrlinger and Dringen, 2005](#) J. Hirrlinger, R. Dringen **Multidrug resistance protein 1-mediated export of glutathione and glutathione disulfide from brain astrocytes** *Methods Enzymol*, 400 (2005), pp. 395-409
- [Hock et al., 1998](#) C. Hock, G. Drasch, S. Golombowski, F. Muller-Spahn, B. Willershausen-Zonnchen, P. Schwarz, *et al.* **Increased blood mercury levels in patients with Alzheimer's disease** *J Neural Transm*, 105 (1998), pp. 59-68
- [Kaur et al., 2006](#) P. Kaur, M. Aschner, T. Syversen **Glutathione modulation influences methyl mercury induced neurotoxicity in primary cell cultures of neurons and astrocytes** *Neurotoxicology*, 27 (2006), pp. 492-500
- [Kaur et al., 2007](#) P. Kaur, M. Aschner, T. Syversen **Role of glutathione in determining the differential sensitivity between the cortical and cerebellar regions towards mercury-induced oxidative stress** *Toxicology*, 230 (2007), pp. 164-177
- [Kaur et al., 2009](#) P. Kaur, L. Evje, M. Aschner, T. Syversen **The in vitro effects of selenomethionine on methylmercury-induced neurotoxicity** *Toxicol In Vitro*, 23 (2009), pp. 378-385
- [Kaur et al., 2010](#) P. Kaur, L. Evje, M. Aschner, T. Syversen **The in vitro effects of Trolox on methylmercury-induced neurotoxicity** *Toxicology*, 276 (2010), pp. 73-78
- [Kawanai et al., 2009](#) T. Kawanai, M. Satoh, K. Murao, Y. Oyama **Methylmercury elicits intracellular Zn²⁺ release in rat thymocytes: its relation to methylmercury-induced decrease in cellular thiol content** *Toxicol Lett*, 191 (2009), pp. 231-235
- [Kitahara et al., 1993](#) J. Kitahara, Y. Seko, N. Imura **Possible involvement of active oxygen species in selenite toxicity in isolated rat hepatocytes** *Arch Toxicol*, 67 (1993), pp. 497-501
- [Koh and Choi, 1987](#) J.Y. Koh, D.W. Choi **Quantitative determination of glutamate mediated cortical neuronal injury in cell culture by lactate dehydrogenase efflux assay** *J Neurosci Methods*, 20 (1987), pp. 83-90
- [Kranich et al., 1998](#) O. Kranich, R. Dringen, M. Sandberg, B. Hamprecht **Utilization of cysteine and cysteine precursors for the synthesis of glutathione in astroglial cultures: preference for cystine** *Glia*, 22 (1998), pp. 11-18
- [Liu et al., 2009](#) X. Liu, T. Rush, J. Zapata, G.D. Lobner **Beta-N-methylamino-L-alanine induces oxidative stress and glutamate release through action on system Xc(-)** *Exp Neurol*, 217 (2009), pp. 429-433
- [Lobner, 2000](#) D. Lobner **Comparison of the LDH and MTT assays for quantifying cell death: validity for neuronal cell death** *J Neurosci Meth*, 96 (2000), pp. 147-152
- [Lobner et al., 2003](#) D. Lobner, S. Golner, J. Hjelmhaug **Neurotrophic factor effects on oxidative stress-induced neuronal death** *Neurochem Res*, 28 (2003), pp. 749-756
- [Lobner et al., 2007](#) D. Lobner, P.M. Piana, A.K. Salous, R.W. Peoples **Beta-N-methylamino-L-alanine enhances neurotoxicity through multiple mechanisms** *Neurobiol Dis*, 25 (2007), pp. 360-366
- [Mahaffey et al., 2004](#) K.R. Mahaffey, R.P. Clickner, C.C. Bodurow **Blood organic mercury and dietary mercury intake: National Health and Nutrition Examination Survey, 1999 and 2000** *Environ Health Perspect*, 112 (2004), pp. 562-570
- [Marty and Atchison, 1997](#) M.S. Marty, W.D. Atchison **Pathways mediating Ca²⁺ entry in rat cerebellar granule cells following in vitro exposure to methyl mercury** *Toxicol Appl Pharmacol*, 147 (1997), pp. 319-330
- [Minich et al., 2006](#) T. Minich, J. Riemer, J.B. Schulz, P. Wielinga, J. Wijnholds, R. Dringen **The multidrug resistance protein 1 (Mrp1), but not Mrp5, mediates export of glutathione and glutathione disulfide from brain astrocytes** *J Neurochem*, 97 (2006), pp. 373-384
- [Monnet-Tschudi et al., 2006](#) F. Monnet-Tschudi, M.G. Zurich, C. Boschat, A. Corbaz, P. Honegger **Involvement of environmental mercury and lead in the etiology of neurodegenerative diseases** *Rev Environ Health*, 21 (2006), pp. 105-117

- [Morken et al., 2005](#) T.S. Morken, U. Sonnewald, M. Aschner, T. Syversen **Effects of methylmercury on primary brain cells in mono- and co-culture** *Toxicol Sci*, 87 (2005), pp. 169-175
- [Myers et al., 2000](#) G.J. Myers, P.W. Davidson, C. Cox, C. Shamlaye, E. Cernichiari, T.W. Clarkson **Twenty-seven years studying the human neurotoxicity of methylmercury exposure** *Environ Res*, 83 (2000), pp. 275-285
- [Mysona et al., 2009](#) B. Mysona, Y. Dun, J. Duplantier, V. Ganapathy, S.B. Smith **Effects of hyperglycemia and oxidative stress on the glutamate transporters GLAST and system x_c⁻ in mouse retinal Muller glial cells** *Cell Tissue Res*, 335 (2009), pp. 477-488
- [Praline et al., 2007](#) J. Praline, A.M. Guennoc, N. Limousin, H. Hallak, B. de Toffol, P. Corcia **ALS and mercury intoxication: a relationship?** *Clin Neurol Neurosurg*, 109 (2007), pp. 880-883
- [Qin et al., 2010](#) Z. Qin, E. Freitas, R. Sullivan, S. Mohan, R. Bacelieri, D. Branch, *et al.* **Upregulation of xCT by KSHV-encoded microRNAs facilitates KSHV dissemination and persistence in an environment of oxidative stress** *PLoS Pathog*, 6 (2010), p. e1000742
- [Ralston and Raymond, 2010](#) N.V. Ralston, L.J. Raymond **Dietary selenium's protective effects against methylmercury toxicity** *Toxicology*, 278 (2010), pp. 112-123
- [Reeves and Hoffmann, 2009](#) M.A. Reeves, P.R. Hoffmann **The human selenoproteome: recent insights into functions and regulation** *Cell Mol Life Sci*, 66 (2009), pp. 2457-2478
- [Rooney, 2007](#) J.P. Rooney **The role of thiols, dithiols, nutritional factors and interacting ligands in the toxicology of mercury** *Toxicology*, 234 (2007), pp. 145-156
- [Rush et al., 2009](#) T. Rush, J. Hjelmhaug, D. Lobner **Effects of chelators on mercury, iron, and lead neurotoxicity in cortical culture** *Neurotoxicology*, 30 (2009), pp. 47-51
- [Rush et al., 2010](#) T. Rush, X.Q. Liu, J. Hjelmhaug, D. Lobner **Mechanisms of chlorpyrifos and diazinon induced neurotoxicity in cortical culture** *Neuroscience*, 166 (2010), pp. 899-906
- [Sakaue et al., 2005](#) M. Sakaue, M. Okazaki, S. Hara **Very low levels of methylmercury induce cell death of cultured rat cerebellar neurons via calpain activation** *Toxicology*, 213 (2005), pp. 97-106
- [Salinas and Wong, 1999](#) A.E. Salinas, M.G. Wong **Glutathione S-transferases – a review** *Curr Med Chem*, 6 (1999), pp. 279-309
- [Sarafian and Verity, 1991](#) T. Sarafian, M.A. Verity **Oxidative mechanisms underlying methyl mercury neurotoxicity** *Int J Dev Neurosci*, 9 (1991), pp. 147-153
- [Schwartz and Wilson, 1992](#) J.P. Schwartz, D.J. Wilson **Preparation and characterization of type 1 astrocytes cultured from adult rat cortex, cerebellum, and striatum** *Glia*, 5 (1992), pp. 75-80
- [Shafer et al., 2002](#) T.J. Shafer, C.A. Meacham, S. Barone Jr. **Effects of prolonged exposure to nanomolar concentrations of methylmercury on voltage-sensitive sodium and calcium currents in PC12 cells** *Brain Res Dev Brain Res*, 136 (2002), pp. 151-164
- [Shanker and Aschner, 2003](#) G. Shanker, M. Aschner **Methylmercury-induced reactive oxygen species formation in neonatal cerebral astrocytic cultures is attenuated by antioxidants** *Brain Res Mol Brain Res*, 110 (2003), pp. 85-91
- [Toyama et al., 2011](#) T. Toyama, Y. Shinkai, A. Yasutake, K. Uchida, M. Yamamoto, Y. Kumagai **Isothiocyanates reduce mercury accumulation via an Nrf2-dependent mechanism during exposure of mice to methylmercury** *Environ Health Perspect* (2011)
- [Tsuda et al., 2009](#) T. Tsuda, T. Yorifuji, S. Takao, M. Miyai, A. Babazono **Minamata disease: catastrophic poisoning due to a failed public health response** *J Public Health Policy*, 30 (2009), pp. 54-67
- [Wang and Joseph, 1999](#) H. Wang, J.A. Joseph **Quantifying cellular oxidative stress by dichlorofluorescein assay using microplate reader** *Free Radic Biol Med*, 27 (1999), pp. 612-616
- [Wang et al., 2009](#) L. Wang, H. Jiang, Z. Yin, M. Aschner, J. Cai **Methylmercury toxicity and Nrf2-dependent detoxification in astrocytes** *Toxicol Sci*, 107 (2009), pp. 135-143
- [Yee and Choi, 1996](#) S. Yee, B.H. Choi **Oxidative stress in neurotoxic effects of methylmercury poisoning** *Neurotoxicology*, 17 (1996), pp. 17-26

[Yin et al., 2007](#) Z. Yin, D. Milatovic, J.L. Aschner, T. Syversen, J.B. Rocha, D.O. Souza, *et al.* **Methylmercury induces oxidative injury, alterations in permeability and glutamine transport in cultured astrocytes** *Brain Res*, 1131 (2007), pp. 1-10