

4-1-2001

Increased mRNA Expression for the α_1 Subunit of the GABA_A Receptor Following Nitrous Oxide Exposure in Mice

Lisa M. Johaneck
Marquette University

William E. Cullinan
Marquette University, william.cullinan@marquette.edu

Linda K. Vaughn
Marquette University, linda.vaughn@marquette.edu

Marquette University

e-Publications@Marquette

Biomedical Sciences Faculty Research and Publications/College of Health Sciences

This paper is NOT THE PUBLISHED VERSION; but the author's final, peer-reviewed manuscript. The published version may be accessed by following the link in the citation below.

Molecular Brain Research. Vol. 89, No. 1-2 (April 2001): 41-49. [DOI](#). This article is © Elsevier and permission has been granted for this version to appear in [e-Publications@Marquette](#). Elsevier does not grant permission for this article to be further copied/distributed or hosted elsewhere without the express permission from Elsevier.

Increased mRNA expression for the α_1 subunit of the GABA_A receptor following nitrous oxide exposure in mice

Lisa M. Johanek

Department of Biomedical Sciences, Marquette University, PO Box 1881, Milwaukee WI 53201-1881, USA

William E. Cullinan

Department of Biomedical Sciences, Marquette University, PO Box 1881, Milwaukee WI 53201-1881, USA

Linda K. Vaughn

Department of Biomedical Sciences, Marquette University, PO Box 1881, Milwaukee WI 53201-1881, USA

Abstract

The mechanisms by which nitrous oxide (N₂O) produces physical dependence and withdrawal seizures are not well understood, but both N₂O and ethanol exert some of their effects via the GABA_A receptor and several lines of evidence indicate that withdrawal from N₂O and ethanol may be produced through similar mechanisms. Expression levels of mRNA transcripts encoding several GABA_A receptor subunits change with chronic ethanol exposure and, therefore, we hypothesized that N₂O exposure would produce changes in mRNA expression for the α_1 subunit. Male, Swiss–Webster mice, 10–12 weeks of age, were exposed for 48 h to either room air or a 75%:25% N₂O:O₂ environment. Brains were sectioned and mRNA for the α_1 subunit was detected by in situ

hybridization using an ³⁵S-labelled cRNA probe. N₂O exposure produced a significant increase in expression levels of the α₁ subunit mRNA in the cingulate cortex, the CA1/2 region of the hippocampus, the dentate gyrus, the subiculum, the medial septum, and the ventral tegmental area. These results lend support to the hypothesis that N₂O effects are produced, at least in part, through the GABA_A receptor and that N₂O produces these effects through actions in the cingulate cortex, hippocampus, ventral tegmental area and medial septum. These results are also further evidence that ethanol and N₂O produce dependence and withdrawal through common mechanisms.

Neurotransmitters, modulators, transporters, and receptors, GABA receptors

Keywords

Nitrous oxide, GABA_A receptor, Alpha₁ subunit, Mouse, In situ hybridization

1. Introduction

Nitrous oxide (N₂O) is an analgesic, anxiolytic, anesthetic drug with euphorogenic properties and abuse potential. Animals with prolonged exposure to N₂O experience physical dependence and withdrawal seizures [3], [18], [31], [36], [50]. Although it is widely believed that nitrous oxide interacts with central opioid mechanisms to produce its analgesic effect (for review see [48]) and through the GABA_A receptor to produce its anxiolytic effect [13], [45], [49], the mechanisms by which N₂O produces physical dependence and withdrawal seizures remain unclear.

Several lines of evidence indicate that withdrawal from N₂O and ethanol may be produced through similar mechanisms. Withdrawal from N₂O and ethanol are both accompanied by withdrawal seizures [3], [18], [31], [36], [50], [53]. N₂O can suppress ethanol withdrawal seizures [3], ethanol can suppress N₂O withdrawal seizures [3], and cross-tolerance between N₂O and ethanol has been demonstrated [22]. In addition, sensitivity of various mouse strains to ethanol and N₂O correlate well [3], [21], and quantitative trait loci analysis of BXD recombinant inbred mouse strains demonstrates a genetic correlation between N₂O and ethanol withdrawal [4].

Some of the effects of both N₂O and ethanol, including anxiolytic effects and withdrawal symptoms, are known to occur through interaction with the GABA_A receptor. The anxiolytic effects of N₂O are sensitive to antagonism by the benzodiazepine antagonist flumazenil and the anxiolytic effects are significantly reduced in benzodiazepine-tolerant animals [13], [45]. There is also evidence that withdrawal responses following exposure to N₂O involve the GABA_A receptor. For example, injection of the benzodiazepine partial inverse agonist Ro 15-4513 significantly increased the frequency of withdrawal seizures in N₂O-exposed mice [56], an effect also seen with ethanol-induced withdrawal seizures [2], [27].

GABA_A receptors are heterooligomeric complexes consisting of numerous isoforms of several classes of subunits (α, β, γ, δ, ρ). Since recombinant expression studies have shown that the pharmacology and the binding characteristics of GABA_A receptor are altered by the expression of different subunit combinations [1], [7], [10], [25], [28], [37], [38], [44], [51], [52], it has been hypothesized that one mechanism by which ethanol alters GABA_A receptor function is by differentially affecting the expression of GABA_A receptor subunits and thus altering the composition of GABA_A receptors and their responsiveness to GABA [34], [39], [41]. Expression levels of mRNA transcripts encoding several GABA_A receptor subunits are altered by chronic ethanol exposure, with consistent findings of decreased mRNA expression for the α₁ subunit in rat cortex following chronic ethanol exposure [8], [12], [32], [33], [34], [39], [41]. A number of studies have detected changes in other subunits of the GABA_A receptor in either rat cortex and cerebellum following chronic ethanol, including changes in α₂₋₆, [8], [11], [12], [33], [34], [39], β₁₋₃ [11], [35], and γ₁ [11], [12], with others reporting no differences in expression of α₃ [34], [39], α₅ [12] or β₁₋₃ [12], [40]. However, relatively few studies have examined brain regions other than cortex or cerebellum. These studies have found that durations longer than the typical 2 weeks of ethanol exposures are

necessary to produce changes in GABA_A receptor subunit composition. Forty days of chronic ethanol consumption were necessary to produce an increase in the α_4 subunit peptide in hippocampus [32]. Gene expression for the hippocampal α_1 , α_2 , α_3 , $\beta_{2/3}$ or γ_2 subunits was unchanged at that time. Others found that a 12-week, but not 4-week, ethanol exposure induced changes in α_1 and α_5 subunit expression in the hippocampus and α_1 subunit levels in the ventral tegmental area [8], [42]. No changes were found in nucleus accumbens or substantia nigra [8].

Results of ethanol exposure on GABA_A receptor subunit composition in mouse brain have been mixed, with results largely dependent upon mouse strain and brain region. For instance, following ethanol treatment, expression of whole brain α_1 mRNA was found to be decreased in Withdrawal Seizure Prone (WSP) and unchanged in Withdrawal Seizure Resistant (WSR) mice [6] but increased in ddy mice [19]. On the other hand, α_1 mRNA expression was reportedly increased in the cerebellum but decreased in the whole brains of C57BL/6J (C6) mice [60]. Levels of α_6 mRNA were decreased in WSR mice, unchanged in WSP mice [6], and increased in C6 mice [60]. Levels of γ_2 mRNA were increased by ethanol in C6 mice [60] but not WSP or WSR mice [6].

Because both N₂O and ethanol exert some of their effects via the GABA_A receptor, and evidence indicates that ethanol and N₂O produce dependence and withdrawal through similar mechanisms, we investigated the effect of N₂O exposure on mRNA expression for the α_1 subunit of the GABA_A receptor.

Further, we felt it important to conduct an extensive neuroanatomical study, both because multiple brain regions are thought to be important in the development of drug dependency and withdrawal responses, and because of the wide-ranging effects of N₂O on the brain. Previous work has shown that withdrawal seizures can be evoked following durations of N₂O exposures ranging from 30 min to 48 h [62]. For this neuroanatomical study of the distribution of N₂O-induced changes in mRNA expression, a 48-h duration was chosen to maximize opportunity for alterations mRNA expression levels.

2. Materials and methods

2.1. Nitrous oxide exposure

Male, Swiss–Webster mice (Charles River, Portage, MI), 10–12 weeks of age, were exposed for 48 h to either room air or a 75%:25% N₂O:O₂ environment. Three cages of five mice were placed in a modified infant incubator [26] with a N₂O delivery rate of 6 liter/min and an O₂ delivery rate of 1.5 liter/min. Gases were delivered using a standard nitrous oxide/oxygen anesthesia machine (Adec, Newburg, Oregon). The exhausted gas was vented to a nearby fumehood. Mice were removed from the N₂O, left in room air for 10 minutes and sacrificed by rapid decapitation. Whole brains were immediately removed, frozen in isopentane at –40°C for 30 s, wrapped in parafilm, and stored at –70°C

2.2. In situ hybridization

2.2.1. Tissue fixation

Serial cryostat sections of brain (14 μ m) were mounted on polylysine-coated slides and stored at –70°C until use. Slices were fixed in 4% buffered paraformaldehyde for 2 h and washed four times with 2 \times SSC (saline–sodium citrate buffer) (pH=7.4). Tissue sections were treated with proteinase K (0.1 μ g/ml) for 8 min at 37°C, washed with distilled water and rinsed with 2 \times SSC. Slides were subsequently acetylated with 0.25% acetic anhydride in a 0.1 M triethanolamine buffer (pH=8.0) for 15 minutes. Finally, slides were washed in 2 \times SSC three times and dehydrated in a successive series of alcohols.

2.2.2. Probe labeling

A riboprobe complementary to mRNA transcript encoding the α_1 subunit of the GABA_A receptor was synthesized according to standard in vitro transcription methodology. Briefly, a cRNA probe was generated in a reaction

(total volume=10 μ l) containing 125 μ Ci 35 S UTP (Amersham, 20 μ Ci/ml), 1 μ l each of 10 mM stocks of ATP, CTP, and GTP, 2 μ l 5 \times transcription buffer, 1 μ l 0.1 M dithiothreitol, 1 μ l linearized plasmid DNA (1 μ g/ μ l), 1 μ l distilled water, 0.5 μ l of RNase inhibitor (40 U/ μ l), and 1.5 μ l T7 polymerase. The reaction was allowed to proceed at 37°C for 2 h followed by separation of labeled probe from reaction contents on a G50-sephadex column. Probes were diluted in hybridization buffer (Amresco, Solon, OH) consisting of 45% formamide, 10% dextran sulfate, 3 \times SSC, 50 mM sodium phosphate buffer (pH=7.4), 1 \times Denhardt's solution, 0.1 mg/ml sheared salmon sperm DNA, and 0.1 mg/ml yeast, to yield approximately 1 500 000 cpm/35 μ l buffer. The GABA_A-receptor cDNA was kindly provided by Dr. A. Tobin (UCLA) and was subcloned into PGem3Z and linearized with Hind III to transcribe a cRNA probe of 842 nts.

2.2.3. Hybridization histochemistry

Tissue sections were apposed to coverslips containing 35 μ l probe and stored for 15 h at 55°C in sealed chambers moistened with a 45% formamide solution. Following hybridization, coverslips were removed in 2 \times SSC, the slides were washed extensively in 2 \times SSC and then treated with RNase A (200 μ g/ml) in Tris buffer for 90 minutes at 37°C. This was followed by several rinses in decreasing concentrations of SSC (2 \times SSC, 1 \times SSC, 0.5 \times SSC, and 0.2 \times SSC), and washing at 65°C for 1 h in 0.2 \times SSC. Slides were then washed in 0.2 \times SSC at room temperature and dehydrated through graded alcohol concentrations. Slides were exposed to Kodak BioMax X-ray film for 16 h and developed in GBX developer.

2.2.4. Imaging

X-ray images were digitized and analyzed using NIH Image software. Identification of brain regions was determined using the atlas of Franklin and Paxinos [14]. Depending on the size of the brain region of interest, between 3 and 80 gray value signals for each specific brain region were obtained per mouse ($n=5$ control mice, $n=8$ N₂O exposed mice). Mean gray values were determined for each brain region per mouse after subtraction of background values determined from adjacent brain regions which lacked specific expression of the GABA_A receptor.

2.3. Statistics

Two-way analysis of variance was performed to test whether there were differences in mean gray values in 20 brain regions between N₂O and room air exposed mice. To correct for heteroscedasticity, the data was transformed using the equation

[62]. Post-hoc analysis to determine which brain regions accounted for the statistically significant main effect seen in treatment groups was performed using the Tukey multiple comparison test.

3. Results

The regional expression pattern of the α_1 subunit mRNA was similar to that previously found in rat [29], [30], [43], [57] and long-sleep and short-sleep mouse brain [61]. Widespread distribution was noted, with highest expression densities in the cerebellar Purkinje layer and the ventral tegmental nucleus, high levels in the cortical primary and secondary somatosensory cortex, dense layer of the piriform cortex, CA1/2 region of the hippocampus and subiculum, granular layer of the cerebellum, red nucleus and the medial septum. Moderate to high levels were found in the cingulate and motor cortex, CA3 region of the hippocampus, dentate gyrus, molecular layer of the cerebellum, and globus pallidus. Low to moderate levels were found in the basolateral, medial and central amygdaloid nucleus, and ventral tegmental area (VTA) (Fig. 1). The regional distribution of the GABA_A receptor α_1 subunit mRNA in N₂O and room air exposed mice is shown in Fig. 2, Fig. 3.

Regional GABA_A α_1 Subunit Expression




Fig. 1. Regional distribution of expression of GABA_A α_1 subunit mRNA in control mice. Mean gray values, determined from multiple sections for each specific brain region, were determined for each mouse. The average mean gray values for 5 control mice \pm S.E.M. are plotted.




Fig. 2. Darkfield photomicrographs of coronal sections at three forebrain levels illustrate the brain distribution of mRNA transcripts encoding the α_1 subunit of the GABA_A receptor in mice exposed to room air (A–E) or 75% N₂O (B–F) for 48 h. Abbreviations: ac, anterior commissure; BL, basolateral amygdaloid nucleus; CA1, CA3, hippocampal subfields; CE, central amygdaloid nucleus; Cg, cingulate gyrus; Co, cortical amygdaloid nucleus; CP,

caudate putamen; DG, dentate gyrus; GP, globus pallidus; HDB, horizontal limb of the diagonal band nucleus; ic, internal capsule, LS, lateral septal nucleus; Me, medial amygdaloid nucleus; MS, medial septal nucleus; Par, parietal cortex, Pir, piriform cortex; VDB, vertical limb of the diagonal band nucleus, 3V, third ventricle.




Fig. 3. Darkfield photomicrographs of coronal sections at three brainstem levels illustrate the brain distribution of mRNA transcripts encoding the α_1 subunit of the GABA_A receptor in mice exposed to room air (A, C, E) or 75% N₂O (B, D, F) for 48 h. Abbreviations: Aq, cerebral aqueduct; gl, granular layer, cerebellum; IC, inferior colliculus; LV, lateral vestibular nucleus; MG, medial geniculate nucleus; ml, molecular layer, cerebellum; MV, medial vestibular nucleus; pl, purkinje layer, cerebellum; py, pyramidal tract; RN, red nucleus; SN, substantia nigra; Sub, subiculum; VTA, ventral tegmental area; Vtg, ventral tegmental nucleus; 4V, fourth ventricle; 7, facial nucleus.

Exposure of mice to 48 h of 75% N₂O produced a statistically significant increase in expression levels of mRNA for the α_1 subunit of the GABA_A receptor ($P < 0.001$, two-way ANOVA). There was also a statistically significant difference in expression levels of mRNA between brain regions ($P < 0.001$) and a statistically significant interaction between treatment and brain region ($P = 0.023$). Post-hoc statistical analysis using the Tukey test found significant increases following N₂O exposure in the cingulate cortex ($P = 0.002$), the CA1/2 region of the hippocampus ($P = 0.005$), the dentate gyrus ($P = 0.042$), the subiculum ($P = 0.047$), the medial septum ($P = 0.004$), and the ventral tegmental area ($P = 0.034$) (Table 1).

Table 1. The effect of nitrous oxide exposure on GABA_A α_1 subunit mRNA expression

Brain region	Room air exposure	N ₂ O exposure	% change	Significance
Cingulate cortex	27.90±1.09	38.03±1.68	36.3	$P=0.002^{**}$
Motor cortex	29.66±0.60	35.44±0.90	19.5	$P=0.082$
Primary somatosensory cortex	41.46±0.63	46.92±1.10	13.2	$P=0.167$
Secondary somatosensory cortex	43.73±0.71	44.50±1.27	1.8	$P=0.851$
Piriform cortex	56.80±1.38	61.51±2.67	8.3	$P=0.308$
CA1/2 region, Hippocampus	39.11±3.63	49.85±2.74	27.5	$P=0.005^{**}$
CA3 region, Hippocampus	19.44±2.22	25.09±1.85	29.1	$P=0.214$
Dentate gyrus	27.58±2.28	34.08±1.78	23.6	$P=0.042^*$
Subiculum	40.50±5.46	47.76±3.04	17.9	$P=0.047^*$
Medial septum	42.04±1.55	53.77±2.57	27.9	$P=0.004^{**}$
Ventral tegmental area	0.76±0.40	2.43±0.47	219.7	$P=0.034^*$
Purkinje layer, cerebellum	109.00±2.96	116.74±3.76	7.1	$P=0.228$

Granular layer, cerebellum	47.20±3.42	49.55±3.82	5.0	<i>P</i> =0.618
Molecular layer, cerebellum	18.23±2.21	17.23±1.98	-5.5	<i>P</i> =0.672
Basolateral amygdaloid nucleus	12.08±3.11	12.31±0.78	1.9	<i>P</i> =0.682
Medial amygdaloid nucleus	6.07±1.51	7.39±0.91	21.7	<i>P</i> =0.279
Central amygdaloid nucleus	1.62±0.79	0.50±0.35	-69.1	<i>P</i> =0.050
Red nucleus	46.19±2.99	42.10±3.23	-8.9	<i>P</i> =0.278
Ventral tegmental nucleus	87.20±4.6	83.58±5.06	-4.2	<i>P</i> =0.484
Globus pallidus	21.75±1.67	22.91±1.16	5.3	<i>P</i> =0.666

a. Levels of expression were measured using computerized image analysis and mean gray values were calculated from multiple sections throughout each region for each mouse. Data are reported as the average mean gray values±S.E.M. for mice exposed to room air (*n*=5) or nitrous oxide (*n*=8).

*Statistical significance was determined by the Tukey multiple comparison test following two-way analysis of variance. **P*<0.05; ***P*≤0.005.

4. Discussion

The present studies are the first to identify specific brain regions affected by N₂O exposure and indicate an up regulation of GABA_A α₁ subunit mRNA in the cingulate cortex, the hippocampus, the medial septum, and the VTA. Interestingly, these brain regions have all been implicated as sites of action of ethanol, and ethanol-induced changes in GABA_A receptor subunit expression have been identified in the cortex, hippocampus and VTA. Further, the VTA, as part of the mesolimbic dopaminergic pathway, has long been thought to be involved in the reinforcing actions of drugs of abuse, including ethanol [23], [24], [58], [59]. Ethanol can stimulate the activity of VTA dopaminergic neurons, and has been found to be reinforcing when administered directly into the VTA in operant conditioning experiments [15]. Mesolimbic GABAergic systems are also thought to be involved in the ethanol reinforcement properties, since muscimol injected into the VTA was shown to alter ethanol-reinforced responding [20].

Actions of ethanol and N₂O on the GABAergic system of the medial septum may be of particular functional importance. While the effects of ethanol on GABA_A receptor subunit composition in the medial septum have not been investigated, other effects of ethanol on this brain region have been extensively examined. The medial septum has been implicated as a brain region that influences the ethanol-induced impairment of nonspatial working memory [17] and depression of locomotion [5], as well as the sedative action of ethanol, enhancing the GABA-mediated inhibition of medial septal neuronal activity [16], [54]. Chronic ethanol exposure also produces long term effects on the septal GABAergic system, decreasing GABA's ability to inhibit ethanol-sensitive cells in the medial septum after ethanol withdrawal [9]. Thus, the changes in GABA_A α₁ subunit expression observed in the present study both support the hypothesis that N₂O effects are mediated, at least in part, through GABA_A receptors and provide further evidence for similar mechanisms of action of N₂O and ethanol.

Importantly, the increases in GABA_A α₁ subunit expression detected in the present study were not generalized to all regions under investigation. Areas examined in which differences could not be detected included the cerebellum, globus pallidus, red nucleus, basolateral and medial amygdaloid nuclei, as well as cortical areas such as the somatosensory and the piriform cortices. Interestingly, within the hippocampal formation, an up-regulation of α₁ subunits was noted within the dentate gyrus, CA1 and subiculum, but not in CA3, further suggesting the site-specific nature of the experimental effect. Moreover, regions in which increases were detected have all been implicated in aspects of ethanol addiction/withdrawal. If the present mRNA changes are indeed paralleled by similar increases at the protein level, the findings provide further evidence for a prominent role for GABAergic mechanisms in the reinforcing properties of N₂O and ethanol, and may yield insights into the sequence of cellular events underlying these processes.

The increase in α_1 subunit mRNA levels seen following N₂O exposure may reflect a generalized increase in the concentration of GABA_A receptors. It is also possible that N₂O exposure causes a modification of receptor composition, with an increase in the proportion of GABA_A receptors containing the α_1 subunit. There is evidence that ethanol can produce such an effect in rat cortex, where it has been shown that exposure to ethanol results in a decrease in α_1 subunit mRNA and protein approximately equal to the increase in the α_4 subunit mRNA and protein [11], [12]. A similar effect was found in the hippocampus, where long term administration of ethanol produced a decrease α_1 subunit mRNA and an increase in α_5 mRNA levels [8]. Such a change in subunit composition, with or without a change in the overall number of GABA_A receptors, could significantly change GABA_A receptor function, since the pharmacology and the binding characteristics of GABA_A receptor have been shown to be influenced by expression of different GABA_A receptor subunit combinations [1], [7], [10], [25], [28], [37], [38], [44], [51], [52]. Moreover, an increase in mRNA expression for the α_1 subunit produced by ethanol exposure in mice was previously shown to correspond to a decrease in GABA-dependent Cl⁻ flux [19]. The functional impact of N₂O-induced increases in the α_1 mRNA expression might similarly alter the efficacy of GABAergic inhibition. It should be noted, however, that N₂O-induced changes in mRNA levels for α_1 subunit may not correlate with changes in protein and/or changes in GABA_A receptor stoichiometry.

The functional significance of the changes in GABA_A receptor subunit expression produced by N₂O remains to be determined. Mouse strain has been shown to be an important variable in responsiveness to both the analgesic effects of N₂O [46], [47] and the ability of N₂O to induce withdrawal seizures [55]. Moreover, ethanol studies have shown that mouse strain can influence the effect of ethanol on GABA_A receptor subunit expression [6], [19], [60]. Therefore, determining the effect of N₂O on GABA_A receptor subunit expression in other mouse strains is an important consideration in the analysis of the functional impact of N₂O-induced changes in the GABA_A receptor. In addition, results of previous ethanol studies have shown that effects on GABA_A receptor subunit expression vary with the duration of drug exposure. Expression of α_1 , α_6 , β_3 and γ_2 subunits were differentially affected by acute and chronic ethanol administration [60] and varying durations of chronic ethanol exposures were necessary for changes in GABA_A receptor subunit expression in different brain regions [8], [42]. Varying the duration of N₂O exposure may also provide insight into the functional impact of N₂O-induced changes in the GABA_A receptor.

The N₂O-induced increase in GABA_A receptor α_1 subunit mRNA expression found in the present experiments supports the hypothesis that N₂O effects are produced, at least in part, through the GABA_A receptor, and localize the N₂O-induced effects to the cingulate cortex, hippocampus, ventral tegmental area and medial septum. These results are also consistent with the hypothesis that ethanol and N₂O produce dependence and withdrawal through common mechanisms. Future studies on the effect of N₂O on the full complement of subunits of the GABA_A receptor, as well as additional receptor types, will be required for a more complete understanding of the neuronal mechanism of action of N₂O.

Acknowledgements

This work was supported by NIH grants NS30708, MH56577, DE09998, and the Howard Hughes Medical Institute Undergraduate Research Award Program. The technical assistance of Elizabeth Dallas, Mary Ann Haasch, Krista Johaneck, Julie Lenza, Julie Otto, and Mike Reilly is gratefully acknowledged.

References

- [1] E.A. Barnard, M. Sutherland, S. Zaman, M. Matsumoto, N. Nayeem, T. Green, M.G. Darlison, A.N. Bateson **Multiplicity, structure, and function in GABAA receptors** Ann. N.Y. Acad. Sci., 707 (1993), pp. 116-125
- [2] H.C. Becker, R.F. Anton **The benzodiazepine receptor inverse agonist RO15-4513 exacerbates, but does not precipitate, ethanol withdrawal in mice** Pharmacol. Biochem. Behav., 32 (1989), pp. 163-167
- [3] J.K. Belknap, S.E. Laursen, J.C. Crabbe **Ethanol and nitrous oxide produce withdrawal-induced convulsions by similar mechanisms in mice** Life Sci., 41 (1987), pp. 2033-2040

- [4] J.K. Belknap, P. Metten, M.L. Helms, L.A. O'Toole, S. Angeli-Gade, J.C. Crabbe, T.J. Phillips **Quantitative trait loci (QTL) applications to substances of abuse: physical dependence studies with nitrous oxide and ethanol in BXD mice** *Behav. Genetics*, 23 (1993), pp. 213-222
- [5] G.R. Breese, G.D. Frye, T.J. McCown, R.A. Mueller **Comparison of the CNS effects induced by TRH and bicuculline after microinjection into medial septum, substantia nigra and inferior colliculus: absence of support for a GABA antagonist action for TRH** *Pharmacol. Biochem. Behav.*, 21 (1984), pp. 145-149
- [6] K.J. Buck, L. Hahner, J. Sikela, R.A. Harris **Chronic ethanol treatment alters brain levels of gamma-aminobutyric acid_A receptor subunit mRNAs: relationship to genetic differences in ethanol withdrawal seizure severity** *J. Neurochem.*, 57 (1991), pp. 1452-1455
- [7] D.R. Burt, G.L. Kamatchi **GABA_A receptor subtypes: from pharmacology to molecular biology** *FASEB J.*, 5 (1991), pp. 2916-2923
- [8] M.E. Charlton, P.M. Sweetnam, L.W. Fitzgerald, R.Z. Terwilliger, E.J. Nestler, R.S. Duman **Chronic ethanol administration regulates the expression of GABA_A receptor α_1 and α_5 subunits in the ventral tegmental area and hippocampus** *J. Neurochem.*, 68 (1997), pp. 121-127
- [9] H.E. Criswell, P.E. Simson, K.B. Johnson, G.R. Breese **Chronic ethanol decreases the ability of GABA to inhibit ethanol-sensitive neurons in the medial septum** *Alcohol. Clin. Exp. Res.*, 17 (1993), p. 477
- [10] T.M. deLorey, R.E. Olsen **Gamma-aminobutyric acid_A receptor structure and function** *J. Biol. Chem.*, 267 (1992), pp. 16747-16750
- [11] L.L. Devaud, J.M. Fritschy, W. Sieghart, A.L. Morrow **Bidirectional alterations of GABA_A receptor subunit peptide levels in rat cortex during chronic ethanol consumption and withdrawal** *J. Neurochem.*, 69 (1997), pp. 126-130
- [12] L.L. Devaud, F.D. Smith, D.R. Grayson, A.L. Morrow **Chronic ethanol consumption differentially alters the expression of γ -aminobutyric acid_A receptor subunit mRNAs in rat cerebral cortex: competitive, quantitative reverse transcriptase-polymerase chain reaction analysis** *Mol. Pharmacol.*, 48 (1995), pp. 861-868
- [13] D.E. Emmanouil, C.H. Johnson, R.M. Quock **Nitrous oxide anxiolytic effect in mice in the elevated plus maze: Mediation by benzodiazepine receptors** *Psychopharmacology*, 115 (1994), pp. 167-172
- [14] K.B.J. Franklin, G. Paxinos *The Mouse Brain in Stereotaxic Coordinates*, Academic Press, NY (1997)
- [15] G.J. Gatto, W.J. McBride, J.M. Murphy, L. Lumeng, T.K. Li **Ethanol self-infusion into the ventral tegmental area by alcohol-preferring rats** *Alcohol*, 11 (1994), pp. 557-564
- [16] B.S. Givens, G.R. Breese **Site-specific enhancement of gamma-aminobutyric acid-mediated inhibition of neural activity by ethanol in the rat medial septal area** *J. Pharmacol. Exp. Ther.*, 254 (1990), pp. 528-538
- [17] B. Givens, K. McMahon **Effects of ethanol on nonspatial working memory and attention in rats** *Behav. Neurosci.*, 111 (1997), pp. 275-282
- [18] M.H. Harper, P.M. Winter, B.H. Johnson, D.D. Koblin, E.I. Eger 2d **Withdrawal convulsions in mice following nitrous oxide** *Anesth. Analgesia*, 59 (1980), pp. 19-21
- [19] M. Hirouchi, T. Hashimoto, K. Kuriyama **Alteration of GABA_A receptor α_1 -subunit mRNA in mouse brain following continuous ethanol inhalation** *Eur. J. Pharmacol.*, 247 (1993), pp. 127-130
- [20] C.W. Hodge, M. Haraguchi, H.H. Samson **Effects of ventral tegmental microinjections of the GABA_A agonist muscimol on ethanol reinforced responding** *Alcohol. Clin. Exp. Res.*, 17 (1993), p. 487
- [21] D.D. Koblin, J.E. Deady **Anaesthetic requirement in mice selectively bred for differences in ethanol sensitivity** *Br. J. Anaesth.*, 53 (1981), pp. 5-10
- [22] D.D. Koblin, J.E. Deady, D.E. Dong, E.I. Eger 2d **Mice tolerant to nitrous oxide are also tolerant to alcohol** *J. Pharmacol. Exp. Ther.*, 213 (1980), pp. 309-312
- [23] G.F. Koob **Drugs of abuse: anatomy, pharmacology and function of reward pathways** *Trends Pharmacol. Sci.*, 13 (1992), pp. 177-184
- [24] G.F. Koob, A.J. Robers, G. Schulteis, L.H. Parsons, C.J. Heyser, P. Hyytia, E. Merlo-Pich, F. Weiss **Neurocircuitry targets in ethanol reward and dependence** *Alcohol. Clin. Exp. Res.*, 22 (1998), pp. 3-9

- [25] E.S. Levitan, P. R Schofield, D.R. Burt, L.M. Rhee, W. Wisden, M. Kohler, N. Fujita, H.F. Rodriguez, A. Stephenson, M.G. Darlison, E.A. Barnard, P.H. Seeburg **Structural and functional basis for GABA_A receptor heterogeneity** *Nature (London)*, 335 (1988), pp. 76-79
- [26] J. L Lewis, R.J. Pruhs, R.M. Quock **Modification of an infant incubator for exposure of experimental animals to nitrous oxide** *J. Pharmacol. Methods*, 18 (1987), pp. 143-150
- [27] R.G. Lister, J.W. Karanian **RO 15-4513 induces seizures in DBA/2 mice undergoing alcohol withdrawal** *Alcohol*, 4 (1987), pp. 409-411
- [28] H. Luddens, P.H. Seeburg, E.R. Korpi **Impact of β and γ variants of ligand-binding properties of γ -aminobutyric acid type A receptors** *Mol. Pharmacol.*, 45 (1994), pp. 810-814
- [29] A.J. MacLennan, N. Brecha, M. Khrestchatsky, C. Sternini, N.J. Tillakaratne, M.Y. Chiang, K. Anderson, M. Lai, A.J. Tobin **Independent cellular and ontogenetic expression of mRNAs encoding three alpha polypeptides of the rat GABA_A receptor** *Neuroscience*, 43 (1991), pp. 369-380
- [30] P. Malherbe, E. Sigel, R. Baur, E. Persohn, J.G. Richards, H. Mohler **Functional characteristics and sites of gene expression of the alpha1, beta1, gamma2-isoform of the rat GABA_A receptor** *J. Neurosci.*, 10 (1990), pp. 2330-2337
- [31] H.J. Manson, G. Dyke, J. Melling, M. Gough **The effect of naloxone and morphine on convulsions in mice following withdrawal from nitrous oxide** *Can. Anaesth. Soc. J.*, 30 (1983), pp. 28-31
- [32] D.B. Mathews, L.L. Devaud, J. M Fritschy, W. Sieghart, A.L. Morrow **Differential regulation of GABA_A receptor gene expression by ethanol in the rat hippocampus versus cerebral cortex** *J. Neurochem.*, 70 (1998), pp. 1160-1166
- [33] M.C. Mhatre, G. Pena, W. Sieghart, M.K. Ticku **Antibodies specific for GABA_A receptor alpha subunits reveal that chronic alcohol treatment down-regulates alpha-subunit expression in rat brain regions** *J. Neurochem.*, 61 (1993), pp. 1620-1625
- [34] M.C. Mhatre, M.K. Ticku **Chronic ethanol administration alters γ -aminobutyric acid A receptor gene expression** *Mol. Pharmacol.*, 42 (1992), pp. 415-422
- [35] M.C. Mhatre, M.K. Ticku **Chronic ethanol treatment upregulates the GABA receptor beta subunit expression** *Brain Res. Mol. Brain Res.*, 23 (1994), pp. 246-252
- [36] B. Milne, F.W. Cervenko, K.H. Jhamandas **Physical dependence on nitrous oxide in mice: resemblance to alcohol but not to opiate withdrawal** *Can. Anaesth. Soc. J.*, 28 (1981), pp. 46-50
- [37] H. Mohler, D. Benke, J. Benson, B. Luscher, J.M. Fritschy **GABAA-receptor subtypes in vivo: cellular localization, pharmacology and regulation** *Adv. Biochem. Psychopharmacol.*, 48 (1995), pp. 41-46
- [38] H. Mohler, J.M. Fritschy, B. Luscher, U. Rudolph, J. Benson, D. Benke **The GABAA receptors. From subunits to diverse functions** *Ion Channels*, 4 (1996), pp. 89-113
- [39] P. Montpied, A. L Morrow, J.W. Karanian, E.I. Ginns, B.M. Martin, S. M Paul **Prolonged ethanol inhalation decreases gamma-aminobutyric acid_A receptor alpha subunit mRNAs in the rat cerebral cortex** *Mol. Pharmacol.*, 39 (1991), pp. 157-163
- [40] A.L. Morrow, L.L. Devaud, D. Bucci, F.D. Smith **GABAA and NMDA receptor subunit expression in ethanol dependent rats** *Alcohol Alcohol. Suppl.*, 2 (1994), pp. 89-95
- [41] A.L. Morrow, P. Montpied, A. Lingford-Huges, S.M. Paul **Chronic ethanol and pentobarbital administration in the rat: effects on GABAA receptor function and expression in brain** *Alcohol*, 7 (1990), pp. 237-244
- [42] J. Ortiz, L.W. Fitzgerald, M. Charlton, S. Lane, L. Trevisan, X. Guitart, W. Shoemaker, R.S. Duman, E.J. Nestler **Biochemical actions of chronic ethanol exposure in the mesolimbic dopamine system** *Synapse*, 21 (1995), pp. 289-298
- [43] E. Persohn, P. Malherbe, J.G. Richards **Comparative molecular neuroanatomy of cloned GABA_A receptor subunits in the rat CNS** *J. Comp. Neurol.*, 326 (1992), pp. 193-216
- [44] D.B. Pritchett, H. Luddens, P.H. Seeburg **Type I and type II GABA_A-benzodiazepine receptors produced in transfected cells** *Science*, 245 (1989), pp. 1389-1392
- [45] R.M. Quock, D.E. Emmanouil, L.K. Vaughn, R.J. Pruhs **Benzodiazepine receptor mediation of behavioral effects of nitrous oxide in mice** *Psychopharmacology*, 107 (1992), pp. 310-314

- [46] R.M. Quock, J.L. Mueller, L.K. Vaughn **Strain-dependent differences in responsiveness of mice to nitrous oxide (N₂O) antinociception** Brain Res., 614 (1993), pp. 52-56
- [47] R.M. Quock, J.M. Mueller, L.K. Vaughn, J.K. Belknap **Nitrous oxide antinociception in BXD recombinant inbred mouse strains and identification of quantitative trait loci** Brain Res., 725 (1996), pp. 23-29
- [48] R.M. Quock, L.K. Vaughn **Nitrous oxide: Mechanism of its antinociceptive action** Analgesia, 1 (1995), pp. 151-159
- [49] R.M. Quock, J.A. Wojcechowskyj, D.E. Emmanoui **Comparison of nitrous oxide, morphine and diazepam effects in the mouse staircase test** Psychopharmacology, 92 (1987), pp. 324-326
- [50] J. Ruprecht, B. Dworacek, R. Ducardus, P.I.M. Schmitz, M.R. Dzoljic **The involvement of the central cholinergic and endorphinergic systems in the nitrous oxide withdrawal syndrome in mice** Anesthesiology, 58 (1983), pp. 524-526
- [51] P.H. Seeburg, W. Wisden, T.A. Verdoorn, D.B. Pritchett, P. Werner, A. Herb, H. Luddens, R. Sprengel, B. Sakmann **The GABA_A receptor family: molecular and functional diversity** Cold Spring Harb. Symp. Quant. Biol., 55 (1990), pp. 29-38
- [52] W. Sieghart **Structure and pharmacology of gamma-aminobutyric acidA receptor subtypes** Pharmacol. Rev., 47 (1995), pp. 181-234
- [53] R.A. Smith, P.M. Winter, M. Smith, E. I Eger II **Convulsions in mice after anesthesia** Anesthesiology, 50 (1979), pp. 501-504
- [54] B.L. Soldo, W.R. Proctor, T.V. Dunwiddie **Ethanol differentially modulates GABA_A receptor-mediated chloride currents in hippocampal, cortical, and septal neurons in rat brain slices** Synapse, 18 (1994), pp. 94-103
- [55] L.K. Vaughn, R.J. Pruhs **Strain-dependent variability in nitrous oxide withdrawal seizure frequency** Life Sci., 57 (1995), pp. 1125-1130
- [56] L.K. Vaughn, D. K Marsee **Ro 15-4513 increases nitrous oxide withdrawal seizure frequency in mice** Neurosci. Abstr., 19 (1993), p. 1457
- [57] W. Wisden, D.J. Laurie, H. Monyer, P.H. Seeburg **The distribution of 13 GABA_A receptor subunit mRNAs in the rat brain. I. Telencephalon, diencephalon, mesencephalon** J. Neurosci., 12 (1992), pp. 1040-1062
- [58] R.A. Wise **Actions of drugs of abuse on brain reward systems** Pharmacol. Biochem. Behav., 13 (Suppl. 1) (1980), pp. 213-223
- [59] R.A. Wise **The role of reward pathways in the development of drug dependence** Pharmacol. Ther., 35 (1987), pp. 227-263
- [60] C.H. Wu, A. Frosthalm, A.L. DeBlas, A. Rotter **Differential expression of GABA_A/benzodiazepine receptor subunit mRNAs and ligand binding sites in mouse cerebellar neurons following in vivo ethanol administration: An autoradiographic analysis** J. Neurochem., 65 (1995), pp. 1229-1239
- [61] N.R. Zahniser, K.J. Buck, P. Curella, S.J. McQuikin, D. Wilson-Shaw, C.L. Miller, R.L. Klein, K.A. Heidenreich, W.J. Keir, J.M. Sikela, R.A. Harris **GABA_A receptor function and regional analysis of subunit mRNAs in long-sleep and short-sleep mouse brain** Brain Res. Mol. Brain Res., 14 (1992), pp. 196-206
- [62] J.H. Zar **Biostatistical Analysis (2nd Edition)**, Prentice Hall, Inc, Englewood Cliffs, NJ (1984)