

Marquette University

e-Publications@Marquette

---

Physics Faculty Research and Publications

Physics, Department of

---

2004

## Mononuclear Nitrogen/sulfur-ligated Cobalt(II) Methoxide Complexes: Structural, EPR, Paramagnetic $^1\text{H}$ NMR, and Electrochemical Investigations

Kyle J. Tubbs  
*Utah State University*

Ewa Szajna  
*Utah State University*

Brian Bennett  
*Marquette University, brian.bennett@marquette.edu*

Jason A. Halfen  
*University of Wisconsin - Eau Claire*

Rex W. Watkins  
*Utah State University*

*See next page for additional authors*

Follow this and additional works at: [https://epublications.marquette.edu/physics\\_fac](https://epublications.marquette.edu/physics_fac)

 Part of the [Physics Commons](#)

---

### Recommended Citation

Tubbs, Kyle J.; Szajna, Ewa; Bennett, Brian; Halfen, Jason A.; Watkins, Rex W.; Arif, Atta M.; and Berreau, Lisa M., "Mononuclear Nitrogen/sulfur-ligated Cobalt(II) Methoxide Complexes: Structural, EPR, Paramagnetic  $^1\text{H}$  NMR, and Electrochemical Investigations" (2004). *Physics Faculty Research and Publications*. 82.

[https://epublications.marquette.edu/physics\\_fac/82](https://epublications.marquette.edu/physics_fac/82)

---

**Authors**

Kyle J. Tubbs, Ewa Szajna, Brian Bennett, Jason A. Halfen, Rex W. Watkins, Atta M. Arif, and Lisa M. Berreau

Marquette University

**e-Publications@Marquette**

***Physics Faculty Research and Publications/College of Arts and Sciences***

***This paper is NOT THE PUBLISHED VERSION; but the author's final, peer-reviewed manuscript.*** The published version may be accessed by following the link in the citation below.

*Dalton Transactions*, (July 20, 2004): 2398-2399. [DOI](#). This article is © Royal Society of Chemistry and permission has been granted for this version to appear in [e-Publications@Marquette](#). Royal Society of Chemistry does not grant permission for this article to be further copied/distributed or hosted elsewhere without the express permission from Royal Society of Chemistry.

# Mononuclear nitrogen/sulfur-ligated cobalt(ii) methoxide complexes: Structural, EPR, paramagnetic $^1\text{H}$ NMR, and electrochemical investigations

Kyle J. Tubbs

Department of Chemistry and Biochemistry, Utah State University, Logan, UT

Ewa Szajna

Department of Chemistry and Biochemistry, Utah State University, Logan, UT

Brian Bennett

Department of Biophysics, Medical College of Wisconsin, Milwaukee, WI

Jason A. Halfen

Department of Chemistry, University of Wisconsin-Eau Claire, Eau Claire, WI

Rex W. Watkins

Department of Chemistry and Biochemistry, Utah State University, Logan, UT

Atta M. Arif


Department of Chemistry, University of Utah, Salt Lake City, Utah

## Lisa M. Berreau

Department of Chemistry and Biochemistry, Utah State University, Logan, UT

### Abstract

The first examples of mononuclear nitrogen/sulfur-ligated Co(II) alkoxide complexes, species of relevance to a reactive intermediate observed for Co(II)-substituted liver alcohol dehydrogenase, have been isolated and characterized by multiple methods including X-ray crystallography, EPR, paramagnetic  $^1\text{H}$  NMR, and cyclic voltammetry.




Mononuclear alkoxide complexes of the late transition metals are rare, due in part to the propensity of alkoxide ligands to bridge metal centers.<sup>1</sup> To date, four mononuclear Co(ii) complexes having terminal alkoxide ligation have been structurally characterized.<sup>2,3</sup> One of these is a tetrahedral cationic  $\text{N}_3$ -ligated Co(ii) ethoxide complex isolated using a chelate ligand that encapsulates the alkoxide moiety within a hydrophobic pocket.<sup>2</sup> The others are tetrahedral or trigonal planar species having bulky  $-\text{O}-\text{C}(t\text{-Bu})_3$  or  $-\text{O}-\text{C}(\text{Ph})_3$  ligation.<sup>3</sup>

In the zinc enzyme liver alcohol dehydrogenase (LADH), a nitrogen/sulfur-ligated zinc alkoxide species is proposed to be the reactive moiety for hydride transfer to  $\text{NAD}^+$  during alcohol oxidation.<sup>4</sup> This alkoxide moiety is stabilized *via* hydrogen-bonding to a nearby amino acid residue (Ser48).<sup>4,5</sup> As zinc is spectroscopically silent ( $d^{10}$ ), evidence for the proposed alkoxide intermediate in the alcohol dehydrogenase catalytic cycle has been obtained *via* UV-vis studies of a Co(ii)-substituted form of enzyme that retains catalytic activity, albeit 50–70% less than that of the native enzyme.<sup>6</sup>


We have previously shown that a mononuclear nitrogen/sulfur-ligated zinc methoxide complex can be isolated using a mixed nitrogen/sulfur chelate ligand which provides two internal hydrogen bond donors to the zinc-bound alkoxide oxygen atom.<sup>7</sup> To date, there are no examples in the chemical literature of structurally-related mononuclear zinc and cobalt alkoxide complexes of relevance to LADH. Thus, there is no chemical precedent for understanding how Co(ii) substitution in LADH may influence the chemical reactivity of the metal alkoxide moiety. Outlined herein are studies toward evaluating the chemical properties of mononuclear cobalt methoxide complexes supported by a specific nitrogen/sulfur chelate ligand.

Treatment of  $\text{ebnpa}^7$  or  $\text{pbnpa}^8$  with an equimolar amount of  $\text{Co}(\text{ii})(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$  and  $\text{Me}_4\text{NOH} \cdot 5\text{H}_2\text{O}$  in dry methanol (Scheme 1), followed by work-up and recrystallization from  $\text{CH}_3\text{OH}/\text{Et}_2\text{O}$  yielded green crystalline samples of  $[(\text{ebnpa})\text{Co}-\text{OCH}_3]\text{ClO}_4$  (**1**) and  $[(\text{pbnpa})\text{Co}-\text{OCH}_3]\text{ClO}_4$  (**2**) in yields of 78–79%. Single crystals of these complexes suitable for X-ray crystallographic studies were obtained *via* crystallization of each from  $\text{MeOH}/\text{Et}_2\text{O}$  in the presence of  $\text{Me}_4\text{NClO}_4$ . $\ddagger$  Complexes **1** and **2** have been characterized by elemental analysis, X-ray crystallography, EPR, paramagnetic  $^1\text{H}$  NMR, cyclic voltammetry, UV-vis and FTIR. Analogues of **1** having deuteration at the alkoxide methyl group and selected positions of the  $\text{ebnpa}$  chelate ligand have been generated to enable assignment of paramagnetic  $^1\text{H}$  NMR resonances (see ESI $\dagger$ ).



### Scheme 1

Representations of the cations of **1** and **2** are shown in Fig. 1. Similar to the zinc ion in  $[(\text{ebnpa})\text{Zn}-\text{OCH}_3]\text{ClO}_4$  ( $\tau = 0.84, 0.88$ ),<sup>7,9,10</sup> the Co(ii) centers of **1** and **2** each exhibit a slightly distorted trigonal bipyramidal geometry (**1**:  $\tau = 0.90$ ; **2**:  $\tau = 0.93$ )<sup>9</sup> with the pyridyl nitrogen donors and the thioether sulfur in the pseudo trigonal plane. For **1**, **2**, and  $[(\text{ebnpa})\text{Zn}-\text{OCH}_3]\text{ClO}_4$ , a prominent equatorial angular deviation from the expected  $120^\circ$  involves an acute  $\text{N}(\text{py})-\text{M}-\text{N}(\text{py})$  bond angle (**1**:  $111.84(5)^\circ$ , **2**:  $110.61(6)^\circ$ ,  $[(\text{ebnpa})\text{Zn}-\text{OCH}_3]\text{ClO}_4$ :<sup>10</sup>  $113.2(4)^\circ/111.9(4)^\circ$ ). The axial  $\text{Co}-\text{O}$  and  $\text{Co}-\text{N}(\text{am})$  distances in **1** (Table 1) are slightly longer ( $\sim 0.02 \text{ \AA}$ ) than similar linkages in **2**. The equatorial  $\text{Co}-\text{N}(\text{py})$  and  $\text{Co}-\text{S}$  distances in **1** are shorter than those found in **2**, which is, in part, an indication of enhanced donor ability of the  $-\text{SEt}$  substituent of **1** versus the  $-\text{SPh}$  donor of **2**. The Co(ii) derivative **1** exhibits similar bond distances to its zinc analogue  $[(\text{ebnpa})\text{Zn}-\text{OCH}_3]\text{ClO}_4$ , with the largest deviation being in the metal-sulfur unit ( $\sim 0.05 \text{ \AA}$ ), where the Co(ii) complex has a shorter M-S interaction (Table 1). The general similarity of the bond distances in **1** and  $[(\text{ebnpa})\text{Zn}-\text{OCH}_3]\text{ClO}_4$  is consistent with the nearly identical ionic radii of five-coordinate Co(ii) and Zn(ii) (81 and 82 pm, respectively).<sup>11</sup> Finally, the  $\text{Co}-\text{O}(1)-\text{C}$  angles in **1** and **2** ( $126.12(11)^\circ$  and  $122.18(12)^\circ$ , respectively) are the most acute reported to date for mononuclear cobalt alkoxide linkages,<sup>2,3</sup> and are generally similar to the angles observed for the two independent cations of  $[(\text{ebnpa})\text{Zn}-\text{OCH}_3]\text{ClO}_4$  ( $123.1(8)^\circ$  and  $124.5(8)^\circ$ ).<sup>10</sup>



**Fig. 1** Representations of the cationic portions of **1** and **2**. See Table 1 for selected bond distances.

**Table 1** Selected bond distances ( $\text{\AA}$ ) in **1**, **2**, and  $[(\text{ebnpa})\text{Zn}-\text{OCH}_3]\text{ClO}_4$ <sup>7</sup>

Compound	$\tau^a$	M-O	M-N(py)	M-N(am)	M-S
<b>1</b>	0.90	1.9302(11)	2.0852(12), 2.0885(12)	2.1919(13)	2.4387(4)
<b>2</b>	0.93	1.9125(12)	2.1023(14), 2.1060(15)	2.1741(14)	2.5023(5)
$[(\text{ebnpa})\text{Zn}-\text{OCH}_3]\text{ClO}_4$ <sup>10</sup>	0.84	1.953(8)	2.095(9), 2.076(9)	2.212(10)	2.490(3)
	0.88	1.948(8)	2.086(10), 2.105(9)	2.192(10)	2.496(3)

a A. W. Addison, T. N. Rao, J. Reedijk, J. van Rijn and G. C. Verschoor, *J. Chem. Soc., Dalton Trans.*, 1984, 1349.

Preliminary EPR analysis of **1** and **2** (0.46 and 0.49 mM, respectively) indicates that both complexes are high spin. As shown in Fig. S1, each complex exhibits a well-defined  $S = 3/2$   $M_s = |\pm 1/2\rangle$  signal indicating a system having modest spin-orbit coupling and a high degree of rhombic distortion of the axial zero-field splitting. Simulation of the line shape for **1** yielded  $g(x,y) = 2.395$ ,  $g_z = 2.060$ ,  $A_{\nu}(^{59}\text{Co}) = 0.0068 \text{ cm}^{-1}$ , and  $E/D = 0.233$ .

These parameters suggest an electronically distorted 5- or 6-coordinate site with little electron delocalization away from the Co(ii) ion. Uneven intensities of the resolved hyperfine lines in the spectrum of **2**, centered at  $g_{\text{eff}} \sim 6$ , indicate the presence of two species. Nevertheless, computer simulations show these species to be extremely similar in terms of electronic structure.

Paramagnetic  $^1\text{H}$  NMR has been used extensively to characterize active site structural features and reactions of Co(ii)-substituted zinc enzymes including LADH.<sup>12</sup> We have found NMR useful for analyzing the solution properties of **1** and **2**. Shown in Fig. S2 is the  $^1\text{H}$  NMR spectrum of a 2.3 mM solution of **1** in  $\text{CD}_3\text{CN}$  at 302 K. On the basis of integration,  $T_1$  values, deuteration and complementary  $^2\text{H}$  NMR studies, and comparison of the observed spectral features of **1** with those of **2** (Fig. S3) at a similar concentration, we have assigned several of the proton resonances of **1** (Table S1). For example, a single neopentyl methyl resonance is easily identifiable at 10.6 ppm. The  $\beta$ -protons of the pyridyl rings of **1** appear as two sharp singlets at 66.3 and 50.7 ppm, respectively, with the  $\gamma$ -H (*para* to the pyridyl nitrogen) being observed at 7.3 ppm. The observation of only one set of pyridyl proton resonances indicates that an effective plane of symmetry is present for **1** in  $\text{CD}_3\text{CN}$  solution. Selective deuteration of the neopentyl methylene position of the ebnpa ligand (using **1-d<sub>4</sub>-n**) enabled the identification of two resonances at 6.1 and 3.7 ppm, each of which integrates to 2H, consistent with the “inner” and “outer” chemical environment of these protons in the solid state structure of **1**. Although a resonance for the alkoxide methyl protons could not be identified in the  $^1\text{H}$  NMR spectrum of **1**, using  $[(\text{ebnpa})\text{Co}-\text{OCD}_3]\text{ClO}_4$  (**1-d<sub>3</sub>**) and  $^2\text{H}$  NMR, the alkoxide methyl proton resonance was identified at  $\sim 79$  ppm. Notably, acquisition of  $^1\text{H}$  NMR spectra of **1** and **2** at higher concentrations (*e.g.*  $\sim 24$ – $26$  mM, Fig. S4 and S5) indicates the presence of two species for each complex. We currently speculate that the second species for each is a dimeric complex, as the single set of resonances for the low concentration species for **1** and **2** can be regenerated *via* dilution of the higher concentration solutions.

Compounds **1** and **2** each exhibit a single, quasi-reversible ( $i_{\text{pa}} \gg i_{\text{pc}}$ ) redox couple (Fig. S6) at  $\sim 1460$  and  $\sim 1483$  mV, respectively, *versus* SCE. The reversibility of these couples is not influenced by changes in potential scan range or rate. The anodic peak current ( $i_{\text{pa}}$ ) for each oxidative wave is of similar magnitude to the anodic current of an internal standard ferrocene/ferrocenium couple. Since both the analyte and ferrocene were present in approximately equal concentrations in these experiments, the parity of the two anodic currents suggests that the observed oxidative processes correspond to the removal of one electron from the original cobalt(ii) alkoxide complexes.

In conclusion, the isolation of **1** under the same conditions used for the zinc analogue  $[(\text{ebnpa})\text{Zn}-\text{OCH}_3]\text{ClO}_4$  suggests similar hydrolytic stability for the Co(ii) and Zn(ii) methoxide derivatives. This observation leads us to hypothesize that the Co(ii)–OR species that is spectroscopically observable for Co(ii)-containing LADH is of similar hydrolytic stability to the proposed Zn(ii)–OR intermediate in the catalytic cycle of the native enzyme. However, from these studies we also note that unlike  $[(\text{ebnpa})\text{Zn}-\text{OCH}_3]\text{ClO}_4$ , **1** and **2** exhibit concentration dependent  $^1\text{H}$  NMR features, suggestive of the presence of a monomer/dimer equilibrium in acetonitrile solution.

We thank the National Science Foundation (CAREER Award CHE-0094066 to LMB, CHE-0243591 to JAH) and the National Institutes of Health (AI056231 to BB, EB001980 to the Medical College of Wisconsin) for funding of this research.

## Notes and references

1. D. C. Bradley, R. C. Mehrotra, I. P. Rothwell and A. Singh, *Alkoxo and Aryloxo Derivatives of Metals*, Academic Press, San Diego, 2001,

2. S. J. Archibald , S. P. Foxon , J. D. Freeman , J. E. Hobson , R. N. Perutz and P. H. Walton , *Dalton Trans.*, 2002, 2797
3. (a) M. M. Olmstead , P. P. Power and G. Sigel , *Inorg. Chem.*, 1986, **25** , 1027. (b) G. A. Sigel , R. A. Bartlett , D. Decker , M. M. Olmstead and P. P. Power , *Inorg. Chem.*, 1987, **26** , 1773
4. S. Ramaswamy , D.-H. Park and B. V. Plapp , *Biochemistry*, 1999, **38** , 13951
5. P. K. Agarwal , S. P. Webb and S. Hammes-Schiffer , *J. Am. Chem. Soc.*, 2000, **122** , 4803
6. C. Sartorius , M. Gerber , M. Zeppezauer and M. F. Dunn , *Biochemistry*, 1987, **26** , 871
7. D. K. Garner , S. B. Fitch , L. H. McAlexander , L. M. Bezold , A. M. Arif and L. M. Berreau , *J. Am. Chem. Soc.*, 2002, **124** , 9970
8. K. J. Tubbs , A. L. Fuller , B. Bennett , A. M. Arif , M. M. Makowska-Grzyska and L. M. Berreau , *Dalton Trans.*, 2003, 3111
9. A. W. Addison , T. N. Rao , J. Reedijk , J. van Rijn and G. C. Verschoor , *J. Chem. Soc., Dalton Trans.*, 1984, 1349
10. Two independent molecules are present in the asymmetric unit of [(ebnpa)Zn–OCH<sub>3</sub>]ClO<sub>4</sub> ([ref. 7](#))
11. J. E. Huheey , E. A. Keiter and R. L. Keiter , *Inorganic Chemistry: Principles of Structure and Reactivity* , HarperCollins, New York, NY, 1993,
12. I. Bertini and C. Luchinat , *Adv. Inorg. Biochem.*, 1984, **6** , 71.

## Footnotes

1. † Electronic supplementary information (ESI) available: Experimental details, and Figs. S1 (EPR spectra of **1** and **2** in MeOH soln.), S2 and S4 (<sup>1</sup>H NMR spectra of **1** in CD<sub>3</sub>CN), S3 and S5 (<sup>1</sup>H NMR spectra of **2** in CD<sub>3</sub>CN), S6 (cyclic voltammetry of **1** and **2**) and Table S1 (<sup>1</sup>H NMR features of **1** and **2** in CD<sub>3</sub>CN at 302 K). See <http://www.rsc.org/suppdata/dt/b4/b408177c/>
2. ‡ *Crystal data*: for **1**: C<sub>27</sub>H<sub>46</sub>ClCoN<sub>5</sub>O<sub>5</sub>S, *M* = 647.13, triclinic, space group *P* $\bar{1}$ , *a* = 8.5503(2), *b* = 13.7775(2), *c* = 14.6350(4) Å,  $\alpha$  = 77.3731(14),  $\beta$  = 89.8851(12),  $\gamma$  = 74.0184(14)°, *V* = 1614.20(6) Å<sup>3</sup>, *Z* = 2,  $\mu$  = 0.721 mm<sup>-1</sup>. Using Mo K $\alpha$  radiation (0.71073 Å), a total of 11777 reflections were collected (6.32 < 2 $\theta$  < 55.02°), of which 7305 were independent. Refinement converged to *R*<sub>1</sub> = 0.0323, *wR*<sub>2</sub> = 0.0768 (*I* > 2 $\sigma$ *I*) and *R*<sub>1</sub> = 0.0410, *wR*<sub>2</sub> = 0.0819 (all data). The perchlorate anion was disordered over two positions. For **2**: C<sub>31</sub>H<sub>46</sub>ClCoN<sub>5</sub>O<sub>5</sub>S, *M* = 695.17, triclinic, space group *P* $\bar{1}$ , *a* = 10.8800(2), *b* = 12.0599(3), *c* = 13.9550(4) Å,  $\alpha$  = 72.8711(10),  $\beta$  = 77.7458(15),  $\gamma$  = 75.1747(14)°, *V* = 1672.92(7) Å<sup>3</sup>, *Z* = 2,  $\mu$  = 0.701 mm<sup>-1</sup>. Using Mo K $\alpha$  radiation (0.71073 Å), a total of 12007 reflections were collected (6.32 < 2 $\theta$  < 55.00°), of which 7611 were independent. Refinement converged to *R*<sub>1</sub> = 0.0350, *wR*<sub>2</sub> = 0.0739 (*I* > 2 $\sigma$ *I*) and *R*<sub>1</sub> = 0.0502, *wR*<sub>2</sub> = 0.0808 (all data). CCDC reference numbers: **1**, 240280; **2**, 240281. See <http://www.rsc.org/suppdata/dt/b4/b408177c/> for crystallographic data in CIF or other electronic format.