Marquette University

e-Publications@Marquette

Exercise Science Faculty Research and
Publications

Exercise Science, Department of

2018

Impaired Trunk Flexor Strength, Fatigability, and Steadiness in Postpartum Women.

Rita Deering
Marquette University

Meredith Cruz
Marquette University

Jonathon Senefeld
Marquette University

Tatyana Pashibin
Marquette University

Sarah Eickmeyer
Marquette University

See next page for additional authors

Follow this and additional works at: https://epublications.marquette.edu/exsci_fac

Recommended Citation

Deering, Rita; Cruz, Meredith; Senefeld, Jonathon; Pashibin, Tatyana; Eickmeyer, Sarah; and Hunter, Sandra K., "Impaired Trunk Flexor Strength, Fatigability, and Steadiness in Postpartum Women." (2018). *Exercise Science Faculty Research and Publications*. 134.

https://epublications.marquette.edu/exsci_fac/134

Authors

Rita Deering, Meredith Cruz, Jonathon Senefeld, Tatyana Pashibin, Sarah Eickmeyer, and Sandra K. Hunter

Marquette University

e-Publications@Marquette

Exercise Sciences Faculty Research and Publications/College of Health Sciences

This paper is NOT THE PUBLISHED VERSION; but the author's final, peer-reviewed manuscript. The published version may be accessed by following the link in the citation below.

Medicine and Science in Sports and Exercise, Vol. 50, No. 8 (August 2018): 1558-1569. [DOI](#). This article is © Wolters Kluwer and permission has been granted for this version to appear in [e-Publications@Marquette](#). Wolters Kluwer does not grant permission for this article to be further copied/distributed or hosted elsewhere without the express permission from Wolters Kluwer.

Impaired Trunk Flexor Strength, Fatigability, and Steadiness in Postpartum Women

Rita E. Deering

Exercise Science Program, Department of Physical Therapy, Marquette University, Milwaukee, WI

Meredith Cruz

Department of Obstetrics and Gynecology, Medical College of Wisconsin, Milwaukee, WI

Jonathon W. Senefeld

Exercise Science Program, Department of Physical Therapy, Marquette University, Milwaukee, WI

Tatyana Pashibin

Exercise Science Program, Department of Physical Therapy, Marquette University, Milwaukee, WI

Sarah Eickmeyer

Department of Rehabilitation Medicine, University of Kansas, Kansas City, KS

Sandra K. Hunter

Exercise Science Program, Department of Physical Therapy, Marquette University, Milwaukee, WI

Abstract

Purpose To determine whether postpartum women (vaginal and cesarean delivery) have deficits in trunk flexor strength, fatigability and steadiness, compared with nulligravid women, up to 26 wk postpartum.

We *hypothesized* that postpartum women would be weaker, more fatigable, and have greater torque fluctuations than controls, with cesarean delivery showing greater deficits than vaginal delivery.

Methods Twenty-two control women (nulligravid) and 29 postpartum women (20–40 yr, 19 who delivered via vaginal birth, 13 via Caesarian section) participated. Postpartum women were tested 8 to 10 wk and 24 to 26 wk postpartum. Control women were tested 16 to 18 wk apart. Maximal voluntary isometric contractions (MVC) were performed at multiple trunk positions with the trunk flexor muscles. To determine trunk flexor fatigability, subjects performed intermittent isometric contractions at 50% MVC (6-s contraction, 4-s rest) in upright sitting until task failure. An MVC was performed during the fatiguing task (one per minute) and at 10 and 20 min of recovery.

Results At 8 and 26 wk, postpartum women (groups pooled) were weaker at all trunk angles (38% and 44% respectively, $P < 0.05$) than controls despite no differences in handgrip strength. Postpartum women were more fatigable (71% and 52% respectively) and had greater torque fluctuations than controls ($P < 0.05$). At 8 wk postpartum, women who had a cesarean delivery, were 59% more fatigable ($P = 0.004$) than the vaginal delivery group, with no difference between delivery types at 26 wk postpartum.

Conclusions Musculoskeletal recovery, including trunk flexor muscle strength and fatigability, is incomplete at 26 wk postpartum. These findings provide a rationale for future studies to address outcomes of rehabilitation programs specifically targeted at improving strength and fatigability of the trunk flexor muscles after pregnancy and childbirth.

Several physiological processes occur during pregnancy that impact the mother's musculoskeletal system. The abdominal muscles experience substantial stretch and increased inter-recti distance (IRD) commonly occurs ([1,2](#)). Hormones act on connective tissues throughout the mother's body, resulting in joint laxity, particularly in the pelvis ([3](#)). The loss of passive lumbopelvic joint stabilization increases the importance and role of muscular stabilization, which is provided by the core muscles, including the abdominal muscles ([4–6](#)). Appropriate function of the abdominal muscles is critical for several life functions ([5,7](#)). The ability to stabilize the pelvis against an external load was reported to be impaired during and after pregnancy ([8](#)). Despite the important daily and protective functions of the abdominal muscles, there is almost no understanding of the effects of pregnancy and mode of delivery on abdominal muscle function and recovery postpartum.

There are several other factors that may affect postpartum abdominal muscle function, including the mode of delivery and physical activity levels. Cesarean delivery involves profound disruption of the anterior abdominal wall and is associated with greater reports of postpartum pain and slower, more arduous recovery ([9](#)) than vaginal birth. Furthermore, physical activity, which can offset and protect against muscle dysfunction, is low for many mothers: approximately 60% of pregnant and 80% of postpartum women report a decline in their physical activity levels compared to prepregnancy activity ([10](#)). Given these factors and the reported changes in pain and recovery that suggest impairments in function of the core musculature in postpartum women, we investigated several aspects of abdominal muscle function including maximal strength at several trunk angles, fatigability, and force control (steadiness) during isometric trunk flexion.

Fatigability (often termed muscle fatigue) is an acute, activity-induced decline in the force or power of a muscle ([11](#)). It is often quantified as a reduction in the maximal force during or after a fatiguing task, but also can be quantified as the time to failure of a submaximal task ([11](#)). This metric of muscle function plays an important role in motor performance, injury prevention, and rehabilitation ([11](#)). However, fatigability of the abdominal muscles is rarely assessed clinically. Control of force, or steadiness of contraction, is also important for

stabilization during movement and for limb muscles it is associated with functional tasks ([12,13](#)). Steadiness is the ability to maintain a target force and can be quantified as the fluctuations in force around a target force as the SD of force or in relative terms as the coefficient of variation (CV) of force ([14](#)). In healthy men and nulligravid women, force fluctuations during trunk flexion contractions were threefold to fourfold larger than that reported for limb muscles and were similar between the sexes ([15](#)). Whether this force control of the trunk flexor muscles is altered after childbirth, or impacted by mode of delivery, is unknown.

Currently, the musculoskeletal system is not assessed as part of standard care in pregnant or postpartum women ([16–18](#)), and musculoskeletal impairments are often dismissed as “normal.” Paid maternity leave in the United States, when available, is typically 6 to 8 wk in duration ([19](#)) and often dictated by recovery of the smooth muscle of the uterus and perineal or surgical incision healing ([16,20](#)). Thus, we assessed abdominal function in postpartum women at approximately 8 wk postpartum, as well as at 26 wk postpartum, to determine longer term recovery of the abdominal muscles.

The primary purpose of this study was to determine function of the abdominal muscles after vaginal and cesarean delivery in postpartum women. The abdominal muscles are the primary movers during trunk flexion. Thus, maximal strength, fatigability, and steadiness of the trunk flexor muscles in women 8 to 10 wk and 24 to 26 wk postpartum were compared with nulligravid controls. Given the large disruption to abdominal muscles during pregnancy and childbirth, particularly in women who delivered via cesarean section, we *hypothesized* that 1) postpartum women (vaginal and cesarean delivery pooled) would demonstrate significant deficits in maximal strength across a range of trunk angles, increased fatigability, and decreased steadiness of contraction of the trunk flexor muscles up to 26 wk after delivery compared with healthy controls; and 2) women who underwent cesarean delivery would have greater deficits in strength, fatigability, and steadiness than those who delivered vaginally. We also assessed the influence of physical activity on abdominal function in postpartum women. Thus, we compared the trunk flexor function between a subgroup of postpartum women at 8 wk and control women who were matched for physical activity levels (estimated with accelerometer and questionnaire). We *hypothesized* that, even when matched for physical activity levels, postpartum women would demonstrate significant deficits in maximal strength, increased fatigability and less steady contractions compared with controls. As interpretive measures, we measured body composition, muscle thickness and IRD with ultrasonography, physical function, hand grip strength, and electromyography of the trunk flexor muscles during the fatiguing contraction.

METHODS

Participants

All participants were women age between 18 and 45 yr, not pregnant at the time of testing, free of chronic health conditions, did not smoke or use smokeless tobacco, had no known neurological impairment, did not take medications that impact neuromuscular excitability (including antidepressants), and had no medical or orthopedic contraindications to exercise. Control women had never been pregnant. All participants provided written informed consent. This study was approved by the institutional review boards at Marquette University and the Medical College of Wisconsin, and the Office of Clinical Research and Innovative Care Compliance at Froedtert Hospital.

Thirty-two postpartum women (vaginal delivery, $n = 19$; cesarean delivery, $n = 13$) and 22 control women participated in the study. Some control women were also included in a previously published article examining sex differences in trunk flexor function ([15](#)). Control women were also tested at two separate time points, separated by 16 to 18 wk.

Variables and Data Sources/Measurement

Trunk flexion torque

Maximal and submaximal trunk flexion torques were measured with a Biodex System 4 dynamometer (Biodex, Shirley, NY) using a Back Flexion-Extension attachment. Maximal voluntary isometric contractions (MVC) were assessed in six positions (upright sitting [0°]; 10°, 20°, 30°, and 40° of extension; 20° of flexion) to determine the strength of the trunk flexor muscles at different muscle lengths, to evaluate the impact of abdominal wall stretch during pregnancy on the torque–angle curve of the trunk flexor muscles after childbirth. Subjects were seated in the back flexion–extension attachment with the anterior superior iliac spine aligned with the axis of rotation of the device. The lower extremities and pelvis were secured with velcro straps, and two conjoined vertical straps on the anterior aspect of the shoulders restrained the trunk. Further details of device setup and subject positioning are described elsewhere ([15](#)). In brief, subjects performed trunk flexion, as if curling the shoulders toward the hips, without engaging the lower extremities. A study investigator (physical therapist) visually inspected all trunk flexion trials, and any trials with lower extremity involvement were excluded.

A minimum of three trials of MVC were performed at each position, with 1 min of rest between trials, until two trials were within 5% of each other. The maximum torque value was considered the MVC. Torque was recorded online using a Power 1401 A-D converter and Spike2 software [Cambridge Electronics Design (CED), Cambridge, UK]. Torque signals were digitized at 500 Hz and displayed on a monitor in front of the participant.

Hand grip strength

To determine the strength of a muscle group that is not typically affected by pregnancy and childbirth, and that is not dependent on the abdominal muscles or stability of the pelvic joints for function, bilateral handgrip strength was assessed as a control muscle group. Strength was measured with a JAMAR handgrip dynamometer (Patterson Medical, Warrenville, IL) with the arm positioned fully extended at the side. Three trials were performed bilaterally, and the highest value for each hand was used.

Fatigability

Fatigability of the trunk flexor muscles was assessed as the time to failure of an intermittent, submaximal isometric fatiguing task, and the reduction in MVC torque during the task. Subjects performed repeated trunk flexion contractions in upright sitting (0°) in the Biodex attachment at 50% MVC (target line displayed on a monitor) for 6 s, followed by 4 s of rest. A 6-s MVC was performed every minute (in lieu of 50% MVC), followed by 4 s of rest. Strong verbal encouragement was provided to continue until failure. Task failure was identified as torque <50% MVC for ≥3 s of the 6-s contraction or an MVC of <50% of baseline MVC. The fatiguing protocol always ended with an MVC. Recovery of maximal strength was assessed with an MVC at 10 and 20 min after task failure.

Steadiness (force fluctuations)

Steadiness of the trunk flexor muscles was quantified as the CV of torque during a 6-s isometric contraction at four contraction intensities (5%, 10%, 20%, and 70% of MVC) while the participant sat upright (0°) in the Biodex attachment. The CV allows comparison of steadiness across target intensities or between individuals of varying strength levels. A target line was displayed on a monitor in front of the participant, who was cued to trace the line as steadily as possible for the 6 s. Two trials were performed at each intensity with adequate rest provided between trials. Coefficient of variation of torque [(standard deviation of torque/mean torque produced) × 100] was calculated from the middle 3 s interval (1.5–4.5 s) of each 6 s trial, and the CV of the two trials at each intensity was averaged ([14](#)).

Steadiness was also assessed during the submaximal contractions (50% MVC) performed during the fatiguing exercise task; the middle 3-s interval was assessed from three submaximal contractions and averaged at the beginning and end of the fatiguing exercise task.

EMG

EMG signals were recorded from the right rectus abdominis, left external oblique, and right rectus femoris using two 8-mm silver chloride surface recording electrodes (Coulbourn Instruments, Whitehall, PA) arranged in a bipolar configuration and with recommended placements ([21](#)). EMG signals were amplified (1000×) and band-pass (13–1000 Hz) and Notch (60 Hz) filtered (Coulbourn Instruments, Allentown, PA). Signals were recorded online using a Power 1401 A-D converter (CED) and were digitized at 2000 Hz. EMG signals for the rectus abdominis and external oblique during the submaximal contractions of the fatiguing exercise protocol were normalized to the maximum RMS EMG of each respective muscle recorded during baseline trunk flexion MVC. EMG signals for the rectus femoris muscle during the submaximal trunk flexion contractions during the fatiguing protocol were normalized to the maximum RMS EMG of the rectus femoris recorded during knee extension MVC (obtained with a strap around the shank of the right lower extremity) before the fatiguing protocol of the trunk flexor muscles.

IRD and rectus abdominis muscle thickness

Real-time ultrasound (GE Vivide; 8 LRS linear probe) was used to assess IRD and thickness of the rectus abdominis muscles while the participant was supine. Measurement of IRD was made 4 cm below the umbilicus, because increased IRD below the umbilicus is usually more severe than increased IRD above the umbilicus ([4](#)). Muscle thickness measurements of the right rectus abdominis were taken 2.5 cm above and below the umbilicus.

Physical activity

Physical activity levels were quantified with accelerometry and questionnaire. Physical activity during a 4-d period at each time point of testing was quantified with ActiGraph accelerometers (ActiGraph Corp, Pensacola, FL) worn around the waist. Average minutes per day of moderate intensity exercise were quantified using ActiLife analysis software (ActiGraph). Self-reported physical activity was also estimated with a questionnaire, which estimated the metabolic cost (METs) and the weekly metabolic equivalents ($\text{MET} \cdot \text{h}^{-1} \cdot \text{wk}^{-1}$) of activities performed over the previous year ([22](#)).

Body composition

Dual x-ray absorptiometry was used to estimate fat mass (GE Lunar Idxa, GE Healthcare, Little Chalfont, United Kingdom).

Functional mobility

The 6-min walk test ([23](#)) was performed on an indoor course to quantify the maximal walk distance in 6 min.

Statistical Methods

Differences between groups (postpartum and control) were determined for subject characteristics, preexercise MVC torque, IRD, and time-to-task failure (fatigability) using independent samples *t* tests. Differences between groups for torque–angle curve, recovery of MVC torque after fatiguing exercise, EMG during fatiguing exercise, CV of the force fluctuations (steadiness) at multiple contraction intensities, and steadiness during fatiguing exercise were assessed using repeated-measures ANOVA with group as a between-subjects factor. Impact of method of delivery (cesarean vs vaginal) was assessed for subject characteristics, preexercise MVC torque, IRD, and time-to-task failure (fatigability) using independent samples *t* tests. Differences between delivery types for torque–angle curve, recovery of MVC torque after fatiguing exercise, EMG during fatiguing exercise, CV of the

force fluctuations (steadiness) at multiple contraction intensities, and steadiness during fatiguing exercise were assessed using repeated-measures ANOVA with delivery type as a between-subjects factor.

A subgroup of participants were matched for physical activity at the initial time point (8 wk postpartum) using both questionnaire and accelerometer estimations of physical activity levels. Differences between activity-matched controls and postpartum women were determined with similar methods to between group (postpartum and control) differences. Physical activity matching was not possible at the follow up time point (26 wk postpartum) due to very low numbers of activity matched individuals, resulting in insufficient statistical power.

Pearson correlation was used to examine relationships between scale level variables that demonstrated a linear relationship. Spearman correlation was used to examine relationships between variables that had a curvilinear relationship and/or were ordinal level data. Significance was set as $P < 0.05$. Missing data was excluded analysis by analysis. Statistical analyses were performed with SPSS version 24 software (IBM, Armonk, NY).

RESULTS

Participants

Twenty-nine postpartum women (vaginal delivery, $n = 18$; cesarean delivery, $n = 11$) and 15 control women completed initial (8–10 wk postpartum) testing. Twenty-eight postpartum women (vaginal delivery, $n = 17$; cesarean delivery, $n = 11$) and 14 control women completed testing at follow up (24–26 wk postpartum). Eight control women and 26 postpartum women (vaginal delivery $n = 17$, cesarean delivery $n = 9$) had complete data sets at both time points. Loss of follow-up for seven control women and two women from the cesarean group was due to schedule conflicts. One woman from the vaginal group did not complete follow up testing due to another pregnancy. Two women from the cesarean group completed testing only at 24 to 26 wk postpartum.

At the initial time point (8 wk postpartum), nine control women and nine postpartum women were matched for physical activity levels.

Descriptive Data

Group characteristics, including body composition, handgrip strength, muscle thickness, distance walked in 6 min, and physical activity, are presented for controls versus postpartum women in [Table 1](#) and vaginal delivery versus cesarean delivery in [Table 2](#). Postpartum women were older, weighed more, had greater body fat, and were less physically active than controls at both time points ($P < 0.05$). Hand grip strength was similar between control and postpartum women at both time points ($P > 0.05$). There were no differences in age, number of pregnancies, body composition, hand grip strength, or physical activity between delivery types at either 8 or 26 wk postpartum ($P > 0.05$).

Maximal Torque Across Different Positions (Torque–Angle Curve)

Postpartum women were weaker than controls across all trunk positions at 8 wk (39.6 ± 22.6 N·m vs 65.8 ± 29.2 N·m, respectively, $P < 0.001$; [Fig. 1A](#)) and 26 wk (36.5 ± 19.8 N·m vs 54.3 ± 24.2 N·m, $P = 0.001$; [Fig. 1B](#)) postpartum. The shape of the torque–angle curve was similar between groups at both time points (8 wk: position $\times$ group, $P = 0.927$; 26 wk: position $\times$ group, $P = 0.766$), with both control and postpartum women generating greatest trunk flexion MVC torque at 40° of extension (effect of position, $P < 0.001$ at both time points).

TABLE 1. Subject characteristics: control vs postpartum.

	Initial (8 wk Postpartum)		Follow-up (26 wk Postpartum)	
Variables	Control (n = 15)	Postpartum (n = 29)	Control (n = 14)	Postpartum (n = 28)
Age (yr)	25.5 ± 5.3	31.4 ± 5.2*	25.8 ± 6.1	32.0 ± 5.1*
Weight (kg)	63.6 ± 12.7	74.4 ± 13.4*	62.7 ± 7.8	71.4 ± 14.0*
Height (cm)	166.6 ± 7.1	164.1 ± 4.6	166.9 ± 8.4	164.1 ± 4.8
BMI (kg/m ²)	22.7 ± 3.8	27.6 ± 5.0*	22.4 ± 1.8	26.9 ± 5.1*
Body fat (%)	31.0 ± 5.4	38.5 ± 6.9*	31.8 ± 4.6	36.7 ± 8.4*
Thickness of right rectus abdominis muscle belly (2.5 cm above umbilicus) (cm)	1.0 ± 0.2	0.8 ± 0.2*	1.0 ± 0.1 (n = 13)	0.8 ± 0.2*
Thickness of right rectus abdominis muscle belly (2.5 cm below umbilicus) (cm)	1.0 ± 0.2	0.8 ± 0.2*	0.9 ± 0.1 (n = 13)	0.8 ± 0.2
Average minutes per day of moderate intensity physical activity	47.8 ± 23.4 (n = 9)	18.8 ± 18.8* (n = 20)	31.2 ± 14.0 (n = 7)	16.7 ± 10.5* (n = 13)
Self-reported physical activity over the previous 12 months (MET/hj1lwj1)	43.0 ± 28.3 (n = 14)	22.2 ± 19.2* (n = 27)	34.0 ± 25.1	14.9 ± 17.3* (n = 25)
Distance walked in 6 min (m)	686.2 ± 58.7	640.0 ± 64.8*	693.3 ± 58.2	652.7 ± 65.1
Hand grip maximal strength, right hand (kg)	33.4 ± 6.7	33.1 ± 5.9	34.2 ± 5.3	34.1 ± 6.5 (n = 27)
Hand grip maximal strength, left hand (kg)	31.4 ± 5.9	31.1 ± 4.9	30.8 ± 4.5	32.3 ± 5.6 (n = 27)
Change in MVC torque at end of fatiguing exercise protocol (% MVC)	-26.2 ± 15.9	-14.6 ± 32.4	-30.0 ± 13.4	-24.0 ± 18.8
External oblique EMG at beginning of fatigue task (%MVC)	48.3 ± 33.	7 54.2 ± 23.9	46.6 ± 16.4	50.3 ± 23.0
External oblique EMG at end of fatigue task (%MVC)	50.2 ± 32.5	60.6 ± 20.8	56.6 ± 17.4	56.2 ± 23.0
Rectus femoris EMG at beginning of fatigue task (%MVC)	12.8 ± 27.6	4.5 ± 2.4	4.7 ± 3.7	6.4 ± 6.0
Rectus femoris EMG at end of fatigue task (%MVC)	12.4 ± 26.9	4.5 ± 2.4	4.8 ± 3.8	6.5 ± 6.0

*P < 0.05.

TABLE 2. Subject characteristics: vaginal vs cesarean delivery.

	8 wk Postpartum		26 wk Postpartum	
Variables	Vaginal (n = 18)	Cesarean (n = 11)	Vaginal (n = 17)	Cesarean (n = 11)
Age (yr)	30.6 ± 6.0	32.8 ± 3.5	31.4 ± 5.8	33.0 ± 4.0
Weeks postpartum	9.4 ± 1.1	9.3 ± 0.5	26.1 ± 1.6	26.2 ± 0.6
Total no. pregnancies	2.1 ± 1.2 (n = 17)	3.0 ± 1.9 (n = 10)	2.1 ± 1.1 (n = 15)	3.0 ± 2.1 (n = 9)
Duration of most recent pregnancy (wk)	39.3 ± 1.0 (n = 14)	38.1 ± 1.2* (n = 10)	39.3 ± 1.0 (n = 12)	38.3 ± 1.4 (n = 9)
Weight (kg)	74.3 ± 14.4	74.2 ± 12.2	70.9 ± 15.3	72.4 ± 12.4
Height (cm)	164.6 ± 4.3	163.1 ± 5.3	165.1 ± 4.8	162.6 ± 4.8
BMI (kg/m ²)	27.5 ± 5.4	27.8 ± 4.3	26.4 ± 5.6	27.5 ± 4.3
Body fat %	39.0 ± 7.0	37.7 ± 7.0	36.1 ± 8.8	37.7 ± 8.2
Thickness of right rectus abdominis muscle belly (2.5 cm above umbilicus) (cm)	0.8 ± 0.2	0.8 ± 0.1	0.8 ± 0.2	0.8 ± 0.1

Thickness of right rectus abdominis muscle belly (2.5 cm below umbilicus) (cm)	0.8 ± 0.2	0.8 ± 0.2	0.8 ± 0.3	0.8 ± 0.2
Average minutes per day of moderate intensity physical activity	17.1 ± 19.1 (n = 10)	20.4 ± 19.4 (n = 10)	16.8 ± 11.2 (n = 10)	13.1 ± 4.9 (n = 3)
Self-reported physical activity over the previous 12 months (METIhj1lwkj1)	22.0 ± 16.3 (n = 17)	22.6 ± 24.3 (n = 10)	15.4 ± 20.8 (n = 15)	14.3 ± 11.3 (n = 10)
Distance walked in 6 min (m)	645.0 ± 62.0	631.8 ± 71.3	663.4 ± 60.8	636.1 ± 70.9
Hand grip maximal strength, right hand (kg)	33.4 ± 5.9	32.5 ± 6.1	34.9 ± 6.5	32.7 ± 6.6 (n = 10)
Hand grip maximal strength, left hand (kg)	31.7 ± 5.2	30.2 ± 4.6	33.2 ± 5.9	30.7 ± 4.9 (n = 10)
Change in MVC torque at end of fatiguing exercise protocol (% MVC)	-12.0 ± 27.4 (n = 16)	-18.4 ± 39.7	-22.9 ± 21.0	-25.8 ± 15.4
External oblique EMG at beginning of fatigue task (%MVC)	52.2 ± 26.7 (n = 14)	57.4 ± 19.7 (n = 9)	46.8 ± 22.6	55.7 ± 23.7
External oblique EMG at end of fatigue task (%MVC)	59.4 ± 23.5 (n = 14)	62.3 ± 16.9 (n = 9)	49.2 ± 16.8	67.0 ± 27.6
Rectus femoris EMG at beginning of fatigue task (%MVC)	4.3 ± 2.1 (n = 14)	4.8 ± 2.9 (n = 9)	5.9 ± 5.1	7.3 ± 7.4
Rectus femoris EMG at end of fatigue task (%MVC)	4.3 ± 2.1 (n = 14)	4.9 ± 2.9 (n = 9)	6.0 ± 5.2	7.3 ± 7.4

*P < 0.05.

FIGURE 1

Torque–angle curve for the trunk flexor muscles at the initial and follow up time points for postpartum vs control (A and B) and vaginal vs cesarean delivery (C and D). Negative numbers indicate positions of trunk extension, upright sitting is 0°, and positive numbers indicate positions of trunk flexion. Postpartum women had lower maximal trunk flexion torque across all positions at both time points compared to control. However, the torque–angle curve shape was similar for all groups. There were no differences between delivery types (vaginal vs cesarean). *Group difference at the time point (postpartum vs control).

Mode of Delivery

There was no difference in the torque–angle curve between delivery types at 8 wk (position: $P < 0.001$; position $\times$ delivery type: $P = 0.169$; delivery type: $P = 0.058$; [Fig. 1C](#)) or 26 wk (position: $P < 0.001$, position $\times$ delivery type: $P = 0.964$, delivery type: $P = 0.485$; [Fig. 1D](#)).

Comparison Across Matched Activity Levels

Despite being matched for physical activity levels, postpartum women demonstrated decreased torque production compared to controls at all trunk positions ($P = 0.002$). The shape of the torque–angle curve was similar for both groups (position: $P = 0.001$, position $\times$ group: $P = 0.826$).

Fatigability

Time-to-task failure

Postpartum women had a shorter time-to-task failure (i.e. greater fatigability) than control women at 8 wk postpartum (189 ± 156 s vs 644 ± 327 s, respectively; $P < 0.001$; [Fig. 2A](#)). By 26 wk, postpartum women demonstrated longer time-to-task failure (i.e., improved fatigability) versus 8 wk postpartum ($P = 0.015$), but were still more fatigable than control women (288 ± 167 s vs 605 ± 396 s; group, $P = 0.011$; [Fig. 2B](#)).

FIGURE 2 Fatigability and recovery. (A–B) Time-to-task failure for the intermittent isometric submaximal trunk flexion fatiguing exercise at time points for postpartum (pooled), control, vaginal delivery, and cesarean delivery at 8 to 10 wk postpartum [A] and 24 to 26 wk postpartum[B]. *Group difference at the time point (postpartum vs control); γ Improvement compared to initial timepoint; #Delivery group difference at the time point (vaginal vs cesarean). (C–D) MVC torque before (baseline), immediately after (task failure, TF), and 10 min (R10) and 20 min (R20) after fatiguing trunk flexor exercise for controls vs postpartum women at 8 to 10 wk (C) and 24–26 wk (D). Postpartum women generate lower maximal torque than control women at all time points. Postpartum women demonstrate impaired recovery of MVC strength up to 10 wk postpartum (C), but show a similar recovery pattern to control women at 26 wk postpartum (D). *Group difference at the time point (postpartum vs control); δ Different from baseline MVC. (E–F) No statistical difference is present between modes of delivery at either time point (E and F). Postpartum women demonstrate a decline in baseline MVC strength from 8 to 26 wk, driven by a loss of strength for women in the vaginal delivery group. δ Different from baseline MVC.

Mode of delivery

At 8 wk postpartum, the cesarean delivery group demonstrated a shorter time-to-task failure (greater fatigability) than the vaginal delivery group (99 ± 58 s vs 244 ± 173 s; $P = 0.004$; [Fig. 2A](#)). By 26 wk postpartum, there was no difference in time-to-task failure between delivery groups (262 ± 168 s vs 304 ± 169 s; $P = 0.523$; [Fig. 2B](#)). The vaginal delivery group showed no change in time-to-task failure from 8 to 26 wk ($P = 0.306$), but the cesarean delivery group demonstrated improvement in time-to-task failure ($P = 0.005$).

Comparison across matched activity levels

In this subgroup, the postpartum women continued to demonstrate increased trunk flexor fatigability compared with controls (195.2 ± 183.7 s vs 618.7 ± 332.1 s, respectively; $P = 0.004$; [Table 3](#)).

TABLE 3. Physical activity matched participants at initial time point (8 wk postpartum).

Variables	Control (n = 9)	Postpartum (n = 9)
Age (yr)	23.3 ± 1.8	$31.6 \pm 4.9^*$
Weight (kg)	63.9 ± 10.5	70.0 ± 12.3
Body fat %	29.9 ± 4.3	$36.7 \pm 6.7^*$
BMI (kg/m^2)	22.5 ± 1.5	26.2 ± 5.1
Rectus abdominis muscle thickness (above umbilicus) (cm)	1.1 ± 0.2	$0.8 \pm 0.2^*$
Rectus abdominis muscle thickness (below umbilicus) (cm)	1.1 ± 0.1	$0.7 \pm 0.2^*$
Sedentary time (average minutes per day)	531.0 ± 118.9	494.8 ± 80.1
Light physical activity (average minutes per day)	240.0 ± 34.2	276.5 ± 52.4
Moderate physical activity (average minutes per day)	50.5 ± 25.9	24.3 ± 29.3
Vigorous physical activity (average minutes per day)	13.9 ± 7.0	7.5 ± 9.2

Average steps per day	11509.0 ± 2850.5	6649.5 ± 4292.4
Distance walked in 6 min (m)	697.0 ± 39.9	657.6 ± 78.9
Self-reported physical activity over the previous 12 months (METIh1Iwkj1)	57.3 ± 25.1	45.4 ± 13.5
Time-to-task failure on the intermittent, isometric trunk flexion fatiguing exercise protocol (s)	618.7 ± 332.1	195.2 ± 183.7
Preexercise MVC (NIm)	55.5 ± 28.9	26.9 ± 14.2*
CV of torque at beginning of fatigue task (%)	5.2 ± 2.0%	11.9 ± 6.6%*
CV of torque at end of fatigue task (%)	8.4 ± 4.1	12.3 ± 4.5%*

*P G 0.05.

Maximal trunk flexion torque before and after the fatiguing task

Trunk flexion MVC torque was less for the postpartum women than control women at both 8 wk (28.6 ± 12 N·m vs 46.5 ± 26.1 N·m; $P = 0.022$; [Fig. 2C](#)) and 26 wk postpartum (24.6 ± 10.1 N·m vs 44.1 ± 16.6 N·m; $P = 0.001$; [Fig. 2D](#)).

At task failure, the postpartum and control women had a similar reduction in MVC torque, at both the 8- and 26-wk time points ([Table 1](#)). At 8 wk, the MVC torque of postpartum women was below preexercise MVC at 20 min after fatiguing exercise, whereas the MVC torque of control women had recovered by 10 min postexercise (time, $P < 0.001$; time $\times$ group, $P < 0.001$; group, $P = 0.001$; [Fig. 2C](#)). At 26 wk, postpartum women demonstrated a recovery (increase) of MVC torque postexercise that was similar to control women, though postpartum women demonstrated lower MVC torque than control women at all time points before and after exercise (time, $P < 0.001$; time $\times$ group, $P = 0.463$; group, $P < 0.001$; [Fig. 2D](#)).

Mode of delivery

There were no differences between vaginal and cesarean delivery for preexercise MVC strength at 8 wk (31.4 ± 13.0 N·m vs 24.0 ± 8.8 N·m, respectively, $P = 0.106$, [Fig. 2E](#)) nor at 26 wk (24.9 ± 11.4 N·m vs 24.3 ± 8.1 N·m, respectively, $P = 0.886$, [Fig. 2F](#)). There was also no difference in MVC torque recovery after fatiguing exercise at 8 wk (time, $P = 0.775$; time $\times$ delivery type, $P = 0.592$; delivery type, $P = 0.829$; [Fig. 2E](#)) and 26 wk (time, $P = 0.006$; time $\times$ delivery type, $P = 0.163$; delivery type, $P = 0.386$; [Fig. 2F](#)).

Comparison across matched activity levels

In the subgroup of physical activity–matched women, postpartum women continued to demonstrate decreased trunk flexor strength compared to control women at 8 wk postpartum (26.9 ± 14.2 N·m vs 55.5 ± 28.9 N·m, respectively; $P = 0.021$; [Table 3](#)). Despite being matched for physical activity levels, postpartum women continued to demonstrate impaired recovery of MVC strength after fatiguing exercise (time, $P = 0.027$; time $\times$ group, $P = 0.019$; group, $P = 0.013$), with MVC strength at 20 min postexercise still below baseline MVC.

Steadiness of contraction at 5%, 10%, 20%, and 70% of MVC

At 8 wk postpartum, the CV of torque at four intensities was quantified for 22 postpartum women and 13 control women. Coefficient of variation of torque decreased as contraction intensity increased for both postpartum and control women (intensity, $P < 0.001$; intensity $\times$ group, $P = 0.068$; [Fig. 3A](#)). Postpartum women, however, had larger CV of torque than controls (group, $P = 0.013$), particularly at low contraction intensities. At 26 wk postpartum, CV of torque was obtained for 25 postpartum women and 15 control women. As was seen at initial testing (8 wk postpartum), CV of torque decreased as contraction intensity increased for both groups (intensity, $P < 0.001$; intensity $\times$ group, $P = 0.219$; [Fig. 3B](#)). Postpartum women continued to have larger CV of torque than controls at 26 wk postpartum (group, $P = 0.039$), with the greatest difference between groups at low contraction intensities.

FIGURE 3 Steadiness. (A–B) Steadiness (CV of torque) of trunk flexor contractions at four intensities (5%, 10%, 20%, and 70% MVC) for control vs postpartum women at 8 to 10 wk [A] and 24 to 26 wk [B] postpartum. Postpartum women have greater torque fluctuations than controls at both time points, particularly at low contraction intensities. * = group difference at the time point (postpartum vs control). (C–D) CV of torque during submaximal (50% MVC) contractions during fatiguing exercise is represented for controls vs postpartum women at 8 to 10 wk (C) and 24 to 26 wk (D). Both groups demonstrated increased force fluctuations from the beginning to end of fatiguing exercise at both time points. Coefficient of variation of torque is similar between groups at 8 to 10 wk postpartum (C), but postpartum women have greater fluctuations in torque than controls at 24 to 26 wk postpartum (D). *Group difference at the time point (postpartum vs control).

Comparison across matched activity levels

Coefficient of variation of torque at four intensities was quantified for eight control and eight postpartum women who were matched for physical activity levels. Coefficient of variation of torque decreased as contraction intensity increased (intensity $P < 0.001$). Postpartum women had greater fluctuations in torque than controls (group, $P = 0.020$) despite being matched for physical activity levels. However, the decrease in CV with higher contraction intensity was greater in postpartum women than controls (intensity $\times$ group, $P = 0.049$) because postpartum women had a significantly higher CV at 5% MVC but not at the higher contraction intensities.

Method of delivery

At 8 wk postpartum, CV was quantified at four intensities for 13 women who had a vaginal delivery and nine women who delivered via cesarean section. Coefficient of variation of torque decreased as intensity increased for both delivery types (intensity $P < 0.001$, intensity $\times$ delivery type $P = 0.202$). Coefficient of variation of torque was similar between delivery types (delivery type $P = 0.846$). At 26 wk postpartum ($n = 15$ vaginal delivery, $n = 10$ cesarean delivery), the CV of torque decreased as contraction intensity increased (intensity, $P < 0.001$; intensity $\times$ delivery type, $P = 0.206$) but was similar between delivery types (delivery type, $P = 0.637$).

Steadiness of contraction during fatiguing exercise

Coefficient of variation of torque during the fatiguing exercise protocol (50% MVC) was quantified at the initial time point (8 wk postpartum) in 15 control women and 23 postpartum women. Coefficient of variation of torque increased similarly across time for both control and postpartum women (time, $P < 0.001$; time $\times$ group, $P =$

0.316; [Fig. 3C](#)) with no group differences (group, $P = 0.180$). At the follow up time point (26 wk postpartum), 14 control women and 28 postpartum women were included in analysis. Coefficient of variation of torque increased across time for both groups (time, $P < 0.001$; time $\times$ group, $P = 0.122$; [Fig. 3D](#)), but postpartum women were less steady (larger CV of torque) than control women (group, $P = 0.026$).

Comparison across matched activity levels

Coefficient of variation of torque during the fatiguing task ($n = 9$ control, $n = 6$ postpartum women) increased with time for both groups (time, $P = 0.036$; time $\times$ group, $P = 0.088$). Despite being matched for physical activity levels, postpartum women were less steady (larger CV of torque) than controls (group, $P = 0.028$) at the beginning of the fatiguing task ($11.9\% \pm 6.6\%$ vs $5.2\% \pm 2.0\%$, respectively) and the end of the fatiguing task ($12.3\% \pm 4.5\%$ vs $8.4\% \pm 4.1\%$, respectively) ([Table 3](#)).

Method of delivery

At 8 wk postpartum, 14 women who delivered vaginally and nine women who delivered via cesarean section were included in analysis. Coefficient of variation of torque increased across time (time, $P = 0.001$), with women who underwent cesarean delivery demonstrating a greater increase in CV of torque across time (time $\times$ delivery type, $P = 0.032$). Coefficient of variation of torque was similar between delivery types (delivery type, $P = 0.146$).

At 26 wk postpartum ($n = 17$ vaginal delivery and $n = 11$ cesarean delivery), CV of torque increased across time for both delivery types (time $P < 0.001$; time $\times$ delivery type, $P = 0.929$) and was similar between delivery types (delivery type, $P = 0.511$).

EMG during fatiguing exercise (%MVC)

EMG activity (%MVC) of the external oblique and rectus abdominis increased over time for both postpartum and control women (external oblique: time, $P = 0.025$; time $\times$ group, $P = 0.229$; rectus abdominis: time, $P = 0.037$; time $\times$ group, $P = 0.283$) with both groups demonstrating similar activation of both muscles at the initial time point, that is, 8 wk postpartum (group, $P = 0.361$ for external oblique, $P = 0.380$ for rectus abdominis; [Table 1](#)).

At the follow-up time point (26 wk postpartum), external oblique EMG increased over time for both groups (time, $P = 0.003$; time $\times$ group, $P = 0.409$), with no difference in relative EMG between control and postpartum women (group $P = 0.800$; [Table 1](#)). Rectus abdominis EMG did not change over time for either group (time, $P = 0.138$; time $\times$ group, $P = 0.231$), and relative rectus abdominis EMG was similar between control and postpartum women (group $P = 0.188$).

At both time points (8 and 26 wk postpartum), rectus femoris EMG did not change across time during the fatiguing task in the control or postpartum women (8 wk: time, $P = 0.351$; time $\times$ group, $P = 0.376$; 26 wk: time, $P = 0.410$; time $\times$ group, $P = 0.725$), and both groups demonstrated similar rectus femoris activation (8 wk: group $P = 0.164$; 26 wk: group $P = 0.328$; [Table 1](#)).

Method of delivery

At both time points (8 and 26 wk postpartum), there were no differences in relative EMG between vaginal and cesarean delivery groups for any muscle groups tested (time $\times$ group, $P > 0.05$, group, $P > 0.05$ for all muscles).

IRD

Postpartum women demonstrated greater IRD 4 cm below the umbilicus than controls at 8 wk (1.4 ± 1.1 cm vs 0.4 ± 0.2 cm, $P < 0.001$) and 26 wk (1.3 ± 1.1 cm vs 0.3 ± 0.2 cm, $P < 0.001$).

Comparison across matched activity levels

Postpartum women demonstrated greater IRD than activity-matched controls (1.2 ± 0.8 cm vs 0.4 ± 0.2 cm, respectively; $P = 0.019$; [Table 3](#)).

Mode of delivery

There was no difference in IRD between vaginal and cesarean delivery types at 8 wk (1.2 ± 1.1 cm vs 1.8 ± 1.2 cm, respectively, $P = 0.271$) or 26 wk (1.2 ± 1.0 cm vs 1.5 ± 1.3 cm, respectively, $P = 0.431$).

Associations (Postpartum and Control)

Trunk flexor MVC torque was associated with fatigability (8 wk: $r = 0.602$, $P < 0.001$ and 26 wk: $r = 0.415$, $P = 0.006$), thickness of the rectus abdominis muscle belly 2.5 cm above the umbilicus (8 wk: $r = 0.311$, $P = 0.040$ and 26 wk: $r = 0.388$, $P = 0.012$), self-reported physical activity (8 wk: $r = 0.430$, $P = 0.005$) and the 6-min walk (8 wk: $r = 0.306$, $P = 0.044$ and 26 wk: $r = 0.449$, $P = 0.003$). Thus, stronger women had a longer time-to-task failure, greater rectus abdominis thickness, were more active, and walked further in 6 min.

Fatigability was associated with IRD (8 wk: $r_s = -0.548$, $P < 0.001$ and 26 wk: $r_s = -0.484$, $P = 0.002$), such that women with greater IRD were more fatigable.

DISCUSSION

The novel findings are that the trunk flexor muscles of postpartum women were weaker, more fatigable, and demonstrated greater fluctuations in torque during isometric contractions than nulligravid women (controls), and that these impairments in trunk flexor muscle function persisted at 6 months (26 wk) after childbirth. The strength deficits do not appear to impact muscle groups throughout the body, because there was no difference in hand grip strength between controls and postpartum women. Fatigability of the trunk flexor muscles was greater in postpartum women at 8 wk postpartum, even when matched with the control women for physical activity levels. Furthermore, mode of delivery impacted abdominal muscle function: women whom had a cesarean delivery were more fatigable, and had a greater increase in force fluctuations during the fatiguing trunk flexor exercise, at 8 wk postpartum compared with women who delivered vaginally. Greater trunk flexor strength was associated with lower fatigability and greater functional mobility. Thus, trunk flexor muscle function was substantially lower in postpartum women than healthy controls, with limited recovery by 6 months.

This is the first study to establish that postpartum women demonstrate impaired strength, fatigability, and torque steadiness of core muscles, which are important to many daily activities, even at 6 months postpartum. The strength deficits in postpartum women, relative to controls, were 38% at 8 wk, and 44% at 26 wk; long past when women have returned to work (~8 wk postpartum). The strength deficits occurred at all trunk angles, so that the stretch experienced by the abdominal muscles during pregnancy did not alter the torque angle curve. The highest torque occurred at the -40° position for both postpartum and control women, a position close to full extension as occurs in upright standing and where the abdominal muscle fibers must be close to the optimal overlap of actin and myosin in the sarcomere and optimal force production ([24](#)). Furthermore, strength deficits in postpartum women in this study were localized to the trunk flexor muscles because the handgrip strength of the postpartum and control women was similar.

Postpartum women (both vaginal and Caesarean delivery groups) were also more fatigable (shorter time-to-task failure) than controls by 71% at 8 wk and by 52% at 26 wk postpartum. Another metric of fatigability was the reduction in the MVC which occurred more rapidly for the postpartum women than the control women during the fatiguing task: the reduction in MVC torque was similar between groups, but the time to failure was considerably briefer in the postpartum women than controls. Furthermore, at 20 min of recovery from the

fatiguing task at 8 wk after delivery, the MVC differed between groups, with postpartum women demonstrating impaired *recovery* of maximal strength. In contrast, control women had returned to baseline strength 10 min after exercise. This prolonged reduction in maximal strength may increase risk of injury when performing repetitive tasks, such as lifting and carrying.

Method of delivery was a contributing factor to the increased fatigability of postpartum women at 8 wk after delivery, with a briefer time to failure for the cesarean delivery group compared with vaginal delivery ([Fig. 2A](#)). The recovery of the MVC torque did not differ with delivery mode. These results are in part consistent with our hypothesis and support that the surgical damage to the abdominal muscles and the surrounding fascia slow recovery of abdominal muscle function, particularly for activities involving repeated contractions and fatiguing exercise. Fatigability was also greater in women matched for physical activity levels indicating that the abdominal muscle endurance was still lower independent of physical activity.

Fatigability did not differ with delivery mode by 26 wk because the women who had undergone cesarean delivery had improved abdominal muscle endurance to similar levels as the vaginal delivery group. In another study, women who had a cesarean delivery reported a more difficult postpartum recovery than women who delivered vaginally, along with greater pain reports in the immediate postpartum period (37). We showed here that trunk flexor strength did not differ at 8 wk postpartum between cesarean and vaginal delivery, so fatigability may be a more sensitive test of abdominal muscle recovery, at least at 8 wk postpartum. No differences exist, however, in the length of paid maternity leave based on delivery method ([3](#)). This research suggests that women who undergo cesarean delivery require a longer period to recover from the musculoskeletal deficits experienced from pregnancy and childbirth than women who experience a vaginal delivery. Given our findings and the severe deficits in abdominal function of postpartum women in general, it appears all postpartum women, regardless of delivery method, would benefit from rehabilitative interventions; however, this rehabilitation may be even more crucial in the immediate postpartum period for women who deliver by cesarean section.

EMG activity (% MVC) of the trunk flexor muscles was similar throughout the fatiguing contraction between postpartum and control women, suggesting similar muscle activation between groups. Similarly, EMG was not different between vaginal and cesarean delivery during the fatiguing contraction. Rectus femoris activation was low for all groups (4%–13%) and did not change during the fatiguing exercise task, indicating that the rectus femoris was not contributing to the observed trunk flexion torque or compensating during the fatiguing contraction. Activation of the abdominal muscles (rectus abdominis and external oblique) increased from the beginning to the end of the fatiguing exercise bout, which is expected, and likely represents an increase in motor unit recruitment and altered discharge rates as the active muscle fibers become fatigued ([25,26](#)). Despite the shorter time to failure of the postpartum women, the relative muscle activity at task failure was similar to controls, and EMG was similar between delivery types. These results suggest that, for the rectus abdominis and external oblique muscles, the activation of the motor neuron pool in postpartum women was similar to controls at task end. Hence, impaired muscle activation probably did not contribute significantly to the difference in fatigability between the postpartum and control women. Alternatively, muscular mechanisms likely played a greater role than neural mechanisms for the increased fatigability of the postpartum women in this study, although this requires confirmation.

Postpartum women also demonstrated larger fluctuations in trunk flexor torque (lower steadiness) than control women, with women who underwent cesarean delivery exhibiting a greater increase in force fluctuations during fatiguing exercise than women who delivered vaginally. Steadiness can be impacted by several factors, including common drive to the motor neuron pool ([14,27,28](#)), physical activity ([14](#)), and stress ([29](#)). The trunk flexor muscles are innervated by multiple spinal levels ([4](#)), which may contribute to higher force fluctuations during trunk flexion contractions than those observed in limb muscles, as has been shown in healthy men and nulligravid

women ([15](#)). Low physical activity levels during and after pregnancy ([10](#)) may also impact force steadiness in this population; however, postpartum women continued to demonstrate greater force fluctuations than controls even when matched for physical activity levels, suggesting that physical activity is not the sole mechanism behind the impaired force control observed during trunk flexion contractions after pregnancy and childbirth. Furthermore, perturbations to the connective tissue of the anterior abdominal wall during pregnancy and childbirth, such as thinning and widening of the linea alba ([12](#)), may compromise the transfer of force generated by the abdominal muscles ([30,31](#)). Large fluctuations in abdominal muscle force may lead to significant fluctuations in intra-abdominal pressure, and thus stability of the lumbar spine and pelvis ([5](#)), making further research of force control and its impact on lumbopelvic stability important in postpartum women.

Trunk flexor strength appears to be a critical factor in musculoskeletal function after pregnancy and childbirth. Stronger women were less fatigable and demonstrated better functional mobility, as evidenced by performance on the 6-min walk test. The association between strength and fatigability of the abdominal muscles has also been observed in healthy young men and women ([15](#)), supporting the importance of core strength. Strength was also associated with thickness of the rectus abdominis muscle and with self-reported physical activity over the previous year. Thus, our results suggest that lower physical activity levels in postpartum women may have contributed to lower trunk flexor strength. Physical activity levels of postpartum women were significantly lower than those of control women, and self-reported physical activity over the previous 12 months declined even further between 8 and 26 wk postpartum. In addition, postpartum women did not meet the recommended average of 30 min·d⁻¹ of moderate intensity physical activity ([32](#)). These results provide a rationale for prescription of individualized exercise for postpartum women to offset the strength decrements, and thus the impairments in fatigability and functional mobility, seen in this study.

However, the strength, fatigability, and torque steadiness deficits after pregnancy and childbirth do not appear to be mediated solely by decreased physical activity levels in the postpartum group, as physical activity matching continued to show that postpartum women were weaker, more fatigable, with reduced control of torque during isometric contractions than control women. In this activity-matched subgroup, the postpartum women continued to demonstrate greater IRD and thinner rectus abdominis muscle thickness than controls, suggesting that the significant stretch experienced by the abdominal muscles during pregnancy and the changes in fascial integrity in postpartum women may contribute to the deficits in abdominal muscle function observed in this population. This highlights the critical need for further research into the mechanisms responsible for impaired strength, fatigability, and steadiness, especially of the trunk flexor muscles, after pregnancy and childbirth.

One limitation of this study is that the postpartum women in this study were statistically older than the women in the control group; however, it has been well documented that age-related changes in muscular strength typically are not apparent until after age 50 yr ([28](#)). As the mean age of both groups is in the mid-20s and early 30s (~5–6 yr difference), age was probably not a significant contributing factor to the differences in strength, fatigability and steadiness between nulligravid and postpartum women. Another limitation is that we only assessed one muscle group outside of the trunk flexors (hand grip muscles). Further research is needed to determine if other major muscle groups, such as the knee extensors, also experience impairments in strength/fatigability in a postpartum population.

CONCLUSIONS

Postpartum women had larger deficits in trunk flexor strength, muscle endurance and steadiness of contraction, up to 6 months postpartum, and that could not be fully accounted for by lower levels of physical activity, at least at 8 wk after childbirth. Furthermore, women who delivered by cesarean section had even greater trunk flexor deficits in fatigability and torque steadiness than those who had a vaginal delivery. These large deficits in trunk flexor strength, fatigability, and steadiness in postpartum women in this study support the need for skilled

assessment of the musculoskeletal system by health care providers who have expertise in the examination, diagnosis, and treatment of musculoskeletal disorders. Incorporating health care professionals, such as physiatrists and physical therapists, in the standard care of pregnant and postpartum women may offset some of the neuromuscular impairments seen in this study and may improve the incidence and severity of other pregnancy-related musculoskeletal disorders, such as low back pain, pelvic girdle pain, and incontinence. Further research is needed to determine the mechanisms for the large deficits in abdominal muscle function postpartum and the impact of exercise and rehabilitation on the musculoskeletal health of pregnant and postpartum women.

The authors thank the following individuals for their contributions to this project: Ms. Jordan Lemens, Marquette University, Exercise Science Intern, for assistance with data collection and analysis. Ms. Sarah Nisivaco, Marquette University, Honors Program Summer Research Program, for assistance with data analysis. Dr. Anna McCormick, Medical College of Wisconsin, for assistance with subject recruitment. Dr. Ruta Brazauskas, Medical College of Wisconsin, for statistical analysis.

Women's Health Research Program Grant, Department of Obstetrics and Gynecology, Medical College of Wisconsin; American Dissertation Fellowship, American Association of University Women; National Center for Advancing Translational Sciences, National Institutes of Health, through grants UL1TR001436 and 1TL1TR001437.

The authors report no conflicts of interest. The results of this study are presented clearly, honestly, and without fabrication, falsification, or inappropriate data manipulation. The results of this study do not constitute endorsement by ACSM, are solely the responsibility of the authors, and do not necessarily represent the official views of the NIH.

REFERENCES

1. Boissonnault JS, Blaschak MJ. Incidence of diastasis recti abdominis during the childbearing year. *Phys Ther*. 1988;68(7):1082–6.
2. Coldron Y, Stokes MJ, Newham DJ, Cook K. Postpartum characteristics of rectus abdominis on ultrasound imaging. *Man Ther*. 2008;13(2):112–21.
3. Kristiansson P, Svärdsudd K, von Schoultz B. Reproductive hormones and aminoterminal propeptide of type III procollagen in serum as early markers of pelvic pain during late pregnancy. *Am J Obstet Gynecol*. 1999;180(1):128–34.
4. Neumann DA. *Kinesiology of the Musculoskeletal System: Foundations for Rehabilitation*. Third ed. St. Louis, MO: Mosby Elsevier; 2016.
5. Hodges PW, Kaigle Holm A, Holm S, et al. Intervertebral stiffness of the spine is increased by evoked contraction of transversus abdominis and the diaphragm: in vivo porcine studies. *Spine (Phila Pa 1976)*. 2003;28(23):2594–601.
6. Stokes IA, Gardner-Morse MG, Henry SM. Abdominal muscle activation increases lumbar spinal stability: analysis of contributions of different muscle groups. *Clin Biomech (Bristol, Avon)*. 2011;26(8):797–803.
7. Lee DG, Lee LJ, McLaughlin L. Stability, continence and breathing: the role of fascia following pregnancy and delivery. *J Bodyw Mov Ther*. 2008;12:333–48.
8. Gilleard WL, Brown JM. Structure and function of the abdominal muscles in primigravid subjects during pregnancy and the immediate postbirth period. *Phys Ther*. 1996;76(7):750–62.
9. Lobel M, DeLuca RS. Psychosocial sequelae of cesarean delivery: review and analysis of their causes and implications. *Soc Sci Med*. 2007;64:2272–84.

10. Gutke A, Lundberg M, Ostgaard HC, Oberg B. Impact of postpartum lumbopelvic pain on disability, pain intensity, health-related quality of life, activity level, kinesiophobia, and depressive symptoms. *Eur Spine J*. 2011;20(3):440–8.
11. Hunter SK. Performance fatigability: mechanisms and task specificity. *Cold Spring Harb Perspect Med*. 2017. [Epub ahead of print]. doi:10.1101/cshperspect.a029728.
12. Almuklass AM, Price RC, Gould JR, Enoka RM. Force steadiness as a predictor of time to complete a pegboard test of dexterity in young men and women. *J Appl Physiol (1985)*. 2016;120(12):1410–7.
13. Marmon AR, Pascoe MA, Schwartz RS, Enoka RM. Associations among strength, steadiness, and hand function across the adult life span. *Med Sci Sports Exerc*. 2011;43(4):560–7.
14. Enoka RM, Christou EA, Hunter SK, et al. Mechanisms that contribute to differences in motor performance between young and old adults. *J Electromyogr Kinesiol*. 2003;13(1):1–12.
15. Deering R, Senefeld JW, Pashibin T, Neumann DA, Hunter SK. Muscle function and fatigability of trunk flexors in males and females. *Biol Sex Differ*. 2017;8:12.
16. Borders N. After the afterbirth: a critical review of postpartum health relative to method of delivery. *J Midwifery Womens Health*. 2006;51(4):242–8.
17. Liddle SD, Pennick V. Interventions for preventing and treating low-back and pelvic pain during pregnancy (review). *Cochrane Database Syst Rev*. 2015;9:1–116.
18. Cheng CY, Fowles ER, Walker LO. Continuing education module: postpartum maternal health care in the United States: a critical review. *J Perinat Educ*. 2006;15(3):34–42.
19. Vahratian A, Johnson TR. Maternity leave benefits in the United States: today's economic climate underlines deficiencies. *Birth*. 2009;36(3):177–9.
20. Archuleta K. Handbook on Leave and Workplace Flexibilities for Childbirth, Adoption, and Foster Care. In: Handbook on Leave and Workplace Flexibilities for Childbirth, Adoption, and Foster Care. Washington, DC: United States Office of Personnel Management. 2015;9–18.
21. Cram JR. *Cram's Introduction to Surface Electromyography*. 2nd ed. Sudbury, Massachusetts: Jones and Bartlett; 2011. p. 412.
22. Kriska AM, Knowler WC, LaPorte RE, et al. Development of questionnaire to examine relationship of physical activity and diabetes in Pima Indians. *Diabetes Care*. 1990;13(4):401–11.
23. Ross RM, Murthy JN, Wollak ID, Jackson AS. The six minute walk test accurately estimates mean peak oxygen uptake. *BMC Pulm Med*. 2010;10:31.
24. Hunter SK. Muscle: the Primary Stabilizer and Mover of the Skeletal System. In: Neumann DA, editor. *Kinesiology of the Musculoskeletal System: Foundations for Rehabilitation*. 3rd ed. St. Louis, MO: Mosby Elsevier; 2016. pp. 47–76.
25. Riley ZA, Maerz AH, Litsey JC, Enoka RM. Motor unit recruitment in human biceps brachii during sustained voluntary contractions. *J Physiol*. 2008;586(8):2183–93.
26. Carpentier A, Duchateau J, Hainaut K. Motor unit behaviour and contractile changes during fatigue in the human first dorsal interosseus. *J Physiol*. 2001;534(Pt 3):903–12.
27. Farina D, Negro F. Common synaptic input to motor neurons, motor unit synchronization, and force control. *Exerc Sport Sci Rev*. 2015;43(1):23–33.
28. Hunter SK, Pereira HM, Keenan KG. The aging neuromuscular system and motor performance. *J Appl Physiol (1985)*. 2016;121:982–95.
29. Hunter SK. Sex differences in human fatigability: mechanisms and insight to physiological responses. *Acta Physiol (Oxf)*. 2014;210(4):768–89.
30. Brown SH, McGill SM. Transmission of muscularly generated force and stiffness between layers of the rat abdominal wall. *Spine*. 2009;34(2):E70–5.
31. Gracovetsky S, Farfan H, Lamy C. A mathematical model of the lumbar spine using an optimized system to control muscles and ligaments. *Orthop Clin North Am*. 1977;8(1):135–53.

32. Garber CE, Blissmer B, Deschenes MR, et al. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc.* 2011;43(7):1334–59.

Keywords:

ABDOMINAL MUSCLES; MUSCLE FATIGUE; STEADINESS; CESAREAN DELIVERY; VAGINAL DELIVERY; WOMEN'S HEALTH

© 2018 American College of Sports Medicine