

Marquette University

e-Publications@Marquette

---

Mathematics, Statistics and Computer Science Faculty Research and Publications    Mathematics, Statistics and Computer Science,  
Department of (- 2019)

---

8-2013

## The Pitman Inequality For Exchangeable Random Vectors

J. Behboodian  
*Shiraz University*

Naveen K. Bansal  
*Marquette University, naveen.bansal@marquette.edu*

Gholamhossein Hamedani  
*Marquette University, gholamhoss.hamedani@marquette.edu*

Hans Volkmer  
*University of Wisconsin - Milwaukee*

Follow this and additional works at: [https://epublications.marquette.edu/mscs\\_fac](https://epublications.marquette.edu/mscs_fac)



Part of the [Computer Sciences Commons](#), [Mathematics Commons](#), and the [Statistics and Probability Commons](#)

---

### Recommended Citation

Behboodian, J.; Bansal, Naveen K.; Hamedani, Gholamhossein; and Volkmer, Hans, "The Pitman Inequality For Exchangeable Random Vectors" (2013). *Mathematics, Statistics and Computer Science Faculty Research and Publications*. 164.

[https://epublications.marquette.edu/mscs\\_fac/164](https://epublications.marquette.edu/mscs_fac/164)

Marquette University

**e-Publications@Marquette**

***Mathematics and Statistical Sciences Faculty Research and Publications/College of Arts and Sciences***

***This paper is NOT THE PUBLISHED VERSION; but the author's final, peer-reviewed manuscript.*** The published version may be accessed by following the link in the citation below.

*Statistics & Probability Letters*, Vol. 83, No. 8 (August 2013): 1825+1829. [DOI](#). This article is © Elsevier and permission has been granted for this version to appear in [e-Publications@Marquette](#). Elsevier does not grant permission for this article to be further copied/distributed or hosted elsewhere without the express permission from Elsevier.

# The Pitman Inequality for Exchangeable Random Vectors

J. Behboodian

Shiraz University, Shiraz, Iran

Naveen Bansal

Department of Mathematics, Statistics & Computer Science, Marquette University, Milwaukee, WI

G.G. Hamedani

Department of Mathematics, Statistics & Computer Science, Marquette University, Milwaukee, WI

Hans Volkmer

Department of Mathematical Sciences, University of Wisconsin-Milwaukee, Milwaukee, WI Shiraz University, Shiraz, Iran

## Abstract

In this short article the following inequality called the “Pitman inequality” is proved for the exchangeable random vector  $(X_1, X_2, \dots, X_n)$  without the assumption of continuity and symmetry for each component  $X_i$ :

$$P\left(\left|\frac{1}{n}\sum_{i=1}^n X_i\right| \leq \left|\sum_{i=1}^n \alpha_i X_i\right|\right) \geq \frac{1}{2},$$

where all  $\alpha_i \geq 0$  are special weights with  $\sum_{i=1}^n \alpha_i = 1$ .

## Keywords

Exchangeability, Equality in distribution, Pitman inequality, Characteristic function

### 1. Introduction

Bose et al. (1993) established the following result: if  $X_1$  and  $X_2$  are *i.i.d.* (independent and identically distributed), continuous and symmetric about  $\theta$ , then  $(X_1 + X_2)/2$  is the Pitman-closest estimator of  $\theta$  within the class of all the estimators of the form  $\alpha X_1 + (1 - \alpha)X_2$  for  $0 \leq \alpha \leq 1$ . In other words we have

$$(1) P(|(X_1 + X_2)/2 - \theta| \leq |\alpha X_1 + (1 - \alpha)X_2 - \theta|) \geq \frac{1}{2}.$$

Assume, without loss of generality, that  $\theta = 0$  and consider the general form of (1) where  $X_1, X_2, \dots, X_n$  are *i.i.d.* continuous and symmetric, i.e.

$$(2) P\left(\left|\frac{1}{n}\sum_{i=1}^n X_i\right| \leq \left|\sum_{i=1}^n \alpha_i X_i\right|\right) \geq \frac{1}{2},$$

where all  $\alpha_i \geq 0$  and  $\sum_{i=1}^n \alpha_i = 1$ . We wish to show that (2) is false for  $n > 2$ . Let  $n = 3, \alpha_1 = \alpha_2 = 0, \alpha_3 = 1$  and let  $f(x)$  be a symmetric *pdf* (probability density function) of the *i.i.d.* random variables  $X_1, X_2, X_3$ . Let  $g = f * f$  be the convolution of  $f$  with itself, so  $g$  is the *pdf* of  $X_1 + X_2$  and is symmetric as well. It is easy to show that

$$1 - P\left(\left|\frac{1}{3}\sum_{i=1}^3 X_i\right| \leq |X_3|\right) = 2 \int_0^\infty \int_{-u/4}^{u/2} f(v)g(u)dvdu = 2 \int_0^\infty \left(F^*\left(\frac{u}{2}\right) + F^*\left(\frac{u}{4}\right)\right)g(u)du,$$

where  $F^*(u) = \int_0^u f(x)dx$ . It appears that (2) may be true for many symmetric *pdf*s; for example, it is true for the standard normal *pdf* with  $n = 3, \alpha_1 = \alpha_2 = 0$  and  $\alpha_3 = 1$ . However, consider

$$f(x) = \frac{1}{4(1 + |x|)^{3/2}}.$$

Then, after some computation, for  $x > 0$ ,

$$g(x) = -\frac{1}{2x^2} + \frac{\sqrt{1+x}(x^2 + 2x + 4)}{2(x+2)^2 x^2}.$$

It can be shown that

$$1 - P\left(\left|\frac{1}{3}\sum_{i=1}^3 X_i\right| \leq |X_3|\right) = \frac{5}{12} - \frac{1}{8}\ln 3 - \frac{1}{4}\sqrt{2}\arctan\left(\frac{1}{4}\sqrt{2}\right) - \frac{1}{16}\ln(3\sqrt{2} + 4) + \frac{5}{12}\sqrt{2} = 0.5281 \dots > \frac{1}{2}.$$

The goal here is to prove (2) for an exchangeable random vector  $(X_1, X_2, \dots, X_n)$  with special weights  $\alpha_i$  via a simple proof and without the assumption of continuity and symmetry. That is, we want to show that for exchangeable random variables  $X_1, X_2, \dots, X_n$ ,

$$(3) P\left(\left|\frac{1}{n}\sum_{i=1}^n X_i\right| \leq \left|\sum_{i=1}^n \alpha_i X_i\right|\right) \geq \frac{1}{2},$$

where all  $\alpha_i \geq 0$  are special weights with  $\sum_{i=1}^n \alpha_i = 1$ .

Exchangeability plays an important role in forecasting. Suppose one is interested in forecasting a random variable  $Y$  based on exchangeable or *i.i.d.* random variables  $Z_1, Z_2, \dots, Z_n$ . An important problem is to find a best predictor of  $Y$  based on a linear function of  $Z_1, Z_2, \dots, Z_n$ . It is well known that  $\bar{Z} = n^{-1} \sum_{i=1}^n Z_i$  is the best predictor in the least squares sense. In the sense of Pitman closeness, one may be interested in proving

$$P\left(|Y - \bar{Z}| \leq \left|Y - \sum_{i=1}^n \alpha_i Z_i\right|\right) \geq \frac{1}{2}.$$

This is equivalent to (3) with  $X_i = (Y - Z_i)$ . Note that in this case  $X_1, X_2, \dots, X_n$  are exchangeable, but not necessarily independent. Therefore, it is necessary to consider exchangeable  $X_i$ 's in (3).

We refer the interested reader to related works by Balakrishnan et al. (2009), Bose et al. (1993) and Ghosh and Sen (1989). We would like to mention here that no results have been reported in the literature regarding (1), (3) for exchangeable random vectors. Over the past six decades many researchers have been working on projects dealing with exchangeability, which is a weaker assumption than that of *i.i.d.* The interested reader is referred, among others, to the book by Chow and Teicher (1997).

In Section 2, we state a useful technique called “the equal in distribution technique (EDT)”, which will be employed in this short article. Section 3 is devoted to the proof of (3) for  $n = 2$ . The final section deals with the proof of (3) for  $n > 2$ .

## 2. The equal in distribution technique (EDT)

Two vectors  $U = (X_1, X_2, \dots, X_n)$  and  $V = (Y_1, Y_2, \dots, Y_n)$  are said to be equal in distribution and denoted by  $U \stackrel{d}{=} V$  if they have the same distributions or characteristic functions. It is clear that

if  $U \stackrel{d}{=} V$  then  $g(U) \stackrel{d}{=} g(V)$  for any measurable function  $g: \mathbb{R}^n \rightarrow \mathbb{R}^k$ . The EDT plays a significant role in the proofs of many results in probability theory. For example, using the exchangeability of  $(X_1, X_2, \dots, X_n)$ ,

i.e.  $(X_1, X_2, \dots, X_n) \stackrel{d}{=} (X_{i_1}, X_{i_2}, \dots, X_{i_n})$  for any one of the  $n!$  permutations of  $(1, 2, \dots, n)$ , we conclude that

(i) the  $X_i$ 's are identically distributed;

(ii)  $(X_1, X_2, \dots, X_m)$  is exchangeable for  $2 \leq m \leq n$ ;

(iii)  $\sum_{j=1}^m X_j \stackrel{d}{=} \sum_{j=1}^m X_{i_j}$  for  $1 \leq m \leq n$ .

In particular,  $(X_1, X_2, \dots, X_n)$  is exchangeable if and only if

$(X_1 - \theta, X_2 - \theta, \dots, X_n - \theta)$  is exchangeable.

### Remark 1

We would like to mention here that although (3) is true for any convex combination of  $X_i$ 's when  $n=2$  (see Section 3 below), it is not true when  $n > 2$ , even in the case of symmetry, as the following example shows. Therefore, in proving (3) for  $n > 2$  we have to restrict the  $\alpha_i$ 's to certain “special weights”.

### Example 1

Let  $\alpha_1 = 1$  and  $\alpha_i = 0, i = 2, 3, \dots, n$  in (3). Then

$$\sum_{i=1}^n P(|\sum_{i=1}^n X_i| \leq |X_j|) \geq 1,$$

$$\sum_{i=j}^n P \left( \left| \frac{1}{n} \sum_{i=1}^n X_i \right| \leq |X_j| \right) \geq 1,$$

since for every  $\omega$  there is a  $j$  such that  $|X_j(\omega)| \geq |X_i(\omega)|$  for all  $i = 1, 2, \dots, n$ . Since the  $X_i$ 's are exchangeable, we obtain

$$(4) P \left( \left| \frac{1}{n} \sum_{i=1}^n X_i \right| \leq |X_1| \right) \geq \frac{1}{n}.$$

To show that  $\frac{1}{n}$  in (4) is optimal, we let  $e_i, i = 1, 2, \dots, n$ , denote the standard unit vectors in  $\mathbb{R}^n$ . Let  $P$  be the probability measure with mass  $\frac{1}{n}$  at each  $e_i$ . Let  $X_i: \mathbb{R}^n \rightarrow \mathbb{R}$  be the  $i$ th projection map. Then, the  $X_i$ 's are exchangeable and

$$P \left( \left| \frac{1}{n} \sum_{i=1}^n X_i \right| \leq |X_1| \right) = P(e_1) = \frac{1}{n}.$$

Note that if we take  $2ne_i$ 's and  $-e_i$ 's, the random variables  $X_i$  will be symmetric as well.

### 3. Proof of (3) for $n = 2$ and $\alpha_1 + \alpha_2 = 1$

Applying the two-variable function

$$g(t_1, t_2) = (\alpha_1 t_1 + \alpha_2 t_2, \alpha_2 t_1 + \alpha_1 t_2)$$

on both sides of  $(X_1, X_2) \stackrel{d}{=} (X_2, X_1)$ , by the EDT we obtain

$$(\alpha_1 X_1 + \alpha_2 X_2, \alpha_2 X_1 + \alpha_1 X_2) \stackrel{d}{=} (\alpha_1 X_2 + \alpha_2 X_1, \alpha_2 X_2 + \alpha_1 X_1).$$

Now, setting  $U_1 = \alpha_1 X_1 + \alpha_2 X_2$  and  $U_2 = \alpha_2 X_1 + \alpha_1 X_2$  we have  $(U_1, U_2) \stackrel{d}{=} (U_2, U_1)$ . Applying the function  $h(t_1, t_2) = |t_1| - |t_2|$  on both sides of  $(U_1, U_2) \stackrel{d}{=} (U_2, U_1)$ , by the EDT we obtain  $|U_1| - |U_2| \stackrel{d}{=} |U_2| - |U_1|$ , i.e.  $|U_1| - |U_2|$  is symmetric. Using the fact that  $U_1 + U_2 = X_1 + X_2$ , we have

$$\begin{aligned} P(|(X_1 + X_2)/2| \leq |\alpha_1 X_1 + \alpha_2 X_2|) &= P(|U_1 + U_2| \leq 2|U_1|) \geq P(|U_1| + |U_2| \leq 2|U_1|) = P(|U_1| - |U_2| \geq 0) \\ &= \frac{1}{2}. \end{aligned}$$

Thus, (3) is proved for  $n = 2$  with  $\alpha_1 + \alpha_2 = 1$ .

### 4. Proof of (3) for $n > 2$

We will prove (3) for  $n = 3$  and  $n = 4$  for some specific weights  $\alpha_i$ . For  $n=3$ , we prove (3) when one of the weights is the average of the other two weights.

#### Theorem A

If  $(X_1, X_2, X_3)$  is exchangeable, then

$$(5) P \left( \left| \frac{1}{3} \sum_{i=1}^3 X_i \right| \leq \left| \frac{2}{3} \alpha X_1 + \frac{1}{3} X_2 + \frac{2}{3} \beta X_3 \right| \right) \geq \frac{1}{2},$$

where  $0 \leq \alpha \leq 1$  and  $\alpha + \beta = 1$ .

## Proof

Let

$$T_1 = \frac{2}{3}X_1 + \frac{1}{3}X_2, T_2 = \frac{2}{3}X_3 + \frac{1}{3}X_2,$$

Then

$$\frac{1}{3} \sum_{i=1}^3 X_i = \frac{1}{2} \sum_{j=1}^2 T_j.$$

Clearly

$$\alpha T_1 + \beta T_2 = \frac{2}{3}\alpha X_1 + \frac{1}{3}X_2 + \frac{2}{3}\beta X_3.$$

The pair  $(T_1, T_2)$  is exchangeable. To show this, we observe that by exchangeability

$$(X_1, X_2, X_3) \stackrel{d}{=} (X_3, X_2, X_1).$$

Define the function  $g: \mathbb{R}^3 \rightarrow \mathbb{R}^2$  as follows:

$$g(u_1, u_2, u_3) = \left( \frac{2}{3}u_1 + \frac{1}{3}u_2, \frac{2}{3}u_3 + \frac{1}{3}u_2 \right).$$

Then EDT implies that  $(T_1, T_2)$  is exchangeable.

Now, inserting  $T_1$  and  $T_2$  in the left hand side of (5), by the case  $n = 2$ , we have

$$P \left( \left| \frac{T_1 + T_2}{2} \right| \leq |\alpha T_1 + \beta T_2| \right) \geq \frac{1}{2}.$$

### Remark 2

Theorem A can easily be generalized for any odd integer  $2n + 1 > 3$ .

For  $n = 4$ , we prove (3) when any pair of weights is the same as the weights of the other pairs. Due to exchangeability, it suffices to consider  $\alpha_1 = \alpha_2 = \frac{\alpha}{2}, \alpha_3 = \alpha_4 = \frac{\beta}{2}$  and  $\alpha \neq \beta$  with  $\alpha + \beta = 1$ .

### Theorem B

If  $(X_1, X_2, X_3, X_4)$  is exchangeable, then

$$(6) P \left( \left| \frac{1}{4} \sum_{i=1}^4 X_i \right| \leq \left| \frac{\alpha}{2} X_1 + \frac{\alpha}{2} X_2 + \frac{\beta}{2} X_3 + \frac{\beta}{2} X_4 \right| \right) \geq \frac{1}{2},$$

where  $0 \leq \alpha \leq 1, \alpha \neq \beta$  and  $\alpha + \beta = 1$ .

## Proof

Let

$$T_1 = \frac{X_1 + X_2}{2}, T_2 = \frac{X_3 + X_4}{2},$$

Then

$$\frac{1}{4} \sum_{i=1}^4 X_i = \frac{1}{2} \sum_{j=1}^2 T_j.$$

Clearly

$$\alpha T_1 + \beta T_2 = \frac{\alpha}{2} X_1 + \frac{\alpha}{2} X_2 + \frac{\beta}{2} X_3 + \frac{\beta}{2} X_4.$$

The pair  $(T_1, T_2)$  is exchangeable. To show this, we observe that by exchangeability

$$(X_1, X_2, X_3, X_4) \stackrel{d}{=} (X_3, X_4, X_2, X_1).$$

Define the function  $g: \mathbb{R}^4 \rightarrow \mathbb{R}^2$  as follows:

$$g(u_1, u_2, u_3, u_4) = \left( \frac{u_1 + u_2}{2}, \frac{u_3 + u_4}{2} \right).$$

Then EDT implies that  $(T_1, T_2)$  is exchangeable.

Now, inserting  $T_1$  and  $T_2$  in the left hand side of (6), by the case  $n = 2$ , we have

$$P\left(\left|\frac{T_1 + T_2}{2}\right| \leq |\alpha T_1 + \beta T_2|\right) \geq \frac{1}{2}.$$

### Remark 3

Theorem B can easily be generalized for any even integer  $2n > 2$ .

### Remarks 4

(i) Let  $X_1, X_2, X_3$  be *i.i.d.* positive continuous random variables, then it is easy to see that for  $(\alpha = 1, \beta = 0)$  or  $(\alpha = \frac{1}{3}, \beta = \frac{2}{3})$ , (5) becomes equality and hence Theorem A is optimal with  $\frac{1}{2}$ . (ii) The same can be said for Theorem B with four *i.i.d.* random variables. (iii) Let  $X_1, X_2, X_3$  be *i.i.d.* with an exponential distribution function  $F$  with parameter  $\frac{1}{2}$  ( $F(x) = 1 - e^{-x/2}, x > 0$ ).

Observe that

$$\frac{1}{3} \sum_{i=1}^3 X_i \leq \frac{1}{6} (X_1 + 4X_2 + X_3)$$

is the same as  $X_1 + X_3 \leq 2X_2$  and

$$P(X_1 + X_3 \leq 2X_2) = P(E) = \iiint_E \frac{1}{8} e^{-\frac{1}{2}(x_1+x_2+x_3)} dx_1 dx_2 dx_3 = \frac{4}{9}.$$

This shows that Theorem A is not true if the conditions on the coefficients do not hold. (iv) If we take four *i.i.d.* random variables in (iii) we observe that

$$\frac{1}{4} \sum_{i=1}^4 X_i \leq \frac{1}{8} (X_1 + X_2 + 2X_3 + 4X_4)$$

is the same as  $X_1 + X_2 \leq 2X_4$  and  $P(X_1 + X_2 \leq 2X_4) = \frac{4}{9}$ . Therefore, Theorem B is not true if the conditions on the coefficients do not hold.

## References

- Balakrishnan et al., 2009. N. Balakrishnan, G. Iliopoulos, J.P. Keating, R.L. Mason. **Pitman closeness of sample median to population median**. *Statistics and Probability Letters*, 79 (2009), pp. 1759-1766
- Bose et al., 1993. S. Bose, G.S. Datta, M. Ghosh. **Pitman's measure of closeness for symmetric stable distributions**. *Statistics and Probability Letters*, 17 (1993), pp. 245-251
- Chow and Teicher, 1997. Y.S. Chow, H. Teicher. **Probability Theory, Independence and Interchangeability**. (third ed.), Springer (1997)
- Ghosh and Sen, 1989. M. Ghosh, P.K. Sen. **Median unbiasedness and Pitman-closeness**. *Journal of the American Statistical Association*, 84 (1989), pp. 1089-1091