

Marquette University

e-Publications@Marquette

Mathematics, Statistics and Computer Science Faculty Research and Publications Mathematics, Statistics and Computer Science,
Department of (- 2019)

2012

Characterizations of Certain Continuous Univariate Distributions Based on the Conditional Distribution of Generalized Order Statistics

M. Ahsanullah
Rider University

Gholamhossein Hamedani
Marquette University, gholamhoss.hamedani@marquette.edu

Follow this and additional works at: https://epublications.marquette.edu/mscs_fac



Part of the [Computer Sciences Commons](#), [Mathematics Commons](#), and the [Statistics and Probability Commons](#)

Recommended Citation

Ahsanullah, M. and Hamedani, Gholamhossein, "Characterizations of Certain Continuous Univariate Distributions Based on the Conditional Distribution of Generalized Order Statistics" (2012). *Mathematics, Statistics and Computer Science Faculty Research and Publications*. 184.
https://epublications.marquette.edu/mscs_fac/184

CHARACTERIZATIONS OF CERTAIN CONTINUOUS UNIVARIATE
DISTRIBUTIONS BASED ON THE CONDITIONAL DISTRIBUTION
OF GENERALIZED ORDER STATISTICS

M. Ahsanullah¹ and G.G. Hamedani²

¹ Department of Management Sciences, Rider University,
Lawrenceville, NJ 08648-3009. Email: ahsan@rider.edu

² Department of Mathematics, Statistics and Computer Science,
Marquette University Milwaukee, WI 53201-1881. Email:
g.hamedani@mu.edu

ABSTRACT

The problem of characterizing probability distributions is an interesting problem which has recently attracted the attention of many researchers. Various characterization results have been established in different directions as reported in the literature. We present here, various characterizations of certain univariate continuous distributions based on the conditional distribution of generalized order statistics.

1. INTRODUCTION

The concept of generalized order statistics (*gos*) from a univariate continuous distribution was introduced by Kamps [9] in terms of their joint *pdf* (probability density function). The order statistics, record values, *k*-record values, Pfeifer's records and progressive type II order statistics are special cases of the *gos*. The *rv*'s (random variables) $X(1, n, m, k)$, $X(2, n, m, k)$, ..., $X(n, n, m, k)$, $k > 0$, $m \in \mathbb{R}$, are *n gos* from an absolutely continuous *cdf* (cumulative distribution function) F with corresponding *pdf* f if their joint *pdf*, $f_{1,2,\dots,n}(x_1, x_2, \dots, x_n)$, can be written as

$$f_{1,2,\dots,n}(x_1, x_2, \dots, x_n) = k \left(\prod_{j=1}^{n-1} \gamma_j \right) \left[\prod_{j=1}^{n-1} (1 - F(x_j))^m f(x_j) \right] \\ \times (1 - F(x_n))^{k-1} f(x_n), \quad F^{-1}(0+) < x_1 < x_2 < \dots < x_n < F^{-1}(1-), \quad (1.1)$$

where $\gamma_j = k + (n - j)(m + 1)$ for all j , $1 \leq j \leq n$, k is a positive integer and $m \geq -1$.

If $k = 1$ and $m = 0$, then $X(r, n, m, k)$ reduces to the ordinary *r*th order statistic and (1.1) will be the joint *pdf* of order statistics $X_{1,n} \leq X_{2,n} \leq \dots \leq X_{n,n}$ from F . If $k = 1$ and $m = -1$, then (1.1) will be the joint *pdf* of the first *n* upper record values of the *i.i.d.* (independent and identically distributed) *rv*'s with *cdf* F and *pdf* f .

Integrating out $x_1, x_2, \dots, x_{r-1}, x_{r+1}, \dots, x_n$ from (1.1) we obtain the *pdf* $f_{r,n,m,k}$ of $X(r, n, m, k)$

$$f_{r,n,m,k}(x) = \frac{c_r}{\Gamma(r)} (1-F(x))^{\gamma_r-1} f(x) g_m^{r-1}(F(x)), \quad (1.2)$$

where $c_r = \prod_{j=1}^r \gamma_j$ and

$$g_m(x) = \frac{1}{m+1} \left[1 - (1-x)^{m+1} \right], \quad m \neq -1$$

$$= -\ln(1-x), \quad m = -1, \quad x \in (0,1).$$

Since $\lim_{m \rightarrow -1} \frac{1}{m+1} \left[1 - (1-x)^{m+1} \right] = -\ln(1-x)$, we will write

$$g_m(x) = \frac{1}{m+1} \left[1 - (1-x)^{m+1} \right], \quad \text{for all } x \in (0,1) \text{ and all } m \text{ with}$$

$$g_{-1}(x) = \lim_{m \rightarrow -1} g_m(x).$$

The joint *pdf* of $X(r, n, m, k)$ and $X(r+1, n, m, k)$, $1 \leq r < n$, is given by (see Kamps [9], p.68)

$$f_{r,r+1,n,m,k}(x, y)$$

$$= \frac{c_{r+1}}{\Gamma(r)} (1-F(x))^m f(x) g_m^{r-1}(F(x)) (1-F(y))^{\gamma_{r+1}-1} f(y), \quad x < y$$

and consequently the conditional *pdf* of $X(r+1, n, m, k)$ given $X(r, n, m, k) = x$, for $m \geq -1$, is

$$f_{r+1|r,n,m,k}(y|x) = \gamma_{r+1} \left(\frac{1-F(y)}{1-F(x)} \right)^{\gamma_{r+1}-1} \cdot \frac{f(y)}{(1-F(x))}, \quad y > x, \quad (1.3)$$

where $\gamma_{r+1} = \gamma_r - 1 - m$ (see [2], p. 383).

2. CHARACTERIZATION RESULTS

In this section we present characterizations of certain univariate continuous distributions which include, as special cases, exponential, Pareto, power function, Rayleigh and Weibull distributions. We refer the interested readers to the references [3]–[8], [10] and [11] for further characterizations of distributions via linearity of regression or conditional distributions of order statistics, record values and *gos*.

Proposition 2.1:

Let $X : \Omega \rightarrow (0, \infty)$ be a continuous random variable with *cdf* F and g be a continuous function on $(0, \infty)$ with $\lim_{x \rightarrow 0} g(x) = 1$ and $\lim_{x \rightarrow \infty} g(x) = 0$. If

$$P(X(r+1, n, m, k) \geq x \mid X(r, n, m, k) = t) = (g(x-t))^{Y_{r+1}}, \quad x \geq t, \quad (2.1)$$

then $F(x) = 1 - e^{-\lambda x}$, $x > 0$ for some $\lambda > 0$.

Proof:

In view of (1.3), from (2.1) we have

$$\left(\frac{1-F(x)}{1-F(t)} \right)^{Y_{r+1}} = (g(x-t))^{Y_{r+1}}, \quad x \geq t,$$

or

$$\frac{1-F(x)}{1-F(t)} = g(x-t), \quad x \geq t. \quad (2.2)$$

Now, letting $t \rightarrow 0$, we arrive at $1-F(x) = g(x)$ for all $x > 0$ and again in view of (2.2) we will have

$$1-F(x) = (1-F(x-t))(1-F(t)), \quad x \geq t. \quad (2.3)$$

Equation (2.3) is the well-known Cauchy functional equation. The non-zero solution of this equation is (see, [1], Theorem 1 page 38) $1-F(x) = e^{c x}$, where c is a constant. Using the boundary conditions $\lim_{x \rightarrow 0} F(x) = 0$ and $\lim_{x \rightarrow \infty} F(x) = 1$ we obtain $F(x) = 1 - e^{-\lambda x}$, $x \geq 0$ for some $\lambda > 0$.

Proposition 2.2:

Let $X : \Omega \rightarrow [1, \infty)$ be a continuous random variable with *cdf* F , $F(1) \neq 1$ and g be a continuous function on $[1, \infty)$ with $g(1) = 1$ and $\lim_{x \rightarrow \infty} g(x) = 0$. If

$$P(X(r+1, n, m, k) \geq x \mid X(r, n, m, k) = t) = \left(g\left(\frac{x}{t}\right) \right)^{Y_{r+1}}, \quad x \geq t, \quad (2.4)$$

then $F(x) = 1 - x^{-\lambda}$, $x \geq 1$ for some $\lambda < 0$.

Proof:

As in the proof of Proposition 2.1, from (1.3) and (2.4) we have

$$\left(\frac{1-F(x)}{1-F(t)} \right)^{Y_{r+1}} = \left(g\left(\frac{x}{t}\right) \right)^{Y_{r+1}}, \quad x \geq t,$$

or

$$\frac{1-F(x)}{1-F(t)} = g\left(\frac{x}{t}\right), \quad x \geq t. \quad (2.5)$$

Now, letting $t=1$, we arrive at $1-F(x) = g(x)$ for all $x \geq 1$ and hence

$$1-F(x) = \left(1-F\left(\frac{x}{t}\right)\right)(1-F(t)), \quad x \geq t. \quad (2.6)$$

Equation (2.6) is a Cauchy functional equation. The non-zero solution of this equation is (see, [1], page 39) $1-F(x) = cx^\lambda$, where c is a constant. Using the boundary conditions $\lim_{x \rightarrow 1} F(x) = 0$ and $\lim_{x \rightarrow \infty} F(x) = 1$ we obtain $F(x) = 1 - x^\lambda$, $x \geq 1$ for some $\lambda < 0$.

Proposition 2.3:

Let $X : \Omega \rightarrow [0,1]$ be a continuous random variable with *cdf* F and g be a continuous function on $[0,1]$ with $g(0) = 1$ and $g(1) = 0$. If

$$P(X(r+1, n, m, k) \geq x \mid X(r, n, m, k) = t) = \left(g\left(\frac{1-x}{1-t}\right)\right)^{r+1}, \quad x \geq t, \quad (2.7)$$

then $F(x) = 1 - (1-x)^\lambda$, $x \geq 0$ for some $\lambda > 0$.

Proof:

As before, from (1.3) and (2.7) we have

$$\frac{1-F(x)}{1-F(t)} = g\left(\frac{1-x}{1-t}\right), \quad x \geq t. \quad (2.8)$$

Now, letting $t=0$, we arrive at $1-F(x) = g(1-x)$ for all $x \geq 0$ and hence

$$1-F(x) = \left(1-F\left(1-\frac{1-x}{1-t}\right)\right)(1-F(t)), \quad x \geq t,$$

or

$$1-F(x) = \left(1-F\left(\frac{x-t}{1-t}\right)\right)(1-F(t)), \quad x \geq t. \quad (2.9)$$

With $g(1-y) = 1-F(y)$, equation (2.9) will be equivalent to

$$g(1-y) = g\left(\frac{1-y}{1-t}\right)g(1-t).$$

Letting $z = 1 - y$, we have $g(z) = g\left(\frac{z}{1-t}\right)g(1-t)$ which is a Cauchy functional equation. The non-zero solution of this equation is (see, [1], page 39) $g(x) = cx^\lambda$, where c is a constant, and hence $F(x) = 1 - c(1-x)^\lambda$. Using the boundary conditions $\lim_{x \rightarrow 0} F(x) = 0$ and $\lim_{x \rightarrow 1} F(x) = 1$, we obtain $F(x) = 1 - (1-x)^\lambda$, $0 \leq x \leq 1$, for some $\lambda > 0$.

Proposition 2.4:

Let $X : \Omega \rightarrow (0, \infty)$ be a continuous random variable with cdf F and g be a continuous function on $(0, \infty)$ with $\lim_{x \rightarrow 0} g(x) = 1$ and $\lim_{x \rightarrow \infty} g(x) = 0$. If

$$P(X(r+1, n, m, k) \geq x \mid X(r, n, m, k) = t) = \left(g(x^\beta - t^\beta)\right)^{r+1}, \quad x \geq t, \quad (2.10)$$

where $\beta > 0$, then $F(x) = 1 - e^{-\lambda x^\beta}$, $x > 0$ for some $\lambda > 0$.

Proof:

From (1.3) and (2.10) we have

$$\frac{1 - F(x)}{1 - F(t)} = g(x^\beta - t^\beta), \quad x \geq t. \quad (2.11)$$

Now, letting $t \rightarrow 0$, we arrive at $1 - F(x) = g(x^\beta)$ for all $x > 0$ and in view of (2.11) we will have

$$\frac{1 - F(x)}{1 - F(t)} = 1 - F\left(\left(x^\beta - t^\beta\right)^{1/\beta}\right), \quad x \geq t. \quad (2.12)$$

Now, let $\varphi(x) = \ln(1 - F(x))$, then from (2.12), we have

$$\exp\{\varphi(x) - \varphi(t)\} = \exp\left\{\varphi\left(\left(x^\beta - t^\beta\right)^{1/\beta}\right)\right\}, \quad x \geq t. \quad (2.13)$$

Differentiating both sides of (2.13) with respect to x , we obtain

$$\varphi'(x) \exp\{\varphi(x) - \varphi(t)\} = \varphi'\left(\left(x^\beta - t^\beta\right)^{1/\beta}\right) \frac{x^{\beta-1}}{\left(x^\beta - t^\beta\right)^{1-1/\beta}} \exp\left\{\varphi\left(\left(x^\beta - t^\beta\right)^{1/\beta}\right)\right\},$$

and in view of (2.13)

$$\frac{\varphi'(x)}{\varphi'\left(\left(x^\beta - t^\beta\right)^{1/\beta}\right)} = \frac{x^{\beta-1}}{\left(x^\beta - t^\beta\right)^{1-1/\beta}}, \quad x > t. \quad (2.14)$$

From (2.14), we have

$$\frac{\varphi'(x)}{x^{\beta-1}} = \frac{\varphi'\left(\left(x^\beta - t^\beta\right)^{1/\beta}\right)}{\left(x^\beta - t^\beta\right)^{1-1/\beta}}.$$

Letting $h(x) = \frac{\varphi'(x)}{x^{\beta-1}}$, then in view of the last equation above, $h(x) = h\left(\left(x^\beta - t^\beta\right)^{1/\beta}\right)$

for all $x > t \geq 0$ and all $\beta > 0$. Setting $y = x^\beta - t^\beta$, we have $h\left(\left(y + t^\beta\right)^{1/\beta}\right) = h\left(y^{1/\beta}\right)$ for all $y \geq 0$, $t \geq 0$ and all $\beta > 0$. Thus $h(y) = c$, where c is a constant and hence $\varphi'(x) = cx^{\beta-1}$. In view of the boundary conditions on F , we obtain $F(x) = 1 - e^{-\lambda x^\beta}$, $x > 0$ for some $\lambda > 0$.

ACKNOWLEDGEMENTS

The authors are grateful to the three anonymous referees for their comments, in particular suggested references, which has certainly improved the presentation of the results.

REFERENCES

1. Aczel, J. (1996). *Lectures on Functional Equations and their Applications*. Academic Press, New York.
2. Ahsanullah, M. and Nevzorov, V.B. (2001). *Ordered Random Variables*. Nova Publishers.
3. Arslan, G., Ahsanullah, M. and Bairamov, I. (2005). On characteristic properties of the uniform distribution. *Sankhya: The Indian Journal of Statistics*, 67, 715-721.
4. Bairamov, I. and Özkal, T. (2007). On characterization of distributions through the properties of conditional expectations of order statistics. *Commun. in Statist.: Theo. and Meths.*, 36, 1319-1326.
5. Bieniek, M. and Szynal, D. (2003). Characterizations via linear regression of generalized order statistics. *Metrika*, 58, 259-271.
6. Blazquez, F.L. and Rebollo, J.L.M. (1997). A characterization of distributions based on linear regression of order statistics and record values. *Sankhya*, Series A, 59, 311-323.
7. Cramer, E., Kamps, U. and Keseling, C. (2004). Characterizations of distributions via linearity of regression of order random variables: A unified approach, *Commun. in Statist.: Theo. and Meths.*, 33, 2885-2911.
8. Dembinska, A. and Wesolowski, J. (1998). Linearity of regression for non-adjacent order statistics. *Metrika*, 48, 215-222.
9. Kamps, U. *A concept of generalized order statistics*. Teubner Skripten zur Mathematischen Stochastik. [Teubner Text on Mathematical Stochastics]. B.G. Teubner, Stuttgart.
10. Keseling, C. (1999). Conditional distributions of generalized order statistics and some characterizations. *Metrika*, 49, 27-40.
11. Yildiz, T. and Bairamov, I. (2008). Characterization of distributions by using the conditional expectations of generalized order statistics. *Selcuk J. Appl. Math.*, 9, 19-27.