

Marquette University

e-Publications@Marquette

Exercise Science Faculty Research and
Publications

Exercise Science, Department of

12-2020

Sex Differences in Fatigability Following Exercise Normalised to the Power–Duration Relationship

Paul Ansdell

Jacob Skarabot

Elliott Atkinson

Sarah Corden

Amber Tygart

See next page for additional authors

Follow this and additional works at: https://epublications.marquette.edu/exsci_fac

Authors

Paul Ansdell, Jacob Skarabot, Elliott Atkinson, Sarah Corden, Amber Tygart, Kristy M. Hicks, Kevin Thomas, Sandra K. Hunter, Glyn Howatson, and Stuart Goodall

Marquette University

e-Publications@Marquette

Department of Exercise Science Faculty Research and Publications/College of Health Sciences

This paper is NOT THE PUBLISHED VERSION.

Access the published version via the link in the citation below.

Journal of Physiology, Vol. 598, No. 24 (December 2020): 5717-5737. [DOI](#). This article is © Wiley and permission has been granted for this version to appear in [e-Publications@Marquette](#). Wiley does not grant permission for this article to be further copied/distributed or hosted elsewhere without the express permission from Wiley.

Sex Differences in Fatigability Following Exercise Normalised to the Power–Duration Relationship

Paul Ansdell

Faculty of Health and Life Sciences, Northumbria University, Newcastle upon Tyne, UK

Jakob Skarabot

Faculty of Health and Life Sciences, Northumbria University, Newcastle upon Tyne, UK

School of Sport, Exercise and Health Sciences, Loughborough University, Loughborough, UK

Elliott Atkinson

Faculty of Health and Life Sciences, Northumbria University, Newcastle upon Tyne, UK

Sarah Corden

Faculty of Health and Life Sciences, Northumbria University, Newcastle upon Tyne, UK

Amber Tygart

Faculty of Health and Life Sciences, Northumbria University, Newcastle upon Tyne, UK

Kirsty M. Hicks

Faculty of Health and Life Sciences, Northumbria University, Newcastle upon Tyne, UK

Kevin Thomas

Faculty of Health and Life Sciences, Northumbria University, Newcastle upon Tyne, UK

Sandra K. Hunter

Department of Physical Therapy, Marquette University, Milwaukee, WI, USA

Glyn Howatson

Faculty of Health and Life Sciences, Northumbria University, Newcastle upon Tyne, UK

Water Research Group, School of Environmental Sciences and Development, North-West University, Potchefstroom, South Africa

Stuart Goodall

Faculty of Health and Life Sciences, Northumbria University, Newcastle upon Tyne, UK

Corresponding author Dr S. Goodall: Faculty of Health and Life Sciences, Department of Sport, Exercise and Rehabilitation, Northumbria University, Newcastle upon Tyne, NE1 8ST, UK.

Abstract

Key points

- Knee-extensors demonstrate greater fatigue resistance in females compared to males during single-limb and whole-body exercise. For single-limb exercise, the intensity–duration relationship is different between sexes, with females sustaining a greater relative intensity of exercise.
- This study established the power–duration relationship during cycling, then assessed fatigability during critical power-matched exercise within the heavy and severe intensity domains.
- When critical power and the curvature constant were expressed relative to maximal ramp test power, no sex difference was observed. No sex difference in time to task failure was observed in either trial.
- During heavy and severe intensity cycling, females experienced lesser muscle de-oxygenation. Following both trials, females experienced lesser reductions in knee-extensor contractile function, and following heavy intensity exercise, females experienced less reduction in voluntary activation.
- These data demonstrate that whilst the relative power–duration relationship is not different between males and females, the mechanisms of fatigability during critical power-matched exercise are mediated by sex.

Abstract

Due to morphological differences, females demonstrate greater fatigue resistance of locomotor muscle during single-limb and whole-body exercise modalities. Whilst females sustain a greater relative intensity of single-limb, isometric exercise than males, limited investigation has been performed during whole-body exercise.

Accordingly, this study established the power–duration relationship during cycling in 18 trained participants (eight females). Subsequently, constant-load exercise was performed at critical power (CP)-matched intensities within the heavy and severe domains, with the mechanisms of fatigability assessed via non-invasive neurostimulation, near-infrared spectroscopy and pulmonary gas exchange during and following exercise.

Relative CP (72 ± 5 vs. $74 \pm 2\% P_{\max}$, $P = 0.210$) and curvature constant (51 ± 11 vs. $52 \pm 10 \text{ J P}_{\max}^{-1}$, $P = 0.733$) of the power–duration relationship were similar between males and females. Subsequent heavy ($P = 0.758$) and severe intensity ($P = 0.645$) exercise time to task failures were not different between sexes. However, females experienced lesser reductions in contractile function at task failure ($P \leq 0.020$), and greater vastus lateralis oxygenation ($P \leq 0.039$) during both trials. Reductions in voluntary activation occurred following both trials

($P < 0.001$), but were less in females following the heavy trial ($P = 0.036$). Furthermore, during the heavy intensity trial only, corticospinal excitability was reduced at the cortical ($P = 0.020$) and spinal ($P = 0.036$) levels, but these reductions were not sex-dependent. Other than a lower respiratory exchange ratio in the heavy trial for females ($P = 0.039$), no gas exchange variables differed between sexes ($P \geq 0.052$). Collectively, these data demonstrate that whilst the relative power–duration relationship is not different between males and females, the mechanisms of fatigability during CP-matched exercise above and below CP are mediated by sex.

Introduction

The exercise intensity–duration relationship is a phenomenon that permits mechanistic insight into the metabolic demands and physiological consequences of exercise within distinct intensity domains (Jones *et al.* **2010**; Poole *et al.* **2016**; Burnley & Jones, **2018**). The relationship between exercise intensity and maximal sustainable duration is hyperbolic at severe intensities, with the asymptote of the curve, the so-called critical power (CP), representing the threshold between exercise that is sustainable via a steady state of substrate utilisation and re-synthesis, and exercise that requires ATP re-synthesis from substrate-level phosphorylation (Poole *et al.* **2016**). Exercising above CP therefore leads to a progressive loss of intramuscular homeostasis, and impairment of the contractile apparatus (Vanhatalo *et al.* **2010**; Schäfer *et al.* **2019**). Below the CP, substrate-level phosphorylation and the associated accumulation of metabolites are maintained at a steady rate, permitting a 4–5 times slower rate of fatigability (Burnley *et al.* **2012**; Thomas *et al.* **2016**).

Recent evidence has shown that the power–duration relationship differs between males and females for intermittent, isometric knee-extensor exercise (Ansdell *et al.* **2019a**). The mechanism for this is probably a result of morphological differences within the exercising musculature in comparison to males. For instance, within the vastus lateralis (VL) the proportional area of type I muscle fibres is greater in females (Simoneau & Bouchard, **1989**; Staron *et al.* **2000**; Roepstorff *et al.* **2006**), and differences in sarcoplasmic reticulum calcium activity in response to fatiguing exercise, are evident between sexes (Harmer *et al.* **2014**). Furthermore, the female VL has a greater capillary density per unit of skeletal muscle (Roepstorff *et al.* **2006**), and an augmented vasodilatory response of the femoral artery to exercise (Parker *et al.* **2007**); collectively, these physiological differences could augment the delivery of oxygen to the working muscle. Indeed, VL type I fibre proportion (Vanhatalo *et al.* **2016**) and capillarisation (Mitchell *et al.* **2018a**) are positively correlated with aerobic exercise performance indices such as CP during cycling in males and mixed-sex samples, providing a potential explanation for why females are able to sustain a greater relative exercise intensity than males during an isometric exercise paradigm (Ansdell *et al.* **2019a**).

Whilst the data from Ansdell *et al.* (**2019a**) provide mechanistic insight into the sex difference in fatigability during single-limb exercise (e.g. Hunter *et al.* **2006**; Yoon *et al.* **2007b**; Russ *et al.* **2008**; Ansdell *et al.* **2017**), it does not fully explain why a similar sex difference is demonstrated during whole-body exercise (Glance *et al.* **2013**; Temesi *et al.* **2015**). To date, it remains unclear whether the power–duration relationship is different between sexes, and whether a sex difference in fatigability exists if exercise to exhaustion is performed relative to CP for whole-body exercise. Sundberg *et al.* (**2016**) provided an initial investigation into this topic, assessing the power–duration relationship during bouts ranging from 8 to 283 s, and average power during a 3 min all-out test. The authors suggested no sex difference in the time constant of ‘performance loss’, or maximum sustainable power. However, exercise bouts spanned both the severe and the extreme intensity domains, with the latter being defined by the attainment of task failure prior to the attainment of maximal oxygen uptake ($\dot{V}_{O_{2max}}$, Burnley & Jones, **2018**). Therefore, limited conclusions can be drawn regarding fatigability within and between the heavy and severe exercise intensity domains.

It is well established that the mechanisms of fatigability differ for whole-body and single-limb exercise (Hureau *et al.* **2018**; Thomas *et al.* **2018**). Indeed, Poole *et al.* (**2016**) suggested that parameters of the intensity–

duration relationship, such as the curvature constant (W'), are probably influenced by different factors in the two modalities of exercise. For example, termination of whole-body exercise above CP coincides with the attainment maximal cardiopulmonary responses (e.g. Vanhatalo *et al.* **2010**; Murgatroyd *et al.* **2011**), whereas for single-limb exercise, equivalent variables do not reach maximal values (e.g. Goodall *et al.* **2010**; Ansdell *et al.* **2019a**). Collectively, this evidence suggested that in healthy humans, whole-body severe intensity exercise performance is limited by a combination of convective and diffusive factors, whereas equivalent intensities of single-limb exercise are solely limited by diffusive factors. As described by Hureau *et al.* (**2018**) and Thomas *et al.* (**2018**), during whole-body exercise afferent feedback from other physiological systems (e.g. respiratory) contribute to the attainment of a 'sensory tolerance limit', in addition to accumulation of intramuscular metabolites and depletion of energy stores (Broxterman *et al.* **2015b**). Therefore, conclusions based on data from single-limb exercise (Ansdell *et al.* **2019a**) are limited in explaining exercise tolerance in a whole-body model.

The fact that females have more fatigue-resistant locomotor (Hunter *et al.* **2006**; Yoon *et al.* **2007b**; Russ *et al.* **2008**; Ansdell *et al.* **2017**) and respiratory musculature (Guenette *et al.* **2010**; Welch *et al.* **2018**) might lead to the hypothesis that females are able to sustain a greater relative work rate during cycling compared to males. However, morphological sex differences within the respiratory system have the potential to reduce high-intensity exercise tolerance in females. For example, when height-matched, females have smaller lung volumes and airway size, weaker respiratory muscles, and smaller alveolar surface area for gas exchange compared to males (Mead, **1980**; Crapo *et al.* **1982**; Martin *et al.* **1987**; Sheel *et al.* **2009**). Combined, these factors elicit a greater expiratory flow limitation in females at near-maximal ventilatory capacity during cycling (Guenette *et al.* **2007**). Furthermore, females demonstrate a greater work of breathing (W_b) than males (Witt *et al.* **2007**), and at peak exercise, the oxygen cost of breathing (expressed as a fraction of whole-body \dot{V}_{O_2}), is greater compared to males; 14 vs. 9%, respectively (Dominelli *et al.* **2015**). Additionally, females typically demonstrate a lower haemoglobin mass compared to males (Murphy, **2014**), which is considered to limit endurance exercise performance (Joyner, **2017**). Collectively, these potentially deleterious factors could counteract the greater fatigue resistance of locomotor and respiratory muscles during whole-body exercise, and negate the sex difference in critical torque demonstrated in single-limb exercise, where central factors are not a limitation to exercise (Ansdell *et al.* **2019a**). Additionally, these physiological sex differences could lead to different contributing mechanisms to exercise intolerance, or the so-called sensory tolerance limit (Hureau *et al.* **2018**; Thomas *et al.* **2018**).

Accordingly, the present study had two primary aims: (1) to compare the power–duration relationship between males and females during cycling; and (2) to determine whether a sex difference in fatigability (time to task failure, TTF), and the mechanisms (neuromuscular fatigue, muscle oxygenation and pulmonary gas exchange), existed when exercise intensity was normalised to the power–duration relationship. To do so, non-invasive neurostimulation was used to quantify the neural and contractile adjustments to cycling exercise, and near-infrared spectroscopy (NIRS) was recorded during exercise to monitor changes in knee-extensor oxygenation. It was hypothesised that: (1) due to the poorer convective aspects of oxygen transport negating the superior diffusive aspects that females demonstrate, no sex difference in the power–duration relationship would exist when expressed relative to maximum exercise performance, and (2) when exercise was CP-matched, TTF would not differ, but females would exhibit greater fatigue resistance, and lesser deoxygenation of the knee-extensors compared to males in both heavy and severe intensity domains.

Methods

Ethical approval

The study received institutional ethical approval from the Northumbria University Health and Life Sciences Research Ethics Committee (submission reference: 12,241) and was conducted according to all aspect of the *Declaration of Helsinki*, apart from registration in a database. Participants provided written informed consent to volunteer for the study.

Participants

Using the effect size from Ansdell *et al.* (2017) for the sex difference in fatigability during isometric exercise, a power calculation ($\alpha = 0.05$, power = 0.80) determined that a sample size of 16 participants was required. Thus, 10 males (mean \pm SD age: 25 ± 5 years, stature: 178 ± 9 cm, mass: 67.0 ± 8.8 kg) and eight females (age: 25 ± 6 years, stature: 169 ± 9 cm, mass: 63.3 ± 7.2 kg) gave written informed consent to participate. The females who volunteered were all using monophasic hormonal contraceptives (>6 months), and those using combined contraceptive pills were tested in the 21 day consumption period of the pill cycle in order to negate the effects of endogenous hormones on neuromuscular function and fatigability (Ansdell *et al.* 2019c). To ensure homogeneity in the training status of participants, minimum criteria were set for relative $\dot{V}_{O_{2max}}$ and maximal ramp test power (P_{max}) attained in the first visit (see below). These values were based upon recommendations by De Pauw *et al.* (2013) for males, and Decroix *et al.* (2016) for females and were as follows: minimum $\dot{V}_{O_{2max}}$ of 55 and 48 ml kg⁻¹ min⁻¹, and P_{max} of 4.6 and 3.8 W kg⁻¹ for males and females, respectively, and a minimum weekly training duration of ≥ 5 h week⁻¹. Participants had to achieve one of the aforementioned criteria in order to proceed to the subsequent experimental visits. In total, 18 males and 16 females were screened to achieve the resultant sample size.

Experimental design

All participants visited the laboratory six or seven times, completing a familiarisation visit, three or four constant intensity trials to estimate CP, then subsequent trials 10% above and below CP. Testing took place over a 3–5 week period, with a minimum of 48 h between visits to allow recovery (Carroll *et al.* 2016). The time of day for each testing session was controlled (± 1 h) to account for diurnal variations in maximal force-generating capacity and corticospinal excitability (Tamm *et al.* 2009). All visits were conducted in an environmentally controlled laboratory facility (TIS Services, Environmental Control Specialists, Alton, UK), where the conditions were pre-set to 20°C and 40% relative humidity.

Experimental protocol

Familiarisation and incremental exercise test

Upon providing written informed consent, participants performed a 5 min warm up (80–100 W) on a cycle ergometer (Velotron Pro, RacerMate Inc., Seattle, WA, USA) at a self-selected cadence (60–100 rpm). Participants were then given 2 min of rest, during which they remained stationary on the cycle ergometer, before an incremental exercise test began. For both sexes, the test started at 100 W, then for males the intensity increased gradually by 25 W min⁻¹ (0.416 W s⁻¹), and for females by 20 W min⁻¹ (0.333 W s⁻¹). The different rate of intensity increase was intended to produce ramp tests of similar duration in both sexes, due to lower absolute power outputs demonstrated in females (Sundberg *et al.* 2016), in an attempt to negate the effects of test duration on cardiopulmonary outcomes (Yoon *et al.* 2007a). Mean ramp test duration was not different between males and females (10.5 ± 1.2 vs. 9.1 ± 2.1 min, $P = 0.103$, respectively). The test was terminated once the participant's self-selected cadence decreased by 10 rpm, despite strong verbal encouragement. During the test, expired gas was analysed breath-by-breath using an online system (Vyntus CPX, Jaeger, CareFusion, Germany). The outcome variables from the ramp test were $\dot{V}_{O_{2max}}$ (ml kg⁻¹ min⁻¹) and

P_{\max} (W). Following the incremental exercise test, participants rested for 15 min, before a neuromuscular familiarisation was performed, including all forms of non-invasive neurostimulation and a full neuromuscular function assessment and maximal respiratory pressure assessments (see below).

Critical power estimation trials

To estimate CP, participants completed a minimum of three constant-load exercise trials to task failure. The intensities for the initial three trials were set at 110, 90 and 80% of P_{\max} and were performed on separate days (minimum 24 h between trials) in a randomised order, designed to elicit task failure within 2–15 min (Poole *et al.* 1988). TTF (s) was recorded as the first time at which participants' cadence fell by 10 rpm. Although strong verbal encouragement was provided throughout the test, no feedback was provided to participants about the power output and time elapsed during the trials. Gas exchange was recorded continuously throughout each trial, and a criterion of an end-exercise \dot{V}_{O_2} of $>95\% \dot{V}_{O_2\max}$ was set; all trials used for estimation achieved this. The parameters of the power–duration relationship (CP and W') were estimated using the inverse linear model (eqn 1), the linear work–time model (eqn 2) and the hyperbolic model (eqn 3). The equation with the highest r^2 and lowest standard error (SE) was selected for each individual and used for all further analysis (Mitchell *et al.* 2018a). The hyperbolic fit was used for 10 participants, the linear fit for 5, and the 1/time fit for 3:

$$P = W' \cdot \left(\frac{1}{t}\right) + CP$$

(1)

$$W = CP \cdot t + W'$$

(2)

$$t + W'/(P - CP)$$

(3)

where t is time to task failure, P is power output and W is total work done. If three estimation trials resulted in a large SE for CP ($>5\%$ of the mean) and W' ($>10\%$), a fourth trial was performed (Mitchell *et al.* 2018a). This occurred for three out of the 18 participants (two males, one female).

Severe and heavy intensity trials

Once CP and W' were estimated, severe (110% CP) and heavy (90% CP) intensity trials were performed on separate days in a randomised order. Each session began with electrical nerve stimulation and transcranial magnetic stimulation (TMS) thresholds being determined. Participants then completed a standardised isometric warm up (Gruet *et al.* 2014), before a baseline assessment of neuromuscular function. Following this, NIRS optodes were affixed over the VL and baseline measures were recorded for 5 min on the cycle ergometer with the right leg relaxed in the fully extended position (crank angle 180° from top dead centre). Resting measures of gas exchange were also recorded in this period, then both NIRS and gas exchange were continuously sampled until task failure. Participants completed a 5 min warm up (80–100 W), followed by 1 min of seated rest on the ergometer. In the 5–10 s before the trial, participants were instructed to obtain their self-selected cadence against no resistance, then when achieved, the resistance was applied in a square wave fashion. TTF was recorded for the severe intensity trial, whereas for the heavy intensity trial participants cycled to task failure, or for 60 min, whichever occurred sooner. Immediately upon task failure (<20 s) participants transitioned from the cycle ergometer to the dynamometer and commenced a neuromuscular assessment which was completed within 2.5 min after exercise (described below).

Experimental procedures

Pulmonary gas exchange

Breath-by-breath pulmonary gas exchange and ventilation were measured continuously during all trials. With minute ventilation (\dot{V}_E), oxygen consumption (\dot{V}_{O_2}), carbon dioxide production (\dot{V}_{CO_2}) and respiratory exchange ratio (RER) were quantified. Prior to each visit, the Vyntus CPX was calibrated for oxygen (O_2) and carbon dioxide (CO_2) with gas of known concentration (16% O_2 and 4.97% CO_2) using an electrochemical fuel cell and non-dispersive infrared cell, respectively. Ventilatory volumes were calibrated using a digital turbine transducer at high (2 l s^{-1}) and low (0.2 l s^{-1}) flow rates.

Neuromuscular function assessments

Measures of neuromuscular function were assessed before and after exercise, starting within 30 s of task failure. Pre-exercise neuromuscular assessments began with two practice maximal voluntary contractions (MVCs) to ensure potentiation of subsequent evoked measures, followed by three ~ 3 s MVCs, all separated by 30 s. During these three MVCs, motor nerve stimulation (MNS) was delivered when peak force plateaued, and then ~ 2 s after the MVC to measure voluntary activation (VA_{MNS}) and quadriceps potentiated twitch amplitude ($Q_{tw.pot}$) of the knee-extensors. Single-pulse TMS was subsequently delivered during two sets of five 3–5 s contractions at 100, 87.5, 75, 62.5 and 50% MVC, with 5 s rest between contractions and 10 s rest between sets, to determine VA_{TMS} (Dekerle *et al.* **2019b**). Finally, 10 single- and 10 paired-pulse TMS stimulations were delivered during a 10% MVC contraction in an alternate order to determine corticospinal excitability and short-interval cortical inhibition (SICI), respectively. Measures of neuromuscular function (MVC, $Q_{tw.pot}$, VA_{MNS}) were measured within 30 s of task failure, and VA_{TMS} was measured within 2–2.5 min, in an attempt to minimise the dissipation of fatigue (Gruet *et al.* **2014**).

Transcranial magnetic stimulation

Single- and paired-pulse stimuli (1 ms duration) were delivered to the contralateral (left) motor cortex via a concave double cone coil oriented to induce a posterior-to-anterior cortical current (110 mm diameter, maximum output 1.4 T) powered by two linked monopulse stimulators (Magstim Bistim and Magstim²⁰⁰, The Magstim Company, Whitland, UK). Optimal coil placement was determined as the position that elicited the greatest rectus femoris (RF) motor evoked potential (MEP) with concomitant smallest antagonist (biceps femoris, BF) MEP during a 10% MVC at 50–70% stimulator output. This position was marked on the scalp with an indelible marker to ensure consistent placement during trials. Stimulator intensity for VA_{TMS} was determined as the intensity that elicited the greatest superimposed twitch (SIT) during a 50% MVC. Stimulator intensity was increased in 5% intervals from 50% stimulator output and two stimuli were delivered during an ~ 5 s isometric contraction, with the mean of two SITs recorded (Dekerle *et al.* **2019a**). Mean stimulator intensity was not different between males and females (65 ± 6 vs. $64 \pm 5\%$, $P = 0.791$) or between visits (66 ± 6 vs. $64 \pm 6\%$, $P = 0.100$). The intensities used, activated a large proportion of the motoneuron pool for the RF with no difference in the RF MEPs between trials at baseline (51 ± 15 vs. $53 \pm 11\% M_{max}$, $P = 0.314$). The TMS pulse also avoided substantial activation of the antagonist (BF), with small MEPs recorded at baseline (0.44 ± 0.23 vs. 0.47 ± 0.23 mV, $P = 0.476$).

Active motor threshold (AMT) was determined as the stimulator intensity that elicited an MEP of $> 200 \mu\text{V}$ in three out of five stimulations during a 10% MVC contraction. Stimulator intensity was increased in 5% steps from 35% of stimulator output until a consistent MEP amplitude $> 200 \mu\text{V}$ was found. Thereafter, stimulus intensity was reduced in 1% steps until the lowest intensity to elicit an MEP of $> 200 \mu\text{V}$ was found. Mean AMT was not different between males and females (43 ± 6 vs. $40 \pm 5\%$, $P = 0.392$), or between visits (42 ± 5 vs. $43 \pm 7\%$, $P = 0.245$). SICI was assessed with 10 paired- and 10 single-pulse stimulations delivered. Paired-pulse TMS consisted of a conditioning pulse at 70% of AMT, and a test pulse at 120% AMT, with an inter-stimulus

interval of 2 ms. Two sets of 10 stimuli were used, with a 10 s rest between contractions. All stimuli were delivered during a 10% contraction. This stimulus paradigm has previously been demonstrated as the optimal configuration for measuring SICl in the RF (Brownstein *et al.* **2018**).

Lumbar electrical stimulation

To assess spinal motoneuron excitability, lumbar-evoked potentials (LEPs) were measured with a constant-current stimulator (1 ms pulse duration; Digitimer DS7AH, Welwyn Garden City, UK). The cathode was centred over the first lumbar spinous process (5 × 9 cm; Nidd Valley Medical Ltd, Bordon, UK) with the electrode aligned to the centre of the vertebral column. The surface area of the cathode covered two spinous processes above and below the centre point (T11–L3). A cathode of large area was chosen as it produced less discomfort and greater tolerance by participants (Ugawa *et al.* **1995**; Kuhn *et al.* **2010**). The anode (2.5 cm²) was placed 5 cm above the upper edge of the cathode (Ugawa *et al.* **1995**), corresponding to the level of the eighth thoracic spinous process (T8) as this stimulating site has recently been shown to activate corticospinal axons at the level of lumbar spinal segments (Škarabot *et al.* **2019a**). The pre-exercise LEP was standardised to 15–25% of M_{\max} . Lumbar stimulation was performed during a 10% MVC contraction alone (unconditioned), and 100 ms into a 200 ms silent period (SP; conditioned) to determine excitability of the spinal cord without the presence of background neural drive (Finn *et al.* **2018**). The mean stimulus intensity for unconditioned LEPs was 172 ± 47 mA for males and 166 ± 24 mA for females ($P = 0.732$). For conditioned LEPs (SP-LEPs), the TMS intensity to produce an SP of 200 ms was not different between males and females (49 ± 8 vs. 51 ± 6%, $P = 0.605$), and likewise the intensity of subsequent lumbar stimulation was not different (176 ± 46 vs. 172 ± 22 mA, $P = 0.747$).

Motor nerve stimulation

Single electrical stimuli (200 μs duration) were delivered to the right femoral nerve using a constant current stimulator (DS7AH Digitimer Ltd) via adhesive surface electrodes (CF3200; Nidd Valley Medical Ltd, Harrogate, UK). The cathode was placed over the nerve, high in the femoral triangle, in the position that elicited the greatest twitch amplitude (Q_{tw}) and M-wave in the RF at rest. The anode was placed halfway between the greater trochanter and iliac crest. Optimum stimulus intensity was determined as the minimum current that elicited maximum values of Q_{tw} and M-wave (M_{\max}) at rest and then subsequently multiplied by 1.3 to ensure a supra-maximal stimulus was delivered. Mean stimulus intensity was not different between sexes (189 ± 62 vs. 210 ± 57 mA, $P = 0.438$) or between visits (194 ± 61 vs. 202 ± 61 mA, $P = 0.620$).

Force and electromyography

During assessments of neuromuscular function, participants were seated on a custom-built chair with knee and hip angles kept constant (both 90° flexion). A calibrated load cell (MuscleLab force sensor 300, Ergotest technology, Langesund, Norway) was attached via a non-compliant cuff positioned 2 cm superior to the ankle malleoli on the participants' right leg, to measure knee extensor force (N). EMG signals were recorded continuously throughout the final two trials using wireless sensors (10 mm inter-electrode distance; Trigno Avanti, Delsys, MA, USA). Sensors were placed over the right RF, VL and BF, consistent with SENIAM guidelines (Hermens *et al.* **2000**), as well as the sternocleidomastoid (SCM), and seventh intercostal space (IC). Prior to placement, the skin–electrode contact area was shaved, abraded and cleaned using a 70% IPA alcohol wipe (FastAid, Robinson Healthcare, Worksop, UK). Signals were amplified: gain × 100 for EMG (Delsys Trigno Wireless EMG systems, Boston, MA, USA) and × 300 for force (CED 1902; Cambridge Electronic Design, Cambridge, UK), bandpass filtered (EMG only: 20–450 Hz), digitized (EMG: 2 kHz; force: 5 kHz; CED 1401, Cambridge Electronic Design), and analysed offline (Spike2 v8, Cambridge Electronic Design).

Near infrared spectroscopy

A multi-distance, continuous-wave, single channel NIRS system (NIRO-200NX, Hamamatsu, Hamamatsu City, Japan) evaluated changes in VL oxy- (HbO₂) and deoxy- (HHb) haemoglobin concentrations (μmol l⁻¹), as well as

tissue oxygenation index ($\text{TOI} = \text{HbO}_2 \div [\text{HbO}_2 + \text{HHb}] \times 100$), sampled at a rate of 1 Hz. The light-emitting probe consisted of diodes operating at three wavelengths (735, 810 and 850 nm), and an emitter–detector distance of 3 cm. The probe was placed over the VL, 20 cm above the fibular head lateral side of the patella (Keane *et al.* **2018**). Optodes were held in place by an elasticised, tensor bandage and covered by an opaque, dark material to avoid motion and ambient light influences. During the fatiguing tasks, the 30 s window around 25, 50 and 75% of the task, as well as the final 30 s of the task (100%) were analysed.

Maximal inspiratory and expiratory pressure measurement

Maximum static inspiratory mouth pressure (MaxInsp) was measured from residual lung volume, while maximum static expiratory mouth pressure (MaxExp) was measured from total lung volume. Manoeuvres were performed using a handheld device (MicroRPM, CareFusion, Hampshire, UK) attached to a phlanged mouthpiece with a 1 mm leak to prevent glottic closure during the MaxInsp manoeuvre, and to reduce the use of buccal muscles during the MaxExp manoeuvre (American Thoracic Society/European Respiratory Society, **2002**). Measures were taken while participants were seated, with strong verbal encouragement given to maintain a maximal effort for ~3 s, and participants were given 30 s rest between efforts. Post-exercise values were taken immediately after the neuromuscular function assessments (~2.5 min after task termination). The largest of three values within 5% variability was used for analysis (Wen *et al.* **1997**). The coefficient of variation ($\text{CV} = [\text{SD} \div \text{mean}] \times 100$) between baseline assessments in the severe and heavy trials for MaxInsp was 3.7 and 4.6%, and the CV for MaxExp was 7.9 and 3.3% for males and females, respectively.

Data analysis

One female participant did not complete any assessment incorporating TMS (e.g. VA_{TMS} , MEP, SP-LEP or SICI) due to a contraindication (metal object in the skull), but she did complete all other measures. Voluntary activation using MNS was determined using the twitch interpolation method (Merton, **1954**) by comparing the amplitude of the superimposed twitch (SIT) with the amplitude of the potentiated resting twitch ($Q_{\text{tw.pot}}$) using the following formula: $\text{VA}_{\text{MNS}} (\%) = (1 - [\text{SIT} \div Q_{\text{tw.pot}}]) \times 100$. Voluntary activation using TMS (VA_{TMS}) was assessed during two sets of contractions at 100, 87.5, 75, 62.5 and 50% MVC (Dekerle *et al.* **2019b**). Single-pulse TMS was delivered during each contraction, and the linear regression between SIT amplitude and contraction intensity was extrapolated to the y intercept to obtain an estimated resting twitch (ERT; Todd *et al.* **2003**). To achieve significant linearity ($r^2 > 0.80$, $P < 0.05$), a total of three out of 720 SITs across all trials were excluded (0.4%), which led to three regressions containing nine data points rather than 10 (all after exercise). As a result, mean r^2 values for ERTs were linear throughout the study (0.92 ± 0.07). The SIT during 100% MVC was compared with the ERT using the following formula: $\text{VA}_{\text{TMS}} (\%) = (1 - [\text{SIT} \div \text{ERT}]) \times 100$.

Short-interval intracortical inhibition was quantified as the percentage ratio between the amplitude of conditioned MEPs to the amplitude of unconditioned MEPs. Corticospinal excitability was determined by expressing the mean MEP amplitude during the 10% MVC as a percentage of M_{max} ($\text{MEP}/M_{\text{max}}$). The rmsEMG was recorded for the 50 ms before each stimulation and compared before and after exercise to measure background muscle activity. The NIRS (O_2Hb , HHb and TOI) and gas exchange (\dot{V}_{O_2} , \dot{V}_{CO_2} , \dot{V}_{E} and RER) data were expressed as a percentage of baseline, and the 30 s epochs throughout exercise are presented as $\Delta\%$. Gas exchange data were also expressed as a percentage of final ramp test values, to facilitate comparisons between sexes.

Statistical analysis

Data are presented as mean \pm SD within the text and figures. A normal Gaussian distribution of data was confirmed using the Kolmogorov–Smirnov test. The significance level for all statistical tests was set at $P < 0.05$. For variables assessed before and during exercise (NIRS and gas exchange) a two-way (2×5) repeated-measures ANOVA was used to assess differences between sex (male vs. female) and over time (Pre, 25, 50, 75 and 100%

TTF). For variables assessed only during exercise (rmsEMG) a two-way (2×5) repeated-measures ANOVA was used to assess differences between sex (male vs. female) and over time (Start, 25, 50, 75 and 100% TTF). For variables that were assessed before and after exercise (neuromuscular function) a two-way (2×2) repeated-measures ANOVA was used to assess differences between sex (male vs. female) and over time (Pre vs. Post). If significant main or interaction effects were observed, these were followed up by *post hoc* Tukey's pairwise comparisons. Paired-samples *t* tests were performed to compare end-exercise \dot{V}_{O_2} to $\dot{V}_{O_{2max}}$ in both exercise intensity domains.

Results

Incremental ramp test

The variables recorded during the ramp test are displayed in Table 1. As shown, males recorded greater values for $\dot{V}_{O_{2max}}$, and P_{max} when expressed in absolute units and also when normalised to body mass (all $P < 0.001$). The average performance level (De Pauw *et al.* 2013, Decroix *et al.* 2016) was similar between males and females (3.4 ± 0.7 vs. 3.3 ± 0.5 , $P = 0.609$).

Table 1. Participant demographics, comparison of the results from the incremental exercise test, and power–duration relationship modelling in males and females

	Males	Females	P value
<i>N</i>	10	8	n/a
Age (years)	25 ± 5	25 ± 6	0.836
Stature (cm)	178 ± 9	169 ± 9	0.054
Mass (kg)	67.0 ± 8.8	63.3 ± 7.2	0.325
Training (h.week ⁻¹)	9 ± 3	10 ± 5	0.581
Incremental test			
$\dot{V}_{O_{2max}}$ (l min ⁻¹)	4.02 ± 0.47	2.85 ± 0.51	<0.001
$\dot{V}_{O_{2max}}$ (ml kg ⁻¹ min ⁻¹)	60.5 ± 8.2	45.1 ± 6.3	<0.001
P_{max} (W)	362 ± 29	241 ± 42	<0.001
P_{max} (W·kg ⁻¹)	5.5 ± 0.6	3.8 ± 0.5	<0.001
Power–duration relationship			
CP (W)	260 ± 28	179 ± 32	<0.001
CP (W·kg body mass)	3.9 ± 0.7	2.8 ± 0.5	<0.001
CP (% P_{max})	72 ± 5	74 ± 2	0.210
W' (J)	18,515 ± 4831	12,684 ± 3155	0.009
W' (J·kg body mass ⁻¹)	276 ± 65	197 ± 41	0.009
W' (J· P_{max} ⁻¹)	51 ± 11	52 ± 10	0.733
r^2	0.98 ± 0.02	0.96 ± 0.02	
CP SE (%)	2 ± 1	3 ± 2	
W' SE (%)	7 ± 4	8 ± 3	

Abbreviations: CP, critical power; P_{max} , maximal power; SE, standard error; $\dot{V}_{O_{2max}}$, maximal oxygen uptake; W' , curvature constant.

Power–duration relationship

The parameter estimates for the power–duration relationship are presented in Table 1. The range of TTF for the shortest estimation trial was 105–185 s, while the range for the longest trial was 568–1192 s. When data were expressed in absolute units, males demonstrated greater values than females ($P \leq 0.009$), but when CP and W' were normalised to P_{max} , no differences between the sexes were observed ($P \geq 0.210$).

Severe intensity exercise

Fatigability

All participants reached task failure, and there was no difference in TTF between sexes during the trial at 110% CP (males: 752 ± 329 vs. females: 681 ± 277 s, $P = 0.645$). The power–duration relationship accurately predicted TTF for both males and females, with no difference between predicted values (719 ± 213 vs. 713 ± 146 s, $P \geq 0.678$).

The changes in neuromuscular variables are displayed in Fig. 1. MVC, $Q_{tw.pot}$, VA_{MNS} and VA_{TMS} decreased after exercise at 110% CP ($P \leq 0.002$), and this decrease was less in females compared to males for $Q_{tw.pot}$ (sex \times time interaction: -36 ± 17 vs. $-15 \pm 10\%$, $F_{1,16} = 8.4$, $P = 0.010$, $\eta^2 = 0.344$). When the percentage change in $Q_{tw.pot}$ was normalised to W' , no sex difference was observed ($-2.1 \pm 1.3\%$ decline per kJ vs. $-1.2 \pm 0.8\%$ decline per kJ, $P = 0.119$).




Figure 1. Neuromuscular function before and after exercise at 110% CP

A, maximum voluntary contraction (MVC); B, potentiated quadriceps twitch ($Q_{tw.pot}$); C, voluntary activation with motor nerve stimulation (VA_{MNS}); D, voluntary activation with transcranial magnetic stimulation (VA_{TMS}). *A greater decrease in males than females ($P < 0.05$). Male data are presented in blue and female data in red. Dashed lines indicate individual participants and the solid lines indicate the group mean.

No other variables demonstrated sex \times time interaction effects ($P \geq 0.058$). The amplitude of MEPs, LEPs and SP-LEPs did not change from before to after exercise ($P \geq 0.094$, Fig. 2), and similarly, M_{max} did not change ($P = 0.980$). Maximum inspiratory and expiratory pressures decreased after exercise ($P \leq 0.005$), with no sex difference in the magnitude of decrease ($P \geq 0.565$, Table 2).




Figure 2. Indices of neural excitability before and after exercise at 110% CP

A, motor evoked potential (MEP) amplitude normalised to maximal compound action potential amplitude (MEP/M_{max}); B, lumbar evoked potential (LEP) amplitude normalised to M_{max} (LEP/M_{max}); C, conditioned lumbar evoked potential (SP-LEP); D, short-interval intracortical inhibition (SICI). Male data are presented in blue and female data in red. Dashed lines indicate individual participants and the solid lines indicate the group mean.

Table 2. Changes throughout exercise above critical power for pulmonary gas exchange, EMG and pulmonary function variables

				Severe intensity					Heavy intensity			
Time to task failure/termination (s)	Males			752 ± 329					3073 ± 835			
	Females			681 ± 277					2937 ± 964			
		Pulmonary gas exchange										
		Pre-exercise	25% TTF	50% TTF	75% TTF	100% TTF	Pre-exercise	25% TTF	50% TTF	75% TTF	100% TTF	
\dot{V}_{O_2} (% \dot{V}_{O_2max})	Males	19 ± 4	87 ± 6*	93 ± 5*	95 ± 7*	98 ± 4*	17 ± 3	76 ± 6*	78 ± 6*	78 ± 5*	81 ± 5*	
	Females	18 ± 3	82 ± 6*	87 ± 5*	93 ± 6*	98 ± 4*	18 ± 3	76 ± 7*	79 ± 7*	81 ± 7*	84 ± 6*	
\dot{V}_{CO_2} (% \dot{V}_{CO_2max})	Males	15 ± 2	76 ± 12*	78 ± 12*	79 ± 12*	81 ± 12*	15 ± 2*	64 ± 9*	63 ± 9*	64 ± 9*	65 ± 9*	
	Females	16 ± 2	77 ± 6*	82 ± 6*	84 ± 7*	86 ± 6*	16 ± 4*	63 ± 8*	63 ± 6*	66 ± 6*	68 ± 6*	
\dot{V}_E (% \dot{V}_{Emax})	Males	15 ± 3	76 ± 12*	78 ± 12*	79 ± 12*	81 ± 12*	14 ± 3	52 ± 8*	56 ± 8*	59 ± 7*	64 ± 10*	
	Females	16 ± 2	77 ± 6*	82 ± 6*	84 ± 7*	86 ± 6*	20 ± 2	66 ± 5*	68 ± 8*	72 ± 7*	77 ± 7*	
RER ($\dot{V}_{CO_2}/\dot{V}_{O_2}$)	Males	0.92 ± 0.04	1.01 ± 0.11	0.98 ± 0.09	0.95 ± 0.06	0.95 ± 0.07	1.00 ± 0.10	0.96 ± 0.05\$	0.94 ± 0.07	0.95 ± 0.07	0.93 ± 0.05\$	
	Females	0.95 ± 0.06	1.02 ± 0.06	1.00 ± 0.05	0.97 ± 0.06	0.94 ± 0.05	0.97 ± 0.14	0.90 ± 0.05	0.88 ± 0.06	0.89 ± 0.05	0.88 ± 0.05	
		Muscle activation										
		Start exercise	25% TTF	50% TTF	75% TTF	100% TTF	Start exercise	25% TTF	50% TTF	75% TTF	100% TTF	
(rmsEMG·M _{max} ⁻¹)	Males	3.3 ± 1.6	4.2 ± 2.0*	4.5 ± 2.2*	5.2 ± 2.0*	5.7 ± 1.9*	4.3 ± 3.8	4.7 ± 3.8	4.6 ± 3.3	5.3 ± 4.2*	5.1 ± 3.7*	
	Females	3.7 ± 1.4	4.5 ± 1.6*	5.0 ± 1.6*	5.0 ± 1.7*	5.2 ± 1.9*	2.9 ± 1.1	3.2 ± 1.2	3.4 ± 1.2	3.4 ± 1.5	3.4 ± 1.4	
(% rmsMaxInsp)	Males	11.6 ± 8.8	18.4 ± 9.4	26.2 ± 14.1	33.3 ± 16.0	44.5 ± 22.2	10.6 ± 8.0	11.6 ± 8.2	11.9 ± 8.8	15.1 ± 13.6	15.1 ± 11.1	
	Females	16.9 ± 10.5	23.4 ± 12.3	24.9 ± 10.0	28.4 ± 11.7	36.2 ± 11.7	12.9 ± 5.0	19.9 ± 12.8	19.9 ± 12.3	17.9 ± 9.4	19.5 ± 10.8	
External intercostal (% rmsMaxExp)	Males	19.5 ± 9.9	27.9 ± 13.4	34.4 ± 16.3	37.1 ± 17.9	49.6 ± 26.1	10.3 ± 6.5	12.6 ± 13.5	13.5 ± 6.5	13.6 ± 7.8	15.2 ± 7.1	
	Females	38.5 ± 18.4	55.5 ± 38.9	54.9 ± 38.9	62.15 ± 38.6	61.0 ± 34.7	24.9 ± 8.3	31.0 ± 11.7	30.2 ± 12.5	29.5 ± 14.2	33.9 ± 17.6	
		Maximal pulmonary pressures										
			Pre-exercise				Post-exercise		Pre-exercise		Post-exercise	

Maximum expiratory pressure (mmHg)	Males		197 ± 52			171 ± 48*			174 ± 39		157 ± 31
	Females		143 ± 37			129 ± 38*			138 ± 34		136 ± 40
Maximum inspiratory pressure (mmHg)	Males		130 ± 37			118 ± 33*			140 ± 53		135 ± 50*
	Females		113 ± 29			104 ± 25*			138 ± 34		136 ± 40

*Significantly different from before exercise ($P < 0.05$). [§]significantly different from females. Abbreviations: MaxExp, maximal expiratory pressure; MaxInsP, maximal inspiratory pressure; RER, respiratory exchange ratio; rmsEMG· M_{\max}^{-1} , root-mean-square EMG activity normalised to maximal compound action potential amplitude; TTF, time to task failure; \dot{V}_E , minute ventilation; \dot{V}_{O_2} , respiratory oxygen uptake; \dot{V}_{CO_2} , carbon dioxide production.

Oxygenation

Both HbO₂ and TOI decreased throughout severe intensity exercise (Fig. 3, time effect $P < 0.001$), whilst HHb increased ($P = 0.017$). A lesser decrease in TOI (sex \times time interaction: $F_{1.3,21.1} = 16.6$, $P < 0.001$, $\eta^2 = 0.509$) was observed for females compared to males, as well as a reduced increase in HHb (sex \times time interaction: $F_{1.7,26.5} = 5.3$, $P = 0.024$, $\eta^2 = 0.254$).




Figure 3. Indices of muscle oxygenation throughout exercise at 110% CP

A, oxyhaemoglobin (O₂Hb); B, deoxyhaemoglobin (HHb); C, tissue oxygenation index (TOI). * Greater in males than in females ($P < 0.05$). Male data are presented in blue and female data in red. Dashed lines indicate individual participants and the solid lines indicate the group mean.

Respiratory and locomotor muscle electromyography

The rmsEMG for VL, SCM and IC all increased throughout the task (Table 2, all time effects $P < 0.001$). Females demonstrated a lesser increase in rmsEMG for the VL (sex \times time interaction: $F_{4,64} = 2.7$, $P = 0.041$, $\eta^2 = 0.142$), but not for the SCM ($P = 0.079$) or IC ($P = 0.255$).

Pulmonary gas exchange

Oxygen consumption, \dot{V}_{CO_2} and \dot{V}_E increased while RER decreased throughout the task (Table 2, $P < 0.001$). The sex \times time interaction effect for \dot{V}_E was not significant ($P = 0.052$), and no other sex differences were observed ($P \geq 0.114$). End-exercise \dot{V}_{O_2} was not significantly different from the $\dot{V}_{O_{2max}}$ measured during the incremental test ($P = 0.442$).

Heavy intensity

Fatigability

Three males (1843 ± 498 s) and three females (1831 ± 568 s) reached task failure prior to the 60 min (3600 s) cut-off and were included in subsequent analyses (whole group mean duration, 3073 ± 835 vs. 2937 ± 964 s, $P = 0.758$).

There were significant decreases in MVC, $Q_{tw.pot}$, VA_{MNS} , VA_{TMS} , MEP and SP-LEP following exercise at 90% CP ($P \leq 0.039$, Figs 4 and 5). Females demonstrated less of a decrease in $Q_{tw.pot}$ (sex \times time interaction: -10 ± 11 vs. $-24 \pm 11\%$, $F_{1,16} = 31.8$, $P = 0.020$, $\eta^2 = 0.655$) and VA_{MNS} (sex \times time interaction: -4 ± 3 vs. $-9 \pm 6\%$, $F_{1,16} = 5.2$, $P = 0.036$, $\eta^2 = 0.246$) compared to males, but no sex difference was demonstrated for VA_{TMS} , MEP and SP-LEP ($P \geq 0.051$). No change in M_{max} was observed following exercise ($P = 0.980$). Maximum inspiratory pressure decreased after exercise ($P = 0.001$), whereas maximum expiratory pressure did not ($P = 0.063$, Table 2). No sex \times time interaction in the magnitude of decrease for the former was observed ($P = 1.000$).




Figure 4. Neuromuscular function changes relative to baseline for exercise at 90% CP

A, maximum voluntary contraction (MVC); B, potentiated quadriceps twitch ($Q_{tw.pot}$); C, voluntary activation with motor nerve stimulation (V_{Amns}); D, voluntary activation with transcranial magnetic stimulation (V_{Atms}). * Greater decrease in males than in females ($P < 0.05$). Male data are presented in blue and female data in red. Dashed lines indicate individual participants and the solid lines indicate the group mean.




Figure 5. Indices of neural excitability before and after exercise at 90% CP

A, motor evoked potential (MEP) amplitude normalised to maximal compound action potential amplitude (MEP/M_{max}); B, lumbar evoked potential amplitude normalised to M_{max} amplitude (LEP/M_{max}); C, conditioned lumbar evoked potential amplitude (SP-LEP); D, short-interval intracortical inhibition (SICI). Male data are presented in blue and female data in red. Dashed lines indicate individual participants and the solid lines indicate the group mean.

Oxygenation

Decreases in HbO_2 ($P < 0.019$) and TOI ($P < 0.001$) were observed during heavy exercise, with females demonstrating less of a decrease for TOI (sex \times time interaction: $F_{1.7,26.9} = 41.0$, $P < 0.001$, $\eta^2 = 0.719$). An increase in HHb was observed for both sexes ($P = 0.008$), with females demonstrating less of an increase than males (Fig. 6, sex \times time interaction: $F_{1.4,22.8} = 20.8$, $P < 0.001$, $\eta^2 = 0.565$).




Figure 6. Indices of muscle oxygenation throughout exercise at 90% CP

A, oxyhaemoglobin (O₂Hb); B, deoxyhaemoglobin (HHb); C, tissue oxygenation index (TOI). * Greater in males than in females ($P < 0.05$). Male data are presented in blue and female data in red. Dashed lines indicate individual participants and the solid lines indicate the group mean.

Respiratory and locomotor muscle electromyography

The rmsEMG signal from the VL, SCM and IC all increased throughout the task (Table 2, all time effects $P \leq 0.033$). However, there were no sex differences in the rate of increase for any muscle (sex \times time interactions: $P \geq 0.063$).

Pulmonary gas exchange

Oxygen consumption, \dot{V}_{O_2} and \dot{V}_E all increased throughout the heavy intensity exercise task ($P < 0.001$, Table 2) while RER decreased ($P = 0.001$). A sex \times time interaction was not observed for \dot{V}_E ($P = 0.052$), but was for RER ($F_{1,16} = 5.08$, $P = 0.039$, $\eta^2 = 0.241$; see Table 2). *Post hoc* tests indicated that females had a lower RER (mean difference: -0.05) throughout the trial. End-exercise \dot{V}_{O_2} was $19 \pm 5\%$ lower than $\dot{V}_{O_{2max}}$ ($P < 0.001$).

Discussion

The present study explored the sex difference in fatigability during locomotor exercise by comparing the power–duration relationship, then muscle oxygenation and neuromuscular responses to CP-matched exercise intensities. The novel findings were that, while males demonstrated a greater absolute critical power and W' , there was no sex difference when these parameters were normalised to the absolute maximal power (P_{max}). Time to task failure/completion was not different in either heavy or severe exercise intensity domains, but females demonstrated lesser reductions in knee-extensor contractile function ($Q_{tw,pot}$) immediately after exercise. These sex differences are probably related to differences in skeletal muscle size and composition influencing the physiological response to exercise in both intensity domains. The change in corticospinal excitability appeared to be domain- but not sex-specific, with a decrease in MEP and SP-LEP amplitude observed in the heavy domain for both males and females. Together, this integrative data set suggests that the mechanism(s) for greater resistance to neuromuscular fatigue in females reside within the musculature.

Incremental test and power–duration relationship

As expected, males produced greater absolute power outputs during cycling for variables such as P_{max} and CP. Female P_{max} was 66% of male values, similar to previous reports (68%; Sundberg *et al.* 2016). This sex difference in maximum power was still evident when P_{max} was normalised to body mass ($W \text{ kg}^{-1}$), probably a result of differences in body composition (Pate & O'Neill, 2007). A similar sex difference was observed for $\dot{V}_{O_{2max}}$, with males having greater absolute values and values relative to body weight (Pate & O'Neill, 2007). However, allometric scaling of $\dot{V}_{O_{2max}}$ to fat-free mass typically eliminates some of this sex difference, with differences in haemoglobin mass explaining the remainder of the sex differences (Joyner, 2017). The normalisation of CP with P_{max} presents a method of making inter-individual comparisons of the power–duration relationship. Indeed, when relative CP (% P_{max}) was compared, no sex difference was demonstrated. Previously, Sundberg *et al.* (2016) demonstrated no sex difference when profiling the power–duration relationship during cycling bouts across the extreme and severe intensity domains (bout durations: 8–283 s), as well as the 3 min 'all-out' test. The present study corroborates this evidence and extends the conclusion to when CP is assessed using multiple severe intensity exercise trials to exhaustion. Collectively, these data obtained from cycling assessments conflict with equivalent data obtained in an isometric exercise setting where females had a $\sim 7\%$ greater critical torque compared to males (Ansdell *et al.* 2019a,b). This discrepancy is probably explained by the modality of exercise.

Critical power during cycling is considered to be limited by oxygen delivery to the working muscle, a result of convective and diffusive capacity (Vanhatalo *et al.* **2010**; Dekerle *et al.* **2012**; Broxterman *et al.* **2015a**; Goulding *et al.* **2017**), and therefore, during single-limb exercise, where convective factors are not a limiting factor (i.e. Ansdell *et al.* **2019a**), the sex difference in critical torque could be a result of greater diffusive capacity of female muscle. For example, it is well established that females have greater capillarisation and type I fibre proportional area of the knee-extensors (Roepstorff *et al.* **2006**), which could permit a greater rate of oxygen extraction and utilisation, and a greater relative critical torque. Whereas during locomotor exercise in the present study, knee-extensor blood flow is limited due to the modality of exercise (Calbet, **2000**), and therefore convective capacity becomes the primary determinant of CP, rather than diffusive capacity of the muscle, leading to the lack of sex difference in relative CP. Other contributing factors to the lack of sex difference in CP could be that haemoglobin concentrations are typically ~12% lower (Murphy, **2014**), and lung volumes are smaller (Schwartz *et al.* **1988**) in females than in males. The negative consequence of these factors is that females are more prone to exercise-induced arterial hypoxaemia (Harms *et al.* **1998**), meaning that when cardiac output is near maximal (i.e. at task failure within the severe intensity domain), there is no possibility for increased oxygen extraction. This leads to a reduced arterio-venous oxygen difference, which has been suggested to negate the sex difference in muscle fatigability (Dominelli *et al.* **2017**) and could conceivably oppose the positive aspects of greater type I muscle fibre proportion on CP (Mitchell *et al.* **2018a**), leading to a lack of sex difference.

Severe intensity exercise

The power–duration relationship successfully predicted TTF in the exercise trial at 110% CP (Table **2**), and the \dot{V}_{O_2} response to exercise confirmed that exercise was indeed in the severe intensity domain. The end-exercise \dot{V}_{O_2} was equivalent with $\dot{V}_{O_{2max}}$ and demonstrated a gradual increase throughout exercise, indicating the presence of a considerable slow component. Fatigability within the severe intensity domain is consistently associated with depletion of high energy phosphates and an accumulation of metabolites within the exercising musculature (Jones *et al.* **2008**; Black *et al.* **2016**; Vanhatalo *et al.* **2016**), which may reduce the contractile capacity of exercised musculature until the attainment of a limiting degree of disruption (Amann, **2011**; Burnley & Jones, **2018**). In contrast to our hypothesis, and despite no difference in TTF when exercise intensity was CP-matched, females demonstrated greater fatigue resistance of the knee-extensors compared to males immediately after the exercise (21% difference in $Q_{tw,pot}$ reduction). There are multiple factors that could explain this sex difference from the present study and previous data. For example, as previously mentioned, females typically have a greater proportional area of type I muscle fibres (Staron *et al.* **2000**; Roepstorff *et al.* **2006**), and whilst in the context of this study it might not contribute to differences in the power–duration relationship, it could provide females with the capacity to tolerate deleterious metabolites when exercising above CP. Similarly, previous studies using ^{31}P MRS during ‘all-out’ exercise have shown lower decreases in muscle pH, phosphocreatine and attenuated increases in ADP (Russ *et al.* **2005**; Willcocks *et al.* **2010**). Currently, it is unknown whether a lesser accumulation of metabolites occurs in the severe intensity domain for females, or whether the sex differences in contractile properties allows lesser peripheral fatigue for equivalent metabolic stress. Slower sarcoplasmic reticulum calcium ATPase and uptake activity in females (Harmer *et al.* **2014**) could reflect a more fatigue-resistant contractile apparatus. Additionally, females demonstrated a smaller increase in rmsEMG during the severe intensity task, and while this has some limitations as a measure of neural activity (Farina *et al.* **2004**), it could represent a reduced rate of increase in neural drive because of less fatigue within the already recruited motor units than males (Vigotsky *et al.* **2018**). Indeed, this finding mirrors previous data (Ansdell *et al.* **2019a**), and exists when rmsEMG is normalised to M_{max} to negate the influence of subcutaneous fat on the EMG signal (Lanza *et al.* **2018**).

Another potential explanation of the lesser degree of $Q_{tw.pot}$ decrease in females could be unearthed when exercise is considered using absolute (i.e. 286 vs. 197 W), rather than relative (110% CP) values, given that the power–duration relationship was not different between sexes (i.e. similar relative values, Table 1). The power output for CP, and W' values were both $\sim 30\%$ lower for females compared to males, and therefore TTF was not different, despite the lower amount of work done for females. Evidence suggests that W' is positively related to the cross-sectional area (CSA) of the exercising muscle (Miura *et al.* 2002; Kordi *et al.* 2018), and whilst the present study did not measure CSA of the thigh musculature, it is established that females have 25–30% smaller knee extensors and flexors (Behan *et al.* 2018). Similarly, Schäfer *et al.* (2019) demonstrated a positive relationship between W' and $Q_{tw.pot}$ decrease following severe intensity exercise. Therefore, in the present study, it is possible that the larger muscle CSA in males probably permitted a greater absolute W' , which consequently elicited a greater decrease in $Q_{tw.pot}$ when severe intensity exercise was performed to task failure. Indeed, when the decrease in $Q_{tw.pot}$ was normalised to W' , no sex difference was observed, providing some support to the possibility that the sex difference in $Q_{tw.pot}$ decrease was a result of the greater absolute workload performed by males.

A final potential factor contributing to the sex difference observed in $Q_{tw.pot}$ decline could be the greater oxygenation within the VL for females during exercise (Fig. 3). In both Ansdell *et al.*'s (2019a) and the present study, this manifested predominantly as a lesser rise in HHb concentration for females during both heavy and severe intensity cycling, which could be a result of the fibre type difference between males and females. Specifically, the greater HHb increase in males could be a result of greater oxygen extraction (Grassi *et al.* 2003), which could be related to a greater oxygen cost within the muscle ($m\dot{V}_{O_2}$) compared to females. Indeed, when assessed at a pulmonary level, individuals with greater type I fibre proportion of the VL demonstrate a lower \dot{V}_{O_2} for a given exercise intensity (Coyle *et al.* 1992). This is speculative, although potentially fertile ground for future research as $m\dot{V}_{O_2}$ can be non-invasively quantified with a combination of NIRS and muscle occlusion (Ryan *et al.* 2012); a combined approach to pulmonary and muscle \dot{V}_{O_2} kinetics could permit further insight into the integrative response to exercise in males and females (Poole & Jones, 2012). One might expect pulmonary \dot{V}_{O_2} to reflect a potential sex difference in $m\dot{V}_{O_2}$; however, females experienced a similar \dot{V}_E to males during severe intensity cycling, which has previously been linked to a greater oxygen cost of breathing in females (Witt *et al.* 2007). When measured at the pulmonary level, the \dot{V}_{O_2} response to exercise is an amalgamation of all physiological systems, so the elevated W_b might have counterbalanced the reduced $m\dot{V}_{O_2}$ in females, leading to no sex difference in pulmonary \dot{V}_{O_2} values attained in the present study. Together, the aforementioned data present evidence that whilst exercise performance (TTF) in the severe intensity domain is not affected by sex, the integrative response differs between males and females. Females experience a lesser decline in $Q_{tw.pot}$, potentially because of differences in muscle oxygenation and Ca^{2+} kinetics. Despite this, females probably have a greater W_b when exercise intensity is CP-matched. Collectively, these data imply that even though TTF was not different, the mechanisms underpinning severe intensity exercise tolerance might differ between males and females.

Although a reduction in voluntary activation occurred during this trial for both sexes, excitability of the corticospinal tract was unaltered at the cortical and spinal level, suggesting that responsiveness of descending neurons did not change after exercise (Weavil & Amann, 2018). Therefore, the CNS adjustments might have been a result of impaired neural drive, or synaptic input into the corticospinal tract (Amann, 2011). Regardless, these central adjustments are not considered to be the limiting factor to exercise within the severe intensity domain during cycling (Burnley & Jones, 2018). One caveat of the present study and other locomotor neuromuscular fatigue studies is that responses were assessed after exercise during an isometric contraction (Sidhu *et al.* 2013; Place & Millet, 2020). Responses evoked *during* exercise could elucidate further details about the time-course and magnitude of fatigue-related changes in activation.

Heavy intensity domain

The \dot{V}_{O_2} response to exercise at 90% CP was typical of heavy intensity exercise. The \dot{V}_{O_2} response exhibited a slow component, but only reached $\sim 83\% \dot{V}_{O_2\max}$ at task termination, indicating that energy provision from aerobic sources was not maximal (i.e. exercise intensity was less than CP). In terms of the before to after exercise change in neuromuscular function, the fatigue observed was not due to an accumulation of disruptive metabolites, or an exhaustion of high-energy phosphates as substrate-level phosphorylation reaches a steady-state (Black *et al.* **2016**; Vanhatalo *et al.* **2016**). Rather, neuromuscular fatigue in the heavy intensity domain is a result of both central and peripheral adjustments, with the latter occurring in response to depletion of intramuscular glycogen concentration and the associated negative consequences for excitation–contraction coupling (Ørtenblad *et al.* **2013**). Furthermore, reactive oxygen species generation and extracellular accumulation of K^+ might also impair contractile function (Allen *et al.* **2008**). The net result in the present study is a decrease in $Q_{tw,pot}$ (Fig. **4B**), which was less profound in females. Given that the mechanisms of peripheral adjustments differ above and below CP, this greater fatigue resistance of female knee-extensors below CP must be a result of different physiological processes as well. One explanation could be that, given RER was lower in females compared to males during the 90% CP trial, the rate of fatty acid utilisation as a substrate was greater, eliciting a glycogen-sparing effect. This notion is supported by previous evidence demonstrating that males utilise $\sim 25\%$ more muscle glycogen at exercise intensities matched below CP (Tarnopolsky *et al.* **1990**; Roepstorff *et al.* **2002, 2006**; Devries *et al.* **2006**). The greater reliance on fat oxidation in females (RER), yet similar exercise economy (\dot{V}_{O_2}) between males and females also implies a lower oxygen cost of exercise, as fat oxidation is less efficient compared to carbohydrate oxidation. To further support this, and similar to 110% CP, the decrease in muscle oxygenation was less in females at 90% CP, potentially reflecting a lower oxygen cost of contraction as a result of greater type I muscle fibre proportion.

The CNS adjustments occurring below CP are thought not to be a result of group III/IV afferent feedback, as there is no progressive metabolite accumulation (Burnley & Jones, **2018**); instead, repetitive activation of motoneurons can alter their intrinsic properties, rendering them less responsive to activation (Carpentier *et al.* **2001**). This phenomenon is reflected in the present study as a decrease in VA_{MNS} and VA_{TMS} , with a greater decline in VA_{MNS} only for males. This discrepancy might indirectly suggest that the aetiology of the sex difference in central fatigue would be located at a sub-cortical level. Indeed, a decrease in MEP and SP-LEP was observed (Fig. **5**) and is probably a result of reduced strength of persistent inward currents (Heckman *et al.* **2008**). However, the sex \times time interaction for these evoked variables was not significant ($P \geq 0.132$). Multiple studies have provided evidence to show reduced motoneuronal excitability with fatigue in single-limb (Kennedy *et al.* **2016**; Finn *et al.* **2018**) and whole-body exercise modalities (Weavil *et al.* **2016**; Sidhu *et al.* **2017**), but the present study is the first to match exercise intensity to CP and assess the neural response. Interestingly, the decrease in LEP was only evident during the SP, with no change in unconditioned LEP (Fig. **5**). Finn *et al.* (**2018**) demonstrated a similar phenomenon in an isometric modality and suggested that SP-LEPs were more sensitive to intrinsic changes in motoneuronal properties, as inhibiting descending drive from the motor cortex (i.e. the TMS-SP; Škarabot *et al.* **2019b**) removes a confound of excitatory synaptic input to the motoneuron. The present data support this notion, as the unconditioned LEPs did not change, due to the compensatory effects of neural drive (rmsEMG in Table **2**) on net motoneuronal output. Therefore, as only SP-LEPs changed, the central fatigue observed at 90% CP in the present study is likely to be a result of a change in intrinsic properties of motoneurons, rendering them less responsive to synaptic input. As mentioned above, this occurred independent of sex for any evoked responses. It is possible that due to far smaller measurement error for VA_{MNS} compared to evoked potentials (Ansdell *et al.* **2019c**), a sex difference in motoneuronal excitability was not discernible due to a lack of statistical power. Collectively, these data suggested that the neuromuscular response to cycling at 90% CP is underpinned by decreases in CNS function and contractile impairment. Similar to severe intensity exercise, there was no sex difference in exercise duration, but the neuromuscular

adjustments were different between males and females, with greater oxygenation and less contractile dysfunction observed in females.

Further considerations

Fatigability of both inspiratory and expiratory muscle groups was demonstrated above CP, which was not sex-dependent. This contrasts with previous evidence suggesting that the diaphragm is a more fatigue-resistant muscle in females (Guenette *et al.* **2010**; Welch *et al.* **2018**), although the assessment modality employed in the present study was not able to provide information on the individual muscles or mechanisms responsible for the reduced pressures observed after exercise. The rise in rmsEMG for respiratory musculature was similar between sexes in both trials, which also contradicts previous findings suggesting females activate 'accessory' respiratory muscle such as the SCM to reduce the diaphragmatic load (Mitchell *et al.* **2018b**). Whilst no sex differences in respiratory muscle fatigability or gas exchange were observed in the present study, the similar \dot{V}_E values attained in the present study in males and females during both severe and heavy intensity cycling probably led to females experiencing a greater relative work of breathing during both trials (Dominelli *et al.* **2015**), which could contribute to greater exertional dyspnoea (Schaeffer *et al.* **2014**; Cory *et al.* **2015**). When taken into consideration with the lesser degree of peripheral adjustments in locomotor muscles in females, it could conceivably be suggested that the 'sensory tolerance limit' consists of different magnitudes of afferent feedback from different physiological systems in males and females (Hureau *et al.* **2018**; Thomas *et al.* **2018**), such that the locomotor muscle component is less, but the respiratory component is greater in females (Cory *et al.* **2015**).

To compare fatigability in different populations, it is necessary to match both the intensity of exercise and the training status of the populations. The former was addressed in the present study by normalising exercise intensity to CP. Attempts were made to recruit populations of males and females of equivalent training status (De Pauw *et al.* **2013**; Decroix *et al.* **2016**), which resulted in similar average performance levels between groups. However, the sex difference in relative $\dot{V}_{O_{2max}}$ was $\sim 25\%$. This is larger than the sex difference suggested for sexes of equivalent training status ($\sim 10\%$, Joyner **2017**), although this was based on a mixture of studies and a small sample of $n = 8$ male and 15 female elite distance runners (Pate & O'Neill, **2007**). Other sources have previously described larger magnitudes in this sex difference (e.g. 17%, Froberg & Pedersen, **1984**), although similarly rely on small sample sizes ($n = 6$ females and $n = 7$ males). Indeed, a meta-analysis of 440 male and 381 female participants demonstrated an average sex difference of 28% in $\dot{V}_{O_{2max}}$ when expressed relative to body mass; this difference remained in trained vs. untrained populations when body composition was accounted for (Sparling, **1980**). Nevertheless, there appears to be a discrepancy in what researchers deem to be an appropriate magnitude for the sex difference in $\dot{V}_{O_{2max}}$. The present study used a minimum performance level (De Pauw *et al.* **2013**; Decroix *et al.* **2016**) to account for $\dot{V}_{O_{2max}}$, relative P_{max} , as well as training history (hours week⁻¹), and the sex differences demonstrated are therefore assumed to be independent of training status. However, as is well established, training status influences aerobic fitness, so this discrepancy in the appropriate magnitude of sex difference in $\dot{V}_{O_{2max}}$ highlights a potential limitation in the present study, if the differences in indices of aerobic fitness are considered to be of too great a magnitude. The precise measurement of, and normalisation of values to, fat-free mass could be an area for future research in order to uncouple the effects of sex and muscle mass in the field of integrative exercise physiology.

Finally, NIRS signals can be influenced by subcutaneous adipose tissue thickness, which manifests as a reduction in the concentration of haeme compounds (Van Beekvelt *et al.* **2001**; Bopp *et al.* **2011**; Bowen *et al.* **2013**). It is well established that females typically have a greater amount of subcutaneous adipose tissue (Westerbacka *et al.* **2004**). The system used in the present study was a spatially resolved spectroscopy one, which enhances the signal from deeper tissues, whilst reducing the contribution from superficial tissues (i.e. skin and subcutaneous fat; Messere & Roatta, **2013**). Additionally, this form of NIRS system provides a relative index

of tissue oxygenation (TOI), in which both the numerator (HbO₂) and denominator (HbO₂ + HHb) are equally affected by adipose tissue thickness, and therefore a correction might not be necessary (Barstow, 2019). Despite adipose tissue's established effects on HbO₂ and HHb values, it is currently unknown whether this also affects the sensitivity of the technique to changes induced by exercise. Therefore, the NIRS data presented in this study must be considered within this context.

Conclusions

This study demonstrated that the power–duration relationship for cycling did not differ between males and females when expressed relative to P_{max}. Subsequent exercise performance in the severe and heavy intensity domains was not different, but the integrative response of cardiopulmonary, respiratory and neuromuscular systems differed. Specifically, muscle de-oxygenation and contractile impairment was less in females during both tasks, potentially related to skeletal muscle size and composition. Additionally, the decline in CNS function was attenuated for females in the heavy intensity domain. Collectively, the present data show that the integrative response of physiological systems differs between males and females, which has important implications for acute and chronic exercise prescription.

References

- Allen DG, Lamb GD & Westerblad H (2008). Skeletal muscle fatigue: cellular mechanisms. *Physiol Rev* **88**, 287– 332.
- Amann M (2011). Central and peripheral fatigue: interaction during cycling exercise in humans. *Med Sci Sports Exerc* **43**, 2039– 2045.
- American Thoracic Society/European Respiratory Society (2002). ATS/ERS statement on respiratory muscle testing. *Am J Respir Crit Care Med* **166**, 518– 624.
- Ansdell P, Brownstein CG, Škarabot J, Hicks KM, Howatson G, Thomas K, Hunter SK & Goodall S (2019a). Sex differences in fatigability and recovery relative to the intensity–duration relationship. *J Physiol* **597**, 5577– 5595.
- Ansdell P, Brownstein CG, Škarabot J, Hicks KM, Howatson G, Thomas K, Hunter SK & Goodall S (2019b). Methodological issues influence determination of critical force during intermittent exercise: authors' reply. *J Physiol* **597**, 5987– 5989.
- Ansdell P, Brownstein CG, Škarabot J, Hicks KM, Simoes DCM, Thomas K, Howatson G, Hunter SK & Goodall S (2019c). Menstrual cycle-associated modulations in neuromuscular function and fatigability of the knee extensors in eumenorrhic women. *J Appl Physiol* **126**, 1701– 1712.
- Ansdell P, Thomas K, Howatson G, Hunter S & Goodall S (2017). Contraction intensity and sex differences in knee-extensor fatigability. *J Electromyogr Kinesiol* **37**, 68– 74.
- Barstow TJ (2019). Understanding near infrared spectroscopy and its application to skeletal muscle research. *J Appl Physiol* **126**, 1360– 1376.
- Van Beekvelt MCP, Colier WNJM, Wevers RA & Van Engelen BGM (2001). Performance of near-infrared spectroscopy in measuring local O₂ consumption and blood flow in skeletal muscle. *J Appl Physiol* **90**, 511– 519.
- Behan FP, Maden-Wilkinson TM, Pain MTG & Folland JP (2018). Sex differences in muscle morphology of the knee flexors and knee extensors. *PLoS One* **13**, e0190903.
- Black MI, Bowtell JL, McDonagh STJ, Blackwell JR, Kelly J, Bailey SJ, Thompson C, Jones AM, Wylie LJ, Mileva KN, Sumners P & Vanhatalo A (2016). Muscle metabolic and neuromuscular determinants of fatigue during cycling in different exercise intensity domains. *J Appl Physiol* **122**, 446– 459.
- Bopp CM, Townsend DK & Barstow TJ (2011). Characterizing near-infrared spectroscopy responses to forearm post-occlusive reactive hyperemia in healthy subjects. *Eur J Appl Physiol* **111**, 2753– 2761.

- Bowen TS, Rossiter HB, Benson AP, Amano T, Kondo N, Kowalchuk JM & Koga S (2013). Slowed oxygen uptake kinetics in hypoxia correlate with the transient peak and reduced spatial distribution of absolute skeletal muscle deoxygenation. *Exp Physiol* **98**, 1585– 1596.
- Brownstein CG, Ansdell P, Škarabot J, Howatson G, Goodall S & Thomas K (2018). An optimal protocol for measurement of corticospinal excitability, short intracortical inhibition and intracortical facilitation in the rectus femoris. *J Neurol Sci* **394**, 45– 56.
- Broxterman RM, Ade CJ, Craig JC, Wilcox SL, Schlup SJ & Barstow TJ (2015a). Influence of blood flow occlusion on muscle oxygenation characteristics and the parameters of the power–duration relationship. *J Appl Physiol* **118**, 880– 889.
- Broxterman RM, Craig JC, Smith JR, Wilcox SL, Jia C, Warren S & Barstow TJ (2015b). Influence of blood flow occlusion on the development of peripheral and central fatigue during small muscle mass handgrip exercise. *J Physiol* **593**, 4043– 4054.
- Burnley M & Jones AM (2018). Power–duration relationship: physiology, fatigue, and the limits of human performance. *Eur J Sport Sci* **18**, 1– 12.
- Burnley M, Vanhatalo A & Jones AM (2012). Distinct profiles of neuromuscular fatigue during muscle contractions below and above the critical torque in humans. *J Appl Physiol* **113**, 215– 223.
- Calbet JA (2000). Oxygen tension and content in the regulation of limb blood flow. *Acta Physiol Scand* **168**, 465– 472.
- Carpentier A, Duchateau J & Hainaut K (2001). Motor unit behaviour and contractile changes during fatigue in the human first dorsal interosseus. *J Physiol* **534**, 903– 912.
- Carroll TJ, Taylor JL & Gandevia SC (2016). Recovery of central and peripheral neuromuscular fatigue after exercise. *J Appl Physiol* **122**, 1068– 1076.
- Cory JM, Schaeffer MR, Wilkie SS, Ramsook AH, Puyat JH, Arbour B, Basran R, Lam M, Les C, Macdonald B, Jensen D & Guenette JA (2015). Sex differences in the intensity and qualitative dimensions of exertional dyspnea in physically active young adults. *J Appl Physiol* **119**, 998– 1006.
- Coyle EF, Sidossis LS, Horowitz JF & Beltz JD (1992). Cycling efficiency is related to the percentage of Type I muscle fibers. *Med Sci Sports Exerc* **24**, 782– 788.
- Crapo RO, Morris AH & Gardner RM (1982). Reference values for pulmonary tissue volume, membrane diffusing capacity, and pulmonary capillary blood volume. *Bull Eur Physiopathol Respir* **18**, 893– 899.
- Decroix L, De Pauw K, Foster C & Meeusen R (2016). Guidelines to classify female subject groups in sport-science research. *Int J Sports Physiol Perform* **11**, 204– 213.
- Dekerle J, Ansdell P, Schäfer L, Greenhouse-Tucknott A & Wrightson J (2019a). Methodological issues with the assessment of voluntary activation using transcranial magnetic stimulation in the knee extensors. *Eur J Appl Physiol* **119**, 991– 1005.
- Dekerle J, Greenhouse-Tucknott A, Wrightson J, Schafer L & Ansdell P (2019b). Improving the measurement of TMS-assessed voluntary activation in the knee extensors. *PLoS One* **14**, e0216981.
- Dekerle J, Mucci P & Carter H (2012). Influence of moderate hypoxia on tolerance to high-intensity exercise. *Eur J Appl Physiol* **112**, 327– 335.
- Devries MC, Hamadeh MJ, Phillips SM & Tarnopolsky MA (2006). Menstrual cycle phase and sex influence muscle glycogen utilization and glucose turnover during moderate-intensity endurance exercise. *Am J Physiol Regul Integr Comp Physiol* **291**, R1120– R1128.
- Dominelli PB, Molgat-Seon Y, Griesdale DEG, Peters CM, Blouin JS, Sekhon M, Dominelli GS, Henderson WR, Foster GE, Romer LM, Koehle MS & Sheel AW (2017). Exercise-induced quadriceps muscle fatigue in men and women: effects of arterial oxygen content and respiratory muscle work. *J Physiol* **595**, 5227– 5244.
- Dominelli PB, Render JN, Molgat-Seon Y, Foster GE, Romer LM & Sheel AW (2015). Oxygen cost of exercise hyperpnoea is greater in women compared with men. *J Physiol* **593**, 1965– 1979.

- Farina D, Merletti R & Enoka RM (2004). The extraction of neural strategies from the surface EMG. *J Appl Physiol* **96**, 1486– 1495.
- Finn HT, Rouffet DM, Kennedy DS, Green S & Taylor JL (2018). Motoneuron excitability of the quadriceps decreases during a fatiguing submaximal isometric contraction. *J Appl Physiol* **124**, 970– 979.
- Froberg K & Pedersen PK (1984). Sex differences in endurance capacity and metabolic response to prolonged, heavy exercise. *Eur J Appl Physiol Occup Physiol* **52**, 446– 450.
- Glance BW, Kremenic IJ & McHugh MP (2013). Sex differences in central and peripheral mechanisms of fatigue in cyclists. *Eur J Appl Physiol* **113**, 1091– 1098.
- Goodall S, Ross EZ & Romer LM (2010). Effect of graded hypoxia on supraspinal contributions to fatigue with unilateral knee-extensor contractions. *J Appl Physiol* **109**, 1842– 1851.
- Goulding RP, Roche DM & Marwood S (2017). Prior exercise speeds pulmonary oxygen uptake kinetics and increases critical power during supine but not upright cycling. *Exp Physiol* **102**, 1158– 1176.
- Grassi B, Pogliaghi S, Rampichini S, Quaresima V, Ferrari M, Marconi C & Cerretelli P (2003). Muscle oxygenation and pulmonary gas exchange kinetics during cycling exercise on-transitions in humans. *J Appl Physiol* **95**, 149– 158.
- Gruet M, Temesi J, Rupp T, Levy P, Verges S & Millet GY (2014). Dynamics of corticospinal changes during and after high-intensity quadriceps exercise. *Exp Physiol* **99**, 1053– 1064.
- Guenette JA, Romer LM, Querido JS, Chua R, Eves ND, Road JD, McKenzie DC & Sheel AW (2010). Sex differences in exercise-induced diaphragmatic fatigue in endurance-trained athletes. *J Appl Physiol* **109**, 35– 46.
- Guenette JA, Witt JD, McKenzie DC, Road JD & Sheel AW (2007). Respiratory mechanics during exercise in endurance-trained men and women. *J Physiol* **581**, 1309– 1322.
- Harmer AR, Ruell PA, Hunter SK, McKenna MJ, Thom JM, Chisholm DJ & Flack JR (2014). Effects of type 1 diabetes, sprint training and sex on skeletal muscle sarcoplasmic reticulum Ca²⁺ uptake and Ca²⁺-ATPase activity. *J Physiol* **592**, 523– 535.
- Harms CA, McClaran SR, Nিকেle GA, Pegelow DF, Nelson WB & Dempsey JA (1998). Exercise-induced arterial hypoxaemia in healthy young women. *J Physiol* **507**, 619– 628.
- Heckman CJ, Johnson M, Mottram C & Schuster J (2008). Persistent inward currents in spinal motoneurons and their influence on human motoneuron firing patterns. *Neuroscience* **14**, 264– 275.
- Hermens HJ, Freriks B, Disselhorst-Klug C & Rau G (2000). Development of recommendations for SEMG sensors and sensor placement procedures. *J Electromyogr Kinesiol* **10**, 361– 374.
- Hunter SK, Butler JE, Todd G, Gandevia SC & Taylor JL (2006). Supraspinal fatigue does not explain the sex difference in muscle fatigue of maximal contractions. *J Appl Physiol* **101**, 1036– 1044.
- Hureau TJ, Romer LM & Amann M (2018). The ‘sensory tolerance limit’: A hypothetical construct determining exercise performance? *Eur J Sport Sci* **18**, 13– 24.
- Jones AM, Vanhatalo A, Burnley M, Morton RH & Poole DC (2010). Critical power: Implications for determination of \dot{V}_{O_2max} and exercise tolerance. *Med Sci Sports Exerc* **42**, 1876– 1890.
- Jones AM, Wilkerson DP, DiMenna F, Fulford J & Poole DC (2008). Muscle metabolic responses to exercise above and below the “critical power” assessed using 31P-MRS. *Am J Physiol Integr Comp Physiol* **294**, R585– R593.
- Joyner MJ (2017). Physiological limits to endurance exercise performance: influence of sex. *J Physiol* **595**, 2949– 2954.
- Keane KM, Bailey SJ, Vanhatalo A, Jones AM & Howatson G (2018). Effects of montmorency tart cherry (*L. Prunus cerasus*) consumption on nitric oxide biomarkers and exercise performance. *Scand J Med Sci Sport* **28**, 1746– 1756.
- Kennedy DS, McNeil CJ, Gandevia SC & Taylor JL (2016). Effects of fatigue on corticospinal excitability of the human knee extensors. *Exp Physiol* **101**, 1552– 1564.

- Kordi M, Menzies C & Parker Simpson L (2018). Relationship between power–duration parameters and mechanical and anthropometric properties of the thigh in elite cyclists. *Eur J Appl Physiol* **118**, 637– 645.
- Kuhn A, Keller T, Lawrence M & Morari M (2010). The influence of electrode size on selectivity and comfort in transcutaneous electrical stimulation of the forearm. *IEEE Trans Neural Syst Rehabil Eng* **18**, 255– 262.
- Lanza MB, Balshaw TG, Massey GJ & Folland JP (2018). Does normalization of voluntary EMG amplitude to M_{MAX} account for the influence of electrode location and adiposity? *Scand J Med Sci Sport* **28**, 2558– 2566.
- Martin TR, Castile RG, Fredberg JJ, Wohl MEB & Mead J (1987). Airway size is related to sex but not lung size in normal adults. *J Appl Physiol* **63**, 2042– 2047.
- Mead J (1980). Dysanapsia in normal lungs assessed by the relationship between maximal flow, static recoil, and vital capacity. *Am Rev Respir Dis* **121**, 339– 342.
- Merton PA (1954). Voluntary strength and fatigue. *J Physiol* **123**, 553– 564.
- Messere A & Roatta S (2013). Influence of cutaneous and muscular circulation on spatially resolved versus standard Beer-Lambert near-infrared spectroscopy. *Physiol Rep* **1**, e00179.
- Mitchell EA, Martin NRW, Bailey SJ & Ferguson RA (2018a). Critical power is positively related to skeletal muscle capillarization and type I muscle fibers in endurance-trained individuals. *J Appl Physiol* **125**, 737– 745.
- Mitchell RA, Schaeffer MR, Ramsok AH, Wilkie SS & Guenette JA (2018b). Sex differences in respiratory muscle activation patterns during high-intensity exercise in healthy humans. *Respir Physiol Neurobiol* **247**, 57– 60.
- Miura A, Endo M, Sato H, Sato H, Barstow TJ & Fukuba Y (2002). Relationship between the curvature constant parameter of the power–duration curve and muscle cross-sectional area of the thigh for cycle ergometry in humans. *Eur J Appl Physiol* **87**, 238– 244.
- Murgatroyd SR, Ferguson C, Ward SA, Whipp BJ & Rossiter HB (2011). Pulmonary O_2 uptake kinetics as a determinant of high-intensity exercise tolerance in humans. *J Appl Physiol* **110**, 1598– 1606.
- Murphy WG (2014). The sex difference in haemoglobin levels in adults - Mechanisms, causes, and consequences. *Blood Rev* **28**, 41– 47.
- Ørtenblad N, Westerblad H & Nielsen J (2013). Muscle glycogen stores and fatigue. *J Physiol* **591**, 4405– 4413.
- Parker BA, Smithmyer SL, Pelberg JA, Mishkin AD, Herr MD & Proctor DN (2007). Sex differences in leg vasodilation during graded knee extensor exercise in young adults. *J Appl Physiol* **103**, 1583– 1591.
- Pate RR & O'Neill JR (2007). American women in the marathon. *Sports Med* **37**, 294– 298.
- De Pauw K, Roelands B, Cheung SS, De Geus B, Rietjens G & Meeusen R (2013). Guidelines to classify subject groups in sport-science research. *Int J Sports Physiol Perform* **8**, 111– 122.
- Place N & Millet GY (2020). Quantification of neuromuscular fatigue: what do we do wrong and why? *Sport Med* **50**, 439– 447.
- Poole DC, Burnley M, Vanhatalo A, Rossiter HB & Jones AM (2016). Critical power: an important fatigue threshold in exercise physiology. *Med Sci Sports Exerc* **48**, 2320– 2334.
- Poole DC & Jones AM (2012). Oxygen uptake kinetics. *Compr Physiol* **2**, 933– 996.
- Poole DC, Ward SA, Gardner GW & Whipp BJ (1988). Metabolic and respiratory profile of the upper limit for prolonged exercise in man. *Ergonomics* **31**, 1265– 1279.
- Roepstorff C, Steffensen CH, Madsen M, Stallknecht B, Kanstrup I-L, Richter EA & Kiens B (2002). Gender differences in substrate utilization during submaximal exercise in endurance-trained subjects. *Am J Physiol Metab* **282**, E435– E447.
- Roepstorff C, Thiele M, Hillig T, Pilegaard H, Richter EA, Wojtaszewski JFP & Kiens B (2006). Higher skeletal muscle α 2AMPK activation and lower energy charge and fat oxidation in men than in women during submaximal exercise. *J Physiol* **574**, 125– 138.
- Russ DW, Lanza IR, Rothman D & Kent-Braun JA (2005). Sex differences in glycolysis during brief, intense isometric contractions. *Muscle and Nerve* **32**, 647– 655.

- Russ DW, Towse TF, Wigmore DM, Lanza IR & Kent-Braun JA (2008). Contrasting influences of age and sex on muscle fatigue. *Med Sci Sports Exerc* **40**, 234– 241.
- Ryan TE, Erickson ML, Brizendine JT, Young HJ & McCully KK (2012). Noninvasive evaluation of skeletal muscle mitochondrial capacity with near-infrared spectroscopy: Correcting for blood volume changes. *J Appl Physiol* **113**, 175– 183.
- Schaeffer MR, Mendonca CT, Levangie MC, Andersen RE, Taivassalo T & Jensen D (2014). Physiological mechanisms of sex differences in exertional dyspnoea: Role of neural respiratory motor drive. *Exp Physiol* **99**, 427– 441.
- Schäfer LU, Hayes M & Deckerle J (2019). The magnitude of neuromuscular fatigue is not intensity dependent when cycling above critical power but relates to aerobic and anaerobic capacities. *Exp Physiol* **104**, 209– 219.
- Schwartz J, Katz SA, Fegley RW & Tockman MS (1988). Sex and race differences in the development of lung function. *Am Rev Respir Dis* **138**, 1415– 1421.
- Sheel AW, Guenette JA, Yuan R, Holy L, Mayo JR, McWilliams AM, Lam S & Coxson HO (2009). Evidence for dysanapsis using computed tomographic imaging of the airways in older ex-smokers. *J Appl Physiol* **107**, 1622– 1628.
- Sidhu SK, Cresswell AG & Carroll TJ (2013). Corticospinal responses to sustained locomotor exercises: moving beyond single-joint studies of central fatigue. *Sports Med* **43**, 437– 449.
- Sidhu SK, Weavil JC, Mangum TS, Jessop JE, Richardson RS, Morgan DE & Amann M (2017). Group III/IV locomotor muscle afferents alter motor cortical and corticospinal excitability and promote central fatigue during cycling exercise. *Clin Neurophysiol* **128**, 44– 55.
- Simoneau JA & Bouchard C (1989). Human variation in skeletal muscle fiber-type proportion and enzyme activities. *Am J Physiol Endocrinol Metab* **257**, E567– E572.
- Škarabot J, Ansdell P, Brownstein CG, Thomas K, Goodall S, Howatson G & Durbaba R (2019a). Electrical stimulation of human corticospinal axons at the level of the lumbar spinal segments. *Eur J Neurosci* **49**, 1254– 1267.
- Škarabot J, Mesquita RNO, Brownstein CG & Ansdell P (2019b). Myths and methodologies: How loud is the story told by the transcranial magnetic stimulation-evoked silent period? *Exp Physiol* **104**, 635– 642.
- Sparling PB (1980). A meta-analysis of studies comparing maximal oxygen uptake in men and women. *Res Q Exerc Sport* **51**, 542– 552.
- Staron RS, Hagerman FC, Hikida RS, Murray TF, Hostler DP, Crill MT, Ragg KE & Toma K (2000). Fiber type composition of the vastus lateralis muscle of young men and women. *J Histochem Cytochem* **48**, 623– 629.
- Sundberg CW, Hunter SK & Bundle MW (2016). Rates of performance loss and neuromuscular activity in men and women during cycling: evidence for a common metabolic basis of muscle fatigue. *J Appl Physiol* **122**, 130– 141.
- Tamm AS, Lagerquist O, Ley AL & Collins DF (2009). Chronotype influences diurnal variations in the excitability of the human motor cortex and the ability to generate torque during a maximum voluntary contraction. *J Biol Rhythms* **24**, 211– 224.
- Tarnopolsky LJ, MacDougall JD, Atkinson SA, Tarnopolsky MA & Sutton JR (1990). Gender differences in substrate for endurance exercise. *J Appl Physiol* **68**, 302– 308.
- Temesi J, Arnal PJ, Rupp T, Féasson L, Cartier R, Gergelé L, Verges S, Martin V & Millet GY (2015). Are females more resistant to extreme neuromuscular fatigue? *Med Sci Sports Exerc* **47**, 1372– 1382.
- Thomas K, Elmeua M, Howatson G & Goodall S (2016). Intensity-dependent contribution of neuromuscular fatigue after constant-load cycling. *Med Sci Sports Exerc* **48**, 1751– 1760.
- Thomas K, Goodall S & Howatson G (2018). Performance fatigability is not regulated to a peripheral critical threshold. *Exerc Sport Sci Rev* **46**, 240– 246.

- Todd G, Taylor JL & Gandevia SC (2003). Measurement of voluntary activation of fresh and fatigued human muscles using transcranial magnetic stimulation. *J Physiol* **551**, 661– 671.
- Ugawa Y, Genba-Shimizu K & Kanazawa I (1995). Electrical stimulation of the human descending motor tracts at several levels. *Can J Neurol Sci* **22**, 36– 42.
- Vanhatalo A, Black MI, DiMenna FJ, Blackwell JR, Schmidt JF, Thompson C, Wylie LJ, Mohr M, Bangsbo J, Krstrup P & Jones AM (2016). The mechanistic bases of the power–time relationship: muscle metabolic responses and relationships to muscle fibre type. *J Physiol* **594**, 4407– 4423.
- Vanhatalo A, Fulford J, Dimenna FJ & Jones AM (2010). Influence of hyperoxia on muscle metabolic responses and the power–duration relationship during severe-intensity exercise in humans: A 31P magnetic resonance spectroscopy study. *Exp Physiol* **95**, 528– 540.
- Vigotsky AD, Halperin I, Lehman GJ, Trajano GS & Vieira TM (2018). Interpreting signal amplitudes in surface electromyography studies in sport and rehabilitation sciences. *Front Physiol* **8**, 985.
- Weavil JC & Amann M (2018). Corticospinal excitability during fatiguing whole body exercise. *Prog Brain Res* **240**, 219– 240.
- Weavil JC, Sidhu SK, Mangum TS, Richardson RS & Amann M (2016). Fatigue diminishes motoneuronal excitability during cycling exercise. *J Neurophysiol* **116**, 1743– 1751.
- Welch JF, Archiza B, Guenette JA, West CR & Sheel AW (2018). Sex differences in diaphragmatic fatigue: the cardiovascular response to inspiratory resistance. *J Physiol* **596**, 4017– 4032.
- Wen AS, Woo MS & Keens TG (1997). How many maneuvers are required to measure maximal inspiratory pressure accurately? *Chest* **111**, 802– 807.
- Westerbacka J, Cornér A, Tiikkainen M, Tamminen M, Vehkavaara S, Häkkinen AM, Fredriksson J & Yki-Järvinen H (2004). Women and men have similar amounts of liver and intra-abdominal fat, despite more subcutaneous fat in women: Implications for sex differences in markers of cardiovascular risk. *Diabetologia* **47**, 1360– 1369.
- Willcocks RJ, Williams CA, Barker AR, Fulford J & Armstrong N (2010). Age- and sex-related differences in muscle phosphocreatine and oxygenation kinetics during high-intensity exercise in adolescents and adults. *NMR Biomed* **23**, 569– 577.
- Witt JD, Guenette JA, Rupert JL, McKenzie DC & Sheel AW (2007). Inspiratory muscle training attenuates the human respiratory muscle metaboreflex. *J Physiol* **584**, 1019– 1028.
- Yoon B-K, Kravitz L & Robergs RA (2007a). Does protocol duration affect VO_{2max} ? *Med Sci Sport Exerc* **39**, 1186– 1192.
- Yoon T, Delap BS, Griffith EE & Hunter SK (2007b). Mechanisms of fatigue differ after low- and high-force fatiguing contractions in men and women. *Muscle and Nerve* **36**, 515– 524.