

Marquette University

e-Publications@Marquette

Biomedical Sciences Faculty Research and
Publications

Biomedical Sciences, Department of

9-2008

Evidence that α 2-Antiplasmin Becomes Covalently Ligated to Plasma Fibrinogen in the Circulation: a New Role for Plasma Factor XIII in Fibrinolysis Regulation

Michael W. Mosesson
Blood Center of Wisconsin

Kevin R. Siebenlist
Marquette University, kevin.siebenlist@marquette.edu

Irene Hernandez
Blood Research Institute

K. N. Lee
University of Oklahoma Health Sciences Center

V. J. Christiansen
University of Oklahoma Health Sciences Center

See next page for additional authors

Follow this and additional works at: https://epublications.marquette.edu/biomedsci_fac



Part of the [Neurosciences Commons](#)

Recommended Citation

Mosesson, Michael W.; Siebenlist, Kevin R.; Hernandez, Irene; Lee, K. N.; Christiansen, V. J.; and McKee, P. A., "Evidence that α 2-Antiplasmin Becomes Covalently Ligated to Plasma Fibrinogen in the Circulation: a New Role for Plasma Factor XIII in Fibrinolysis Regulation" (2008). *Biomedical Sciences Faculty Research and Publications*. 217.

https://epublications.marquette.edu/biomedsci_fac/217

Authors

Michael W. Mosesson, Kevin R. Siebenlist, Irene Hernandez, K. N. Lee, V. J. Christiansen, and P. A. McKee

Marquette University

e-Publications@Marquette

Biomedical Sciences Faculty Research and Publications/College of Health Sciences

This paper is NOT THE PUBLISHED VERSION.

Access the published version via the link in the citation below.

Journal of Thrombosis and Haemostasis, Vol. 6, No. 9 (September 2008): 1565-1570. [DOI](#). This article is © Wiley and permission has been granted for this version to appear in [e-Publications@Marquette](#). Wiley does not grant permission for this article to be further copied/distributed or hosted elsewhere without express permission from Wiley.

Evidence that α 2-Antiplasmin Becomes Covalently Ligated to Plasma Fibrinogen in the Circulation: A New Role for Plasma Factor XIII in Fibrinolysis Regulation

M. W. Mosesson

Blood Research Institute, BloodCenter of Wisconsin, Milwaukee, WI

K. R. Siebenlist

Department of Biomedical Sciences, Marquette University, Milwaukee, WI

I. Hernandez

Blood Research Institute, BloodCenter of Wisconsin, Milwaukee, WI

K. N. Lee

Department of Medicine, University of Oklahoma Health Sciences Center, Oklahoma, OK

V. J. Christiansen

Department of Medicine, University of Oklahoma Health Sciences Center, Oklahoma, OK

P. A. Mckee

Department of Medicine, University of Oklahoma Health Sciences Center, Oklahoma, OK

Abstract

Summary. Background: Plasma alpha₂-antiplasmin (α₂AP) is a rapid and effective inhibitor of the fibrinolytic enzyme plasmin. Congenital α₂AP deficiency results in a severe hemorrhagic disorder due to accelerated fibrinolysis. It is well established that in the presence of thrombin-activated factor XIII (FXIIIa), α₂AP becomes covalently ligated to the distal α chains of fibrin or fibrinogen at lysine 303 (two potential sites per molecule). Some time ago we showed that α₂AP is covalently linked to plasma fibrinogen. That singular observation led to our hypothesis that native plasma factor XIII (FXIII), which is known to catalyze covalent cross-linking of fibrinogen in the presence of calcium ions, can also incorporate α₂AP into fibrinogen in the circulation. **Results and Conclusions:** We now provide evidence that FXIII incorporates I¹²⁵-labelled α₂AP into the Aα-chain sites on fibrinogen or fibrin. We also measured the content of α₂AP in isolated plasma fibrinogen fractions by ELISA and found that substantial amounts were present (1.2–1.8 moles per mole fibrinogen). We propose that α₂AP becomes ligated to fibrinogen while in the circulation through the action of FXIII, and that its immediate presence in plasma fibrinogen contributes to regulation of *in vivo* fibrinolysis.

Introduction

Plasma alpha₂-antiplasmin (α₂AP) is a rapid and effective serine protease inhibitor (serpin) of the fibrinolytic enzyme plasmin¹⁻⁴. When this inhibitor is absent from plasma, as occurs in congenital homozygous α₂AP-deficiency, a severe hemorrhagic disorder results that is characterized by increased susceptibility of intravascular thrombi to fibrinolysis⁵⁻⁷. The fibrinolytic defect is reversible by the addition of α₂AP. It is well known that in the presence of thrombin-activated factor XIIIa (FXIIIa) the inhibitor becomes covalently ligated ('cross-linked') to a single site, lysine 303, on the fibrin- or fibrinogen Aα-chain^{8,9}. Unbound α₂AP inhibits tPA-induced fibrinogenolysis; however, only after it has been incorporated into fibrin via FXIIIa, is α₂AP an effective inhibitor of fibrinolysis¹⁰⁻¹³. In addition, plasmin that becomes bound to fibrin is protected from inhibition by unbound α₂AP¹⁴, thus emphasizing the importance of prior α₂AP incorporation into fibrin(ogen) for effectively mediating inhibition of fibrinolysis. Finally, α₂AP is incorporated into fibrin at only 20% to 30% of the potentially available lysine 303 sites¹⁵⁻¹⁷, suggesting that the unreactive sites may already be occupied or otherwise unavailable for ligation.

There are two forms of α₂AP in human plasma, a full length 464-residue protein with amino-terminal methionine (Met-α₂AP) that accounts for ~30% of the total, the remainder being composed of a shortened 452-residue form with amino-terminal asparagine (Asn-α₂AP)^{18,19}. The proportions of the two forms of α₂AP in plasma are related to the R/W6 single nucleotide polymorphism, with R being associated with higher Asn-α₂AP levels²⁰. Incorporation of Asn-α₂AP into fibrin is 3–13 times greater than with the full length precursor form, Met-α₂AP^{21,22}, but otherwise both forms display the same inhibitory activities.

A plasmin inhibitory activity like that of plasma α₂AP was discovered in fibrinogen many years ago²³, although it was not identified as α₂AP until much later when immunochemical analyses showed that α₂AP was a constituent of normal fibrinogen as well as a dysfibrinogenemic fibrinogen (fibrinogen Cedar Rapids)²⁴. These observations combined with prior knowledge that native non-thrombin-activated plasma factor XIII (FXIII) can efficiently introduce covalent cross-links into fibrinogen as well

as fibrin in the presence of calcium ions²⁵, and the retrospective insight that α_2 AP is incompletely incorporated into fibrin by FXIIIa¹⁵⁻¹⁷, suggested to us that the α_2 AP found in plasma fibrinogen might have been incorporated in the circulation through the action of FXIII. In this present study we show that substantial amounts of α_2 AP are present in circulating fibrinogen, and we present evidence that this process is mediated by plasma FXIII. We also introduce the concept that ligation of α_2 AP to plasma fibrinogen prior to initiation of clotting serves an important role in down-regulating the rate of fibrinolysis.

Materials and methods

Chemicals and reagents were of the highest purity available. Trasylol (aprotinin) was obtained from Miles Inc. (Kankakee, IL, USA), and DE-52 cellulose was from Whatmann Inc. (Clifton, NJ, USA). Human α -thrombin (3188 μ mg⁻¹) was obtained from Enzyme Research Laboratories (South Bend, IN, USA), and fibrinogen was isolated from human citrated plasma pools (obtained through the BloodCenter of Wisconsin) by glycine precipitation followed by sub-fractionation to 'fraction I-2' as described²⁶. Most A α -chains in fraction I-2 fibrinogen are full length and therefore they contain the α_2 AP ligation site at A α 303. This plasma fraction was further separated into fibrinogen 1 (γ_A, γ_A) and fibrinogen 2 (γ_A, γ') by ion exchange chromatography²⁷. Des- α C fibrinogen (old terminology, 'fraction I-9D') was produced by limited plasmin digestion of fibrinogen²⁸, lacked ~390 residues of the C-terminal region of the A α chain^{29, 30}, and therefore lacked the sequence containing A α 303. Fibrinogen fraction I-2 from the fibrinogen Cedar Rapids proposita, which previously had been demonstrated to contain α_2 AP²⁴, was also analyzed. FXIII was prepared from pooled plasma³¹ and assayed as FXIIIa on a FXIII-free fibrin substrate (fibrin 1) in the presence of 10 mm CaCl₂ as described^{25, 32}. FXIII concentrations were determined spectrophotometrically at 280 nm, using an absorbance coefficient ($A_{1\text{cm}}^{1\%}$, 280 nm) of 13.8³³. The specific activity was 2100 to 2300 Loewy u mg⁻¹. Normal plasma levels of factor XIII are 80–110 Loewy u mL⁻¹. Recombinant Asn- α_2 AP comprised of 464 AA, was prepared as described³⁴, and had a mass of 50 583 Da. A portion of this material was labeled with ¹²⁵I by a lactoperoxidase method³⁵, and stored at a stock concentration of 0.87 mg mL⁻¹ (17 μ m).

Incorporation of α_2 AP into fibrinogen or fibrin by FXIII

For investigating FXIII-mediated incorporation of α_2 AP, FXIII-free fibrinogen 1 (8.8 μ m, final) in 50 mm Tris, 100 mm NaCl, 5.0 mm CaCl₂, 0.4 mm DTT, pH 7.4, was mixed with ¹²⁵I-labeled α_2 AP (1.0 to 6.0 μ m), and ligation initiated at room temperature by adding 100 Loewy u mL⁻¹ FXIII (final). After 6 h incubation, the reaction was terminated by adding an equal volume of 2-fold concentrated Laemmli buffer containing 1% β -mercaptoethanol, and the products of the reaction were analyzed by SDS-PAGE on 9% polyacrylamide gels employing a discontinuous buffer system³⁶. Dried Coomassie Brilliant Blue-stained gels were subjected to autoradiography using Kodak X-Omat film. Autoradiograms were digitized on a CanoScan 9950F flat bed scanner (Canon USA Inc, Lake Success, NY, USA), and α_2 AP-containing bands quantified using Image J (<http://rsb.info.nih.gov/ij/>). Results from stained gels were normalized against the B β region of the gel whereas results from autoradiograms were normalized against known quantities of ¹²⁵I-labeled α_2 AP run on identical gels. Control samples included ¹²⁵I-labeled α_2 AP, fibrinogen 1, and fibrinogen 1 plus FXIII without ¹²⁵I-labeled α_2 AP. To determine the amount of ¹²⁵I-labeled α_2 AP that could be incorporated into cross-linked fibrin by FXIIIa, identical samples were prepared and the reaction initiated by adding 100 Loewy u mL⁻¹ FXIII and 0.5 u mL⁻¹

thrombin. For determining incorporation rates, FXIII-free fibrinogen 1 (8.8 μm final) in 50 mM Tris, 100 mM NaCl, 5.0 mM CaCl_2 , 0.4 mM DTT, pH 7.4, was mixed with ^{125}I -labeled $\alpha_2\text{AP}$ (1.0 or 5.0 μm), and the reaction initiated by adding 100 Loewy u mL^{-1} FXIII or FXIIIa (containing 0.5 u mL^{-1} thrombin). At selected intervals of up to 18 h the incubation was terminated by adding an equal volume of 2X Laemmli buffer containing 1% β -mercaptoethanol, and the products of the reaction were analyzed as described above.

Immunoassay of fibrinogen and $\alpha_2\text{AP}$

Fibrinogen concentrations were determined by ELISA³⁷ using biotinylated rabbit anti-human fibrinogen IgG (Dako, Carpinteria, CA, USA) for tagging the fibrinogen that had been bound to wells. Antigen loading, equilibration and processing conditions were the same as described below for $\alpha_2\text{AP}$ detection. Fibrinogen fraction I-2 (>97% coagulable) was used for constructing a standard curve at test concentrations ranging from 20 to 0.0125 $\mu\text{g mL}^{-1}$. In order to measure $\alpha_2\text{AP}$ in fibrinogen-containing samples, we applied fibrinogen at concentrations that were five to eight times higher (2.5–20 $\mu\text{g mL}^{-1}$, 0.74–5.9 nM) than those usually used for obtaining a linear plot³⁷. Thus, the amount of fibrinogen bound to the plate was determined from a non-linear portion of the fibrinogen calibration curve.

To measure $\alpha_2\text{AP}$ in fibrinogen-containing samples by ELISA, we first constructed an $\alpha_2\text{AP}$ calibration curve as follows: Asn- $\alpha_2\text{AP}$ (0.125–1.5 $\mu\text{g mL}^{-1}$) was added to immunoplates (MG Scientific, Pleasant Prairie, WI, USA), incubated overnight at 4°C, washed with PBS, blocked with 2% non-fat dried milk in PBS for 1 h at room temperature, and rinsed with 200 μL PBS-Tween 20 (0.05%). Washed plates were then treated with goat anti-human $\alpha_2\text{AP}$ (Nordic Immunology, Tilberg, The Netherlands) at 1:5000 dilution in PBS-Tween 20, incubated for 1 h at 37°C, and washed four times with PBS-Tween (0.05%). Horseradish peroxidase-labeled rabbit anti-goat IgG (Zymed, South San Francisco, CA, USA) at 1:15 000 dilution was added, incubated for 1 h at 37°C, and washed with PBS-Tween. After the final rinse, 100 μL o-phenylenediamine (OPD) solution (Zymed, South San Francisco, CA, USA) in 0.22 M citric acid, 0.05 M sodium phosphate, pH 5.0 buffer, was added and incubated in subdued light at room temperature for 10–15 min. The reaction was terminated with 50 μL 2 N H_2SO_4 and the plate read at 490 nm on a Versamax Plate Reader. For determining the $\alpha_2\text{AP}$ content in fibrinogen-containing specimens, replicate wells were processed exactly as described above for $\alpha_2\text{AP}$ calibration. The amount of $\alpha_2\text{AP}$ in fibrinogen was expressed as a molar ratio.

Results

Incorporation of $\alpha_2\text{AP}$ radioactivity into fibrinogen or fibrin

Earlier studies had demonstrated that native FXIII (FXIII) was able to cross-link fibrinogen in the presence of calcium ions²⁵, and these findings prompted us to investigate whether FXIII might also incorporate $\alpha_2\text{AP}$ into fibrinogen without requiring thrombin activation. Incubating fibrinogen at a physiological concentration (8.8 μm) with ^{125}I -labeled $\alpha_2\text{AP}$ (1–6 μm) plus physiological concentrations of FXIII (100 Loewy units mL^{-1}), as assessed from autoradiograms of SDS-PAGE gels, showed new radiolabeled bands (Fig. 1, panel A). These corresponded to an $\alpha_2\text{AP}/\text{A}\alpha$ -chain heterodimer (' $\alpha_2\text{AP}/\text{A}\alpha$ chain'), an $\alpha_2\text{AP}/\text{A}\alpha$ - $\text{A}\alpha$ chain heterotrimer (' $\alpha_2\text{AP}/\text{A}\alpha$ dimer') and an $\alpha_2\text{AP}/\text{A}\alpha$ -chain heteropolymer (' $\alpha_2\text{AP}/\text{A}\alpha$ polymer'). In the Coomassie stained gel, we found new bands corresponding to the radiolabeled $\alpha_2\text{AP}/\text{A}\alpha$ heterodimer and $\alpha_2\text{AP}/\text{A}\alpha$ -chain heteropolymer positions, plus non-radiolabeled

γ -dimers (panel B). The α_2 AP/ α - α chain heterotrimer position was faintly stained but readily detected in the autoradiogram.




Figure 1 Incorporation of 125 I-labeled α_2 AP into fibrinogen in the presence of FXIII (panels A and B) or, after adding thrombin to form FXIIIa and fibrin (panels C and D). Autoradiograms are shown in panels A and C and corresponding Coomassie blue stained gels in B and D).

Adding thrombin to the reaction mixtures to convert fibrinogen to fibrin and to activate FXIII to FXIIIa (Fig. 1, panels C and D) resulted in similarly higher molecular weight stained bands, but the vast majority of radioactive α_2 AP had been incorporated into the α -polymer region rather than the α_2 AP/ α heterodimer or heterotrimer positions (panel C). Figure 2 shows densitometric scans of the Fig. 1 panels A and C gels. It is evident that the radioactivity incorporated into fibrinogen or fibrin increased proportionately with the concentration of α_2 AP, and that incorporation into fibrin was slightly higher than for fibrinogen at any given α_2 AP concentration. At the highest α_2 AP concentration that we studied, only about 40% of the potential α_2 AP sites had been labeled.




Figure 2 Molar ratio of α_2 AP incorporation into fibrinogen (FXIII) or fibrin (FXIIIa) as a function of the concentration of α_2 AP (μM). Data points are derived from densitometric scans of the gels in Fig. 1, panels A and C.

We also studied time-dependent incorporation of radioactivity into fibrinogen at α_2 AP concentrations of 1 and 5 μM (Fig. 3) as well as into fibrin. At the lower α_2 AP concentration, radioactively-labeled α_2 AP/ α chain polymers were detected in fibrinogen within 5 min (Fig. 3, panels A and B), and to a

similar extent in fibrin (gels not shown). Inspection and densitometric scans of these gels revealed that incorporation into either fibrinogen or fibrin began to plateau at about 30 min. At $5 \mu\text{M}$ $\alpha_2\text{AP}$, incorporation of radioactivity began to plateau at 15 min (Fig. 3, panels C and D) for fibrinogen as well as fibrin (gels not shown). Densitometric scans of the fibrinogen or fibrin SDS-PAGE gels (Fig. 4) indicated that maximal incorporation of radioactivity was about 0.25 moles per mole for fibrinogen, and slightly greater for fibrin (~ 0.30 moles per mole).




Figure 3 Incorporation of $\alpha_2\text{AP}$ into fibrinogen at 1 and $5 \mu\text{M}$ $\alpha_2\text{AP}$, as a function of time in minutes (m) or hours (h). Stained gels, left. Autoradiograms, right.




Figure 4 Molar ratio of $\alpha_2\text{AP}$ incorporation into fibrinogen by FXIII or into fibrin by FXIIIa as a function of time (min). The reaction is presented at an $\alpha_2\text{AP}$ concentration of $5 \mu\text{M}$. The data points for fibrinogen are based upon the experiment shown in Fig. 3, whereas the gel for fibrin is not shown.

Fibrinogen $\alpha_2\text{AP}$ content

The $\alpha_2\text{AP}$ content in plasma fibrinogen fraction I-2 from four separate pools of normal citrated plasma and that from the proposita of the Cedar Rapids dysfibrinogenemia ($\gamma 275\text{R}$ to C)²⁴ was assessed by immunoassay and is represented as the molar ratio of $\alpha_2\text{AP}$ to fibrinogen (Table 1). Given the fact that there is a single site at A α 303 on each A α chain to which $\alpha_2\text{AP}$ can become ligated^{8, 15-17, 38}, the maximum ratio of $\alpha_2\text{AP}$ to fibrinogen is two. With the sole exception of des- αC fibrinogen, the negative control, which lacks C-terminal portions of A α chains containing the $\alpha_2\text{AP}$ binding site, each of the fibrinogen fractions tested contained $\alpha_2\text{AP}$. In every case, the $\alpha_2\text{AP}$:fibrinogen ratio was between one and two, indicating that considerable amounts of $\alpha_2\text{AP}$ had been incorporated into plasma fibrinogen, a result that is consistent with earlier observations of the presence of $\alpha_2\text{AP}$ in plasma fibrinogen^{23, 24}. As discussed in the next section, the $\alpha_2\text{AP}$ content of Cedar Rapids fibrinogen was the same or possibly

even lower than that in normal fibrinogen, and thus did not account for delayed fibrinolysis that has been observed in this fibrinogen²⁴.

Table 1. α_2 AP/fibrinogen molar ratio

Fibrinogen fraction	<i>n</i>	Ratio	Range
Normal fraction I-2	4	1.5 (\bar{X})	1.25–1.8
Cedar Rapids fraction I-2	1	1.2	–
Des- α C fibrinogen	1	0	0

Discussion

A plasmin inhibitory activity corresponding functionally to α_2 AP was discovered in plasma fibrinogen many years ago²³, although it was not specifically identified as α_2 AP until years later, when it was shown by immunochemical analysis to be a constituent of normal fibrinogen as well as a dysfibrinogenemic fibrinogen, fibrinogen Cedar Rapids²⁴. This discovery suggested that α_2 AP might have been incorporated covalently into fibrinogen through the action of a calcium ion-dependent transglutaminase like FXIII, which circulates with fibrinogen in plasma. This postulation, combined with prior knowledge that ‘native’ FXIII is an active enzyme that efficiently introduces covalent cross-links into fibrinogen or fibrin molecules²⁵, prompted us to investigate this possibility in detail. Our present experiments confirm that α_2 AP is a covalently bound constituent of plasma fibrinogen, that it is present in substantial amounts, and that plasma FXIII can mediate its incorporation into fibrinogen. These findings support the hypothesis that α_2 AP becomes ligated to plasma fibrinogen in the circulation, and is very likely to have been incorporated through the action of circulating FXIII.

We were particularly interested in quantifying the α_2 AP content in the Cedar Rapids dysfibrinogen²⁴, not only because members of this kindred had experienced severe thrombophilia associated with delayed fibrinolysis, but also because Cedar Rapids fibrinogen had been the index case for measuring the presence of α_2 AP in plasma fibrinogen; we were still seeking to explain that phenomenon in terms of the α_2 AP content. Our present results confirm those in the first study, and indicate that the α_2 AP content of Cedar Rapids fibrinogen is the same or possibly even lower than that in normal fibrinogen, and thus cannot alone account for delayed fibrinolysis²⁴. Another explanation will be required.

Although it is well established that in the presence of thrombin-activated FXIIIa, α_2 AP becomes ligated to the fibrin or fibrinogen A α -chain^{8,9,15-17,38}, it was not known prior to this study that native FXIII, in addition to introducing covalent cross-links into fibrinogen or fibrin in the presence of calcium ions²⁵, can also incorporate α_2 AP into fibrinogen. The rate of α_2 AP incorporation into fibrinogen by FXIII was nearly as rapid as the rate of incorporation into fibrin by FXIIIa. These findings provide an attractive mechanistic explanation for why circulating fibrinogen contains α_2 AP. Furthermore, as the content of α_2 AP in fibrinogen is so relatively high (i.e. 1.2–1.8 moles per mole fibrinogen (Table 1) it readily explains why we (this study) as well as previous investigators found that the number of available sites on A α -chains for α_2 AP incorporation were only 20–30% of those potentially available¹⁵⁻¹⁷. In contrast to the relatively high content of α_2 AP that we found in plasma fibrinogen few, if any, intermolecular fibrinogen γ chain cross-links are found in normal plasma, although substantial amounts of these products occur under pathological circumstances, including subjects with familial Mediterranean fever³⁹ and disseminated intravascular coagulation syndromes⁴⁰. Thus, the effective down-regulation

of FXIII-mediated fibrin(ogen) cross-linking in blood contrasts with the relatively high content of α_2 AP in plasma fibrinogen, implying that this interaction is less effectively regulated.

Our present estimations for the α_2 AP content in fibrinogen are between 1.2 and 1.8 moles per mole fibrinogen. Since the plasma fibrinogen level is $\sim 9 \mu\text{M}$, the plasma concentration of α_2 AP that has been incorporated in the fibrinogen compartment lies between 11 and 16 μM . The reported level of plasma α_2 AP is, however, only $\sim 1 \mu\text{M}$ (17 *inter alia*), a value that vastly underestimates the total amount of α_2 AP in plasma. The lower estimate is derived from radial immunodiffusion or rocket immunoelectrophoresis measurements that would be mainly if not solely, a measure of 'free' α_2 AP, because these techniques are not likely to have taken into account the relatively large amount of more slowly diffusing fibrinogen-bound α_2 AP. Thus, except for this present study and two earlier ones that focused on the inhibitor content of fibrinogen itself (23, 24), the existence of α_2 AP covalently bound to fibrinogen in plasma has been overlooked.

Congenital homozygous α_2 AP deficiency causes a severe hemorrhagic disorder characterized by increased susceptibility of intravascular thrombi to fibrinolysis⁵⁻⁷, underscoring the importance of α_2 AP for down-regulating fibrinolysis. Observations on this catastrophic hemorrhagic condition have not, however, addressed an obverse possibility, namely that incorporation of α_2 AP into plasma fibrinogen prior to the initiation of clotting and subsequent fibrinolysis, serves an important role in its own right in down-regulating the rate of fibrinolysis. This notion is supported by the fact that α_2 AP is an effective inhibitor of fibrinolysis only after it has been incorporated into fibrin¹⁰⁻¹³. The extent to which the α_2 AP incorporation into fibrinogen varies from individual to individual remains to be investigated, but it is tempting to speculate that the degree of α_2 AP incorporation plays an important role in the fibrinolytic response *in vivo*. In summary, we propose that incorporation of α_2 AP into circulating fibrinogen prior to initiation of blood clotting plays an important role in down-regulating fibrinolysis, thus suggesting a new role for plasma FXIII in regulation of fibrinolysis.

Acknowledgements

This study was supported by NIH Grants R01 HL-70627 and HL-72995. We thank J. S. Finlayson for his detailed review of this manuscript.

References

- 1 Aoki H, Harpel PC. Inhibitors in the fibrinolytic enzyme system. *Semin Thromb Haemost* 1984; 10: 24–41.
- 2 Collen D. Identification and some properties of a new fast reacting plasmin inhibitor in human plasma. *Eur J Biochem* 1976; 69: 209–16.
- 3 Moroi M, Aoki N. Isolation and characterization of alpha2-plasmin inhibitor from human plasma. A novel proteinase inhibitor which inhibits activator-induced clot lysis. *J Biol Chem* 1976; 251: 5956–65.
- 4 Mullertz S, Clemmensen I. The primary inhibitor of plasmin in human plasma. *Biochem J* 1976; 159: 545–53.
- 5 Aoki N, Saito H, Kamiya T, Koie K, Sakata Y, Kobakura M. Congenital deficiency of alpha 2-plasmin inhibitor associated with severe hemorrhagic tendency. *J Clin Invest* 1979; 63: 877–84.
- 6 Aoki N, Sakata Y. Influence of alpha 2-plasmin inhibitor on adsorption of plasminogen to fibrin. *Thromb Res* 1980; 19: 149–55.

- 7 Saito H. Alpha 2-plasmin inhibitor and its deficiency states. *J Lab Clin Med* 1988; 112: 671– 8.
- 8 Kimura S, Aoki N. Cross-linking site in fibrinogen for alpha 2-plasmin inhibitor. *J Biol Chem* 1986; 261: 15591– 5.
- 9 Ritchie H, Lawrie LC, Crombie PW, Mosesson MW, Booth NA. Cross-linking of plasminogen activator inhibitor 2 and {alpha}2- antiplasmin to fibrin(ogen). *J Biol Chem* 2000; 275: 24915– 20.
- 10 Moroi M, Aoki N. On the interaction of alpha2-plasmin inhibitor and proteases. Evidence for the formation of a covalent crosslinkage and non-covalent weak bondings between the inhibitor and proteases. *Biochim Biophys Acta* 1977; 482: 412– 20.
- 11 Aoki N, Moroi M, Tachiya K. Effects of alpha2-plasmin inhibitor on fibrin clot lysis. Its comparison with alpha2-macroglobulin. *Thromb Haemost* 1978; 39: 22– 31.
- 12 Sakata Y, Aoki H. Significance of cross-linking of α_2 -plasmin inhibitor to fibrin in inhibition of fibrinolysis and in hemostasis. *J Clin Invest* 1982; 69: 536– 42.
- 13 Weitz JI, Leslie B, Hirsh J, Klement P. Alpha 2-antiplasmin supplementation inhibits tissue plasminogen activator-induced fibrinogenolysis and bleeding with little effect on thrombolysis. *J Clin Invest* 1993; 91: 1343– 50.
- 14 Lee AY, Fredenburgh JC, Stewart RJ, Rischke JA, Weitz JI. Like fibrin, (DD)E, the major degradation product of crosslinked fibrin, protects plasmin from inhibition by alpha2-antiplasmin. *Thromb Haemost* 2001; 85: 502– 8.
- 15 Tamaki T, Aoki H. Cross-linking of α_2 -plasmin inhibitor and fibronectin to fibrin by fibrin-stabilizing factor. *Biochim Biophys Acta* 1981; 661: 280– 6.
- 16 Ichinose A, Aoki N. Reversible cross-linking of alpha 2-plasmin inhibitor to fibrinogen by fibrin-stabilizing factor. *Biochim Biophys Acta* 1982; 706: 158– 64.
- 17 Sakata Y, Aoki N. Cross-linking of alpha 2-plasmin inhibitor to fibrin by fibrin-stabilizing factor. *J Clin Invest* 1980; 65: 290– 7.
- 18 Koyama T, Koike Y, Toyota S, Miyagi F, Suzuki N, Aoki N. Different NH2-terminal form with 12 additional residues of alpha 2-plasmin inhibitor from human plasma and culture media of Hep G2 cells. *Biochem Biophys Res Commun* 1994; 200: 417– 22.
- 19 Bangert K, Johnsen AH, Christensen U, Thorsen S. Different N-terminal forms of alpha 2-plasmin inhibitor in human plasma. *Biochem J* 1993; 291 (Pt 2): 623– 5.
- 20 Christiansen VJ, Jackson KW, Lee KN, McKee PA. The effect of a single nucleotide polymorphism on human alpha 2-antiplasmin activity. *Blood* 2007; 109: 5286– 92.
- 21 Lee KN, Jackson KW, Christiansen VJ, Chung KH, McKee PA. A novel plasma proteinase potentiates alpha2-antiplasmin inhibition of fibrin digestion. *Blood* 2004; 103: 3783– 8.
- 22 Sumi Y, Ichikawa Y, Nakamura Y, Miura O, Aoki N. Expression and characterization of pro alpha 2-plasmin inhibitor. *J Biochem (Tokyo)* 1989; 106: 703– 7.
- 23 Mosesson MW, Finlayson JS. Biochemical and chromatographic studies of certain activities associated with human fibrinogen preparations. *J Clin Invest* 1963; 42: 747– 55.
- 24 Siebenlist KR, Mosesson MW, Meh DA, DiOrio JP, Albrecht RM, Olson JD. Coexisting dysfibrinogenemia (gammaR275C) and factor V Leiden deficiency associated with thromboembolic disease (fibrinogen Cedar Rapids). *Blood Coagul Fibrinolysis* 2000; 11: 293– 304.
- 25 Siebenlist KR, Meh D, Mosesson MW. Protransglutaminase (factor XIII) mediated crosslinking of fibrinogen and fibrin. *Thromb Haemost* 2001; 86: 1221– 8.
- 26 Mosesson MW, Sherry S. The preparation and properties of human fibrinogen of relatively high solubility. *Biochemistry* 1966; 5: 2829– 35.

- 27 Siebenlist KR, Mosesson MW. Evidence for intramolecular cross-linked A α chain heterodimers in plasma fibrinogen. *Biochemistry* 1996; 35: 5817– 21.
- 28 Mosesson MW, Galanakis DK, Finlayson JS. Comparison of human plasma fibrinogen subfractions and early plasmic fibrinogen derivatives. *J Biol Chem* 1974; 249: 4656– 64.
- 29 Henschen A, Lottspeich F, Kehl M, Southan C. Covalent structure of fibrinogen. *Ann NY Acad Sci* 1983; 408: 28– 43.
- 30 Veklich YI, Gorkun OV, Medved LV, Niewenhuizen W, Weisel JW. Carboxyl-terminal portions of the α chains of fibrinogen and fibrin. *J Biol Chem* 1993; 268: 13577– 85.
- 31 Lorand L, Gotoh T. Fibrinolytic factor. The fibrin stabilizing factor. *Methods Enzymol* 1970; 19: 770– 82.
- 32 Loewy AG, Dunathan K, Kriegl R, Wolfinger HL. Fibrinase I. Purification of substrate and enzyme. *J Biol Chem* 1961; 236: 2625– 33.
- 33 Schwartz ML, Pizzo SV, Hill RL, McKee PA. Human Factor XIII from plasma and platelets. Molecular weights, subunit structures, proteolytic activation, and cross-linking of fibrinogen and fibrin. *J Biol Chem* 1973; 248: 1395– 407.
- 34 Lee KN, Lee CS, Tae WC, Jackson KW, Christiansen VJ, McKee PA. Cross-linking of wild-type and mutant alpha 2-antiplasmins to fibrin by activated factor XIII and by a tissue transglutaminase. *J Biol Chem* 2000; 275: 37382– 9.
- 35 Martin BE, Wasiewski WW, Fenton JW II, Detwiler TC. Equilibrium binding of thrombin to platelets. *Biochemistry* 1976; 15: 4886– 93.
- 36 Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970; 227: 680– 5.
- 37 Mosesson MW, Hernandez I, Raife TJ, Medved L, Yakovlev S, Simpson-Haidaris PJ, Uitte de Willige S, Bertina R. Plasma fibrinogen gamma' chain content in the thrombotic microangiopathy syndrome. *J Thromb Haemost* 2007; 5: 62– 9.
- 38 Tamaki T, Aoki N. Cross-linking of alpha 2-plasmin inhibitor to fibrin catalyzed by activated fibrin-stabilizing factor. *J Biol Chem* 1982; 257: 14767– 72.
- 39 Mosesson MW, Wautier JL, Amrani DL, Dervichian M, Cattani D. Evidence for circulating fibrin in familial Mediterranean fever. *J Lab Clin Med* 1982; 99: 559– 67.
- 40 Mosesson MW, Colman RW, Sherry S. Chronic intravascular coagulation syndrome. *N Engl J Med* 1968; 278: 815– 21.