

7-24-2013

Extremal H-Colorings of Graphs with Fixed Minimum Degree

John Engbers

Marquette University, john.engbers@marquette.edu

Extremal H -colorings of graphs with fixed minimum degree

John Engbers*

July 24, 2013

Abstract

For graphs G and H , a homomorphism from G to H , or H -coloring of G , is a map from the vertices of G to the vertices of H that preserves adjacency. When H is composed of an edge with one looped endvertex, an H -coloring of G corresponds to an independent set in G . Galvin showed that, for sufficiently large n , the complete bipartite graph $K_{\delta, n-\delta}$ is the n -vertex graph with minimum degree δ that has the largest number of independent sets.

In this paper, we begin the project of generalizing this result to arbitrary H . Writing $\text{hom}(G, H)$ for the number of H -colorings of G , we show that for fixed H and $\delta = 1$ or $\delta = 2$,

$$\text{hom}(G, H) \leq \max\{\text{hom}(K_{\delta+1}, H)^{\frac{n}{\delta+1}}, \text{hom}(K_{\delta, \delta}, H)^{\frac{n}{2\delta}}, \text{hom}(K_{\delta, n-\delta}, H)\}$$

for any n -vertex G with minimum degree δ (for sufficiently large n). We also provide examples of H for which the maximum is achieved by $\text{hom}(K_{\delta+1}, H)^{\frac{n}{\delta+1}}$ and other H for which the maximum is achieved by $\text{hom}(K_{\delta, \delta}, H)^{\frac{n}{2\delta}}$. For $\delta \geq 3$ (and sufficiently large n), we provide an infinite family of H for which $\text{hom}(G, H) \leq \text{hom}(K_{\delta, n-\delta}, H)$ for any n -vertex G with minimum degree δ . The results generalize to weighted H -colorings.

1 Introduction and statement of results

Let $G = (V(G), E(G))$ be a finite simple graph. A *homomorphism* from G to a finite graph $H = (V(H), E(H))$ (without multi-edges but perhaps with loops) is a map from $V(G)$ to $V(H)$ that preserves edge adjacency. We write

$$\text{Hom}(G, H) = \{f : V(G) \rightarrow V(H) \mid v \sim_G w \implies f(v) \sim_H f(w)\}$$

*johnengbers@gmail.com; Department of Mathematics, University of Notre Dame, Notre Dame IN 46556

for the set of all homomorphisms from G to H , and $\text{hom}(G, H)$ for $|\text{Hom}(G, H)|$. All graphs mentioned in this paper will be finite without multiple edges. Those denoted by G will always be loopless, while those denoted by H may possibly have loops. We will also assume that H has no isolated vertices.

Graph homomorphisms generalize a number of important notions in graph theory. When $H = H_{\text{ind}}$, the graph consisting of a single edge and a loop on one endvertex, elements of $\text{Hom}(G, H_{\text{ind}})$ can be identified with the independent sets in G . When $H = K_q$, the complete graph on q vertices, elements of $\text{Hom}(G, K_q)$ can be identified with the proper q -colorings of G . Motivated by this latter example, elements of $\text{Hom}(G, H)$ are sometimes referred to as H -colorings of G , and the vertices of H are referred to as *colors*. We will utilize this terminology throughout the paper.

In statistical physics, H -colorings have a natural interpretation as configurations in *hard-constraint spin systems*. Here, the vertices of G are thought of as sites that are occupied by particles, with the edges of G representing pairs of bonded sites (for example by spatial proximity). The vertices of H represent the possible spins that a particle may have, and the occupation rule is that spins appearing on bonded sites must be adjacent in H . A valid configuration of spins on G is exactly an H -coloring of G . In the language of statistical physics, independent sets are configurations in the *hard-core gas model*, and proper q -colorings are configurations in the *zero-temperature q -state antiferromagnetic Potts model*. Another example comes from the *Widom-Rowlinson* graph $H = H_{\text{WR}}$, the fully-looped path on three vertices. If the endpoints of the path represent different particles and the middle vertex represents empty space, then the Widom-Rowlinson graph models the occupation of space by two mutually repelling particles.

Fix a graph H . A natural extremal question to ask is the following: for a given family of graphs \mathcal{G} , which graphs G in \mathcal{G} maximize $\text{hom}(G, H)$? If we assume that all graphs in \mathcal{G} have n vertices, then there are several cases where this question has a trivial answer. First, if $H = K_q^{\text{loop}}$, the fully looped complete graph on q vertices, then every map $f : V(G) \rightarrow V(H)$ is an H -coloring (and so $\text{hom}(G, K_q^{\text{loop}}) = q^n$). Second, if the empty graph \overline{K}_n is contained in \mathcal{G} , then again every map $f : V(\overline{K}_n) \rightarrow V(H)$ is an H -coloring (and so $\text{hom}(\overline{K}_n, H) = |V(H)|^n$). Motivated by this second trivial case, it is interesting to consider families \mathcal{G} for which each $G \in \mathcal{G}$ has many edges.

For the family of n -vertex m -edge graphs, this question was first posed for $H = K_q$ around 1986, independently, by Linial and Wilf. Lazebnik provided an answer for $q = 2$ [15], but for general q there is still not a complete answer. However, much progress has been made (see [16] and the references therein). Recently, Cutler and Radcliffe answered this question for $H = H_{\text{ind}}$, $H = H_{\text{WR}}$, and some other small H [2, 3]. A feature of the family of n -vertex, m -edge graphs emerging from the partial results mentioned is that there seems to be no uniform answer to the question, “which G in the family maximizes $\text{hom}(G, H)$?”, with the answers depending very

sensitively on the choice of H .

Another interesting family to consider is the family of n -vertex d -regular graphs. Here, Kahn [13] used entropy methods to show that every *bipartite* graph G in this family satisfies $\text{hom}(G, H_{\text{ind}}) \leq \text{hom}(K_{d,d}, H_{\text{ind}})^{\frac{n}{2d}}$, where $K_{d,d}$ is the complete bipartite graph with d vertices in each partition class. Notice that when $2d|n$ this bound is achieved by $\frac{n}{2d}K_{d,d}$, the disjoint union of $n/2d$ copies of $K_{d,d}$. Galvin and Tetali [11] generalized this entropy argument, showing that for *any* H and any bipartite G in this family,

$$\text{hom}(G, H) \leq \text{hom}(K_{d,d}, H)^{\frac{n}{2d}}. \quad (1)$$

Kahn conjectured that (1) should hold for $H = H_{\text{ind}}$ for all (not necessarily bipartite) G , and Zhao [19] resolved this conjecture affirmatively, deducing the general result from the bipartite case. Interestingly, (1) does not hold for general H when biparticity is dropped, as there are examples of n , d , and H for which $\frac{n}{d+1}K_{d+1}$, the disjoint union of $n/(d+1)$ copies of the complete graph K_{d+1} , maximizes the number of H -colorings of graphs in this family. (For example, take H to be the disjoint union of two looped vertices; here $\log_2(\text{hom}(G, H))$ equals the number of components of G .) Galvin proposes the following conjecture in [8].

Conjecture 1.1. *Let G be an n -vertex d -regular graph. Then, for any H ,*

$$\text{hom}(G, H) \leq \max\{\text{hom}(K_{d+1}, H)^{\frac{n}{d+1}}, \text{hom}(K_{d,d}, H)^{\frac{n}{2d}}\}.$$

When $2d(d+1)|n$, this bound is achieved by either $\frac{n}{2d}K_{d,d}$ or $\frac{n}{d+1}K_{d+1}$. Evidence for this conjecture is given by Zhao [19, 20], who provided a large class of H for which $\text{hom}(G, H) \leq \text{hom}(K_{d,d}, H)^{\frac{n}{2d}}$. Galvin [8, 9] provides further results for various H (including triples (n, d, H) for which $\text{hom}(G, H) \leq \text{hom}(K_{d+1}, H)^{\frac{n}{d+1}}$) and asymptotic evidence for the conjecture.

It is clear that Conjecture 1.1 is true when $d = 1$, since the graph consisting of $n/2$ disjoint copies of an edge is the only 1-regular graph on n vertices. We prove the conjecture for $d = 2$ and also characterize the cases of equality.

Theorem 1.2. *Let G be an n -vertex 2-regular graph. Then, for any H ,*

$$\text{hom}(G, H) \leq \max\{\text{hom}(C_3, H)^{\frac{n}{3}}, \text{hom}(C_4, H)^{\frac{n}{4}}\}.$$

If $H \neq K_q^{\text{loop}}$, the only graphs achieving equality are $G = \frac{n}{3}C_3$ (when $\text{hom}(C_3, H)^{\frac{n}{3}} > \text{hom}(C_4, H)^{\frac{n}{4}}$), $G = \frac{n}{4}C_4$ (when $\text{hom}(C_3, H)^{\frac{n}{3}} < \text{hom}(C_4, H)^{\frac{n}{4}}$), or the disjoint union of copies of C_3 and copies of C_4 (when $\text{hom}(C_3, H)^{\frac{n}{3}} = \text{hom}(C_4, H)^{\frac{n}{4}}$).

It is possible for each of the equality conditions in Theorem 1.2 to occur. The first two situations arise when H is a disjoint union of two looped vertices and $H = K_2$, respectively. For the third situation, we utilize that if G is connected and H is the

disjoint union of H_1 and H_2 , then $\text{hom}(G, H) = \text{hom}(G, H_1) + \text{hom}(G, H_2)$. Letting H be the disjoint union of 8 copies of a single looped vertex and 4 copies of K_2 gives $\text{hom}(C_3, H)^{\frac{1}{3}} = \text{hom}(C_4, H)^{\frac{1}{4}} = 2$.

Another natural and related family to study is $\mathcal{G}(n, \delta)$, the set of all n -vertex graphs with minimum degree δ . Our question here becomes: for a given H , which $G \in \mathcal{G}(n, \delta)$ maximizes $\text{hom}(G, H)$? Since removing edges increases the number of H -colorings, it is tempting to believe that the answer to this question will be a graph that is δ -regular (or close to δ -regular). This in fact is not the case, even for $H = H_{\text{ind}}$. The following result appears in [7].

Theorem 1.3. *For $\delta \geq 1$, $n \geq 8\delta^2$, and $G \in \mathcal{G}(n, \delta)$, we have*

$$\text{hom}(G, H_{\text{ind}}) \leq \text{hom}(K_{\delta, n-\delta}, H_{\text{ind}}),$$

with equality only for $G = K_{\delta, n-\delta}$.

Recently, Cutler and Radcliffe [4] have extended Theorem 1.3 to the range $n \geq 2\delta$. Further results related to maximizing the number of independent sets of a fixed size for $G \in \mathcal{G}(n, \delta)$ can be found in e.g. [1, 6].

With Conjecture 1.1 and Theorem 1.3 in mind, the following conjecture is natural.

Conjecture 1.4. *Fix $\delta \geq 1$ and H . There exists a constant $c(\delta, H)$ (depending on δ and H) such that for $n \geq c(\delta, H)$ and $G \in \mathcal{G}(n, \delta)$,*

$$\text{hom}(G, H) \leq \max\{\text{hom}(K_{\delta+1}, H)^{\frac{n}{\delta+1}}, \text{hom}(K_{\delta, \delta}, H)^{\frac{n}{2\delta}}, \text{hom}(K_{\delta, n-\delta}, H)\}.$$

This conjecture stands in marked contrast to the situation for the family of n -vertex m -edge graphs, where each choice of H seems to create a different set of extremal graphs. Here, we conjecture that for *any* H , one of exactly three situations can occur. For $2\delta(\delta+1)|n$ and n large, this represents the best possible conjecture, since for H consisting of a disjoint union of two looped vertices, $H = K_2$, and $H = H_{\text{ind}}$, the number of H -colorings of a graph $G \in \mathcal{G}(n, \delta)$ is maximized by $G = \frac{n}{\delta+1}K_{\delta+1}$, $G = \frac{n}{2\delta}K_{\delta, \delta}$, and $G = K_{\delta, n-\delta}$, respectively.

The purpose of this paper is to make progress toward Conjecture 1.4. We resolve the conjecture for $\delta = 1$ and $\delta = 2$, and characterize the graphs that achieve equality. We also find an infinite family of H for which $\text{hom}(G, H) \leq \text{hom}(K_{\delta, n-\delta}, H)$ for all $G \in \mathcal{G}(n, \delta)$ (for sufficiently large n), with equality only for $G = K_{\delta, n-\delta}$. Before we formally state these theorems, we highlight the degree conventions and notations that we will follow for the remainder of the paper.

Convention. For $v \in V(H)$, let $d(v)$ denote the degree of v , where loops count *once* toward the degree. While δ will always refer to the minimum degree of a graph G , Δ will always denote the maximum degree of a graph H (unless explicitly stated otherwise).

Theorem 1.5. ($\delta = 1$). Fix H , $n \geq 2$ and $G \in \mathcal{G}(n, 1)$.

1. Suppose that $H \neq K_{\Delta}^{\text{loop}}$ satisfies $\sum_{v \in V(H)} d(v) \geq \Delta^2$. Then

$$\text{hom}(G, H) \leq \text{hom}(K_2, H)^{\frac{n}{2}},$$

with equality only for $G = \frac{n}{2}K_2$.

2. Suppose that H satisfies $\sum_{v \in V(H)} d(v) < \Delta^2$, and let $n_0 = n_0(H)$ be the smallest integer in $\{3, 4, \dots\}$ satisfying $\sum_{v \in V(H)} d(v) < \left(\sum_{v \in V(H)} d(v)^{n_0-1}\right)^{\frac{2}{n_0}}$.

- (a) If $2 \leq n < n_0$, then

$$\text{hom}(G, H) \leq \text{hom}(K_2, H)^{\frac{n}{2}},$$

with equality only for $G = \frac{n}{2}K_2$ [unless $n = n_0 - 1$ and $\sum_{v \in V(H)} d(v) = \left(\sum_{v \in V(H)} d(v)^{n_0-2}\right)^{\frac{2}{n_0-1}}$, in which case $G = K_{1, n-1}$ also achieves equality].

- (b) If $n \geq n_0$, then

$$\text{hom}(G, H) \leq \text{hom}(K_{1, n-1}, H),$$

with equality only for $G = K_{1, n-1}$.

Remark. Notice that $\text{hom}(K_2, H) = \sum_{v \in V(H)} d(v)$, so the conditions on H in Theorem 1.5 may also be written as $\text{hom}(K_2, H)^{\frac{1}{2}} \geq \Delta$ and $\text{hom}(K_2, H)^{\frac{1}{2}} < \Delta$.

Theorem 1.6. ($\delta = 2$). Fix H .

1. Suppose that $H \neq K_{\Delta}^{\text{loop}}$ satisfies $\max\{\text{hom}(C_3, H)^{\frac{1}{3}}, \text{hom}(C_4, H)^{\frac{1}{4}}\} \geq \Delta$. Then for all $n \geq 3$ and $G \in \mathcal{G}(n, 2)$,

$$\text{hom}(G, H) \leq \max\{\text{hom}(C_3, H)^{\frac{n}{3}}, \text{hom}(C_4, H)^{\frac{n}{4}}\},$$

with equality only for $G = \frac{n}{3}C_3$ (when $\text{hom}(C_3, H)^{\frac{n}{3}} > \text{hom}(C_4, H)^{\frac{n}{4}}$), $G = \frac{n}{4}C_4$ (when $\text{hom}(C_3, H)^{\frac{n}{3}} < \text{hom}(C_4, H)^{\frac{n}{4}}$), or the disjoint union of C_3 and copies of C_4 (when $\text{hom}(C_3, H)^{\frac{n}{3}} = \text{hom}(C_4, H)^{\frac{n}{4}}$).

2. Suppose that H satisfies $\max\{\text{hom}(C_3, H)^{\frac{1}{3}}, \text{hom}(C_4, H)^{\frac{1}{4}}\} < \Delta$. Then there exists a constant c_H such that for $n > c_H$ and $G \in \mathcal{G}(n, 2)$,

$$\text{hom}(G, H) \leq \text{hom}(K_{2, n-2}, H),$$

with equality only for $G = K_{2, n-2}$.

Theorems 1.5 and 1.6 are easily seen to resolve Conjecture 1.4 when $\delta = 1$ and $\delta = 2$, respectively. Notice that if G' is obtained from G by deleting some edges from G , then $\text{hom}(G, H) \leq \text{hom}(G', H)$. Because of this, their proofs focus on G which are *edge-min-critical* for δ , meaning that the minimum degree of G is δ and for any edge e the minimum degree of $G - e$ is $\delta - 1$.

The edge-min-critical graphs in $\mathcal{G}(n, 1)$ are disjoint unions of stars, and the proof of Theorem 1.5 critically uses this fact. Theorem 1.6 relies on a structural characterization of edge-min-critical graphs in $\mathcal{G}(n, 2)$ (see Lemma 5.1) and also uses Theorem 1.2. The global structure of edge-min-critical graphs in $\mathcal{G}(n, \delta)$ for $\delta \geq 3$ is not very well understood.

We also make some progress in the general δ case of Conjecture 1.4 by providing an infinite family of H for which $K_{\delta, n-\delta}$ has the largest number of H -colorings.

Theorem 1.7. *Fix δ and H . Suppose that H satisfies $\sum_{v \in V(H)} d(v) < \Delta^2$. Then there exists a constant c_H such that for all $n \geq (c_H)^\delta$ and $G \in \mathcal{G}(n, \delta)$,*

$$\text{hom}(G, H) \leq \text{hom}(K_{\delta, n-\delta}, H),$$

with equality only for $G = K_{\delta, n-\delta}$.

If H also has the property that all vertices of degree Δ share the same Δ neighbors, then the same result holds for all $n \geq c_H \delta^2$.

The following corollary warrants special attention, and is immediate.

Corollary 1.8. *Suppose that $H \neq K_{\Delta}^{\text{loop}}$ has a looped dominating vertex, or that H satisfies $\sum_{v \in V(H)} d(v) < \Delta^2$ and H has a unique vertex of degree Δ . Then there exists a constant c_H such that for all $n \geq c_H \delta^2$ and $G \in \mathcal{G}(n, \delta)$,*

$$\text{hom}(G, H) \leq \text{hom}(K_{\delta, n-\delta}, H),$$

with equality only for $G = K_{\delta, n-\delta}$.

The graphs $H = H_{\text{ind}}$ and $H = H_{\text{WR}}$ satisfy the conditions of Corollary 1.8, so in particular we provide an alternate proof of Galvin's result for $H = H_{\text{ind}}$ [7]. Another graph H which satisfies the conditions of Corollary 1.8 is the k -state hard-core constraint graph $H(k)$ ($k \geq 1$), the graph with vertex set $\{0, 1, \dots, k\}$ and edge $i \sim_{H(k)} j$ if $i + j \leq k$. This graph naturally occurs in the study of multicast communications networks, and has been considered in e.g. [10, 17].

Notice that the condition on H in Theorem 1.7 is necessary but not sufficient for $\text{hom}(G, H) \leq \text{hom}(K_{\delta, n-\delta}, H)$ for all $G \in \mathcal{G}(n, \delta)$. Indeed, if H is a path on three vertices with a loop on one endpoint of the path, then $\sum_{v \in V(H)} d(v) = 5$ while $\Delta = 2$. However, for large enough n and $G \in \mathcal{G}(n, 2)$, $\text{hom}(G, H) \leq \text{hom}(K_{2, n-2}, H)$, as can be seen by computing $\text{hom}(C_3, H)$, $\text{hom}(C_4, H)$, and applying Theorem 1.6.

It is also interesting to consider a maximum degree condition in addition to a minimal degree condition (see e.g. [1, 12, 13]). Let $\mathcal{G}(n, \delta, D)$ denote the set of graphs on n vertices with minimum degree δ and maximum degree at most D . Which graphs $G \in \mathcal{G}(n, \delta, D)$ maximize $\text{hom}(G, H)$? For a fixed δ , this question is interesting for the H with the property that $\text{hom}(G, H) \leq \text{hom}(K_{\delta, n-\delta}, H)$ for all $G \in \mathcal{G}(n, \delta)$, as $K_{\delta, n-\delta} \in \mathcal{G}(n, \delta, D)$ only when $D \geq n - \delta$.

For $\delta = 1$ and any $D \geq 1$, we provide an answer.

Theorem 1.9. *Fix H and $D \geq 1$. For any $G \in \mathcal{G}(n, 1, D)$,*

$$\text{hom}(G, H) \leq \max\{\text{hom}(K_2, H)^{\frac{n}{2}}, \text{hom}(K_{1,D}, H)^{\frac{n}{1+D}}\},$$

with the cases of equality as in Theorem 1.5 (where $K_{1, n-1}$ is replaced by $\frac{n}{1+D}K_{1,D}$).

From a statistical physics standpoint, there is a very natural family of probability distributions that can be put on $\text{Hom}(G, H)$. Fix a set of positive weights $\Lambda = \{\lambda_i : i \in V(H)\}$ indexed by the vertices of H . We think of λ_i as representing the likelihood of particle i appearing at a site in G , and formalize this by giving an element $f \in \text{Hom}(G, H)$ weight $w_\Lambda(f) = \prod_{v \in V(G)} \lambda_{f(v)}$ and probability

$$p_\Lambda(f) = \frac{w_\Lambda(f)}{Z_\Lambda(G, H)},$$

where $Z_\Lambda(G, H) = \sum_f w_\Lambda(f)$ is the appropriate normalizing constant (or *partition function*) of the model. By taking $\lambda_i = 1$ for each i , $p_\Lambda(f)$ is the uniform distribution on $\text{Hom}(G, H)$ and in this case $Z_\Lambda(G, H) = \text{hom}(G, H)$.

Interestingly, several proofs of structural results about H -colorings *require* passing to the weighted model first; see e.g. [5, 13, 14]. Our results generalize naturally to weighted H -colorings. Although the proofs of the weighted versions come with almost no extra effort, for the clarity of presentation we defer this discussion until Section 6.

The paper is laid out as follows. In Section 2, we prove Theorem 1.7 by partitioning $\mathcal{G}(n, \delta)$ based on the size of a maximal matching. Section 3 utilizes the structure of edge-min-critical graphs for $\delta = 1$ to prove Theorems 1.5 and 1.9. By analyzing H -colorings of cycles, we prove Theorem 1.2 in Section 4. Following some preliminary lemmas about edge-min-critical graphs for $\delta = 2$ and H -colorings of paths, we prove Theorem 1.6 in Section 5. Finally, in Section 6 we comment on the generalization of our results to weighted H -colorings and also present some related questions.

2 Proof of Theorem 1.7

Suppose that we have a graph H satisfying

$$\sum_{v \in V(H)} d(v) < \Delta^2, \tag{2}$$

and let G be a graph with minimum degree δ . Let M be the edge set of a matching of maximum size in G and I the set of unmatched vertices.

We first derive some structural properties of our graph G based on M . Since M is maximal, I forms an independent set. Furthermore, suppose that x_1 and x_2 in $V(G)$ are matched in M . If x_1 has at least two edges into I , then x_2 cannot be adjacent to any vertex in I , as this would create an augmenting path of length 3 and therefore a matching of larger size. In summary:

$$\begin{aligned} \text{At most one vertex in an edge of the matching } M \text{ can have degree at} \\ \text{least 2 into } I, \text{ and if one has degree at least 2 into } I \text{ then the other} \quad (3) \\ \text{has degree 0 into } I. \end{aligned}$$

For each edge in M , put the endpoint with the largest degree into I in a set $J \subset V(G)$, and put the other endpoint in a set $K \subset V(G)$. (If the degrees are equal, make an arbitrary choice.) A schematic picture of G is shown in Figure 2; there are at most $|K| = |M|$ total edges between I and K .



Figure 1: The relevant structure for G .

Also, if there are more than $|M|$ vertices in I that are adjacent to both endpoints of some edge in M , then by the pigeonhole principle there are distinct $y_1, y_2 \in I$ that are adjacent to both endpoints of some fixed edge in M . This would force both endpoints of an edge in M to have degree at least 2 into I , contradicting (3). Therefore we have:

$$\text{There are at most } |M| \text{ vertices in } I \text{ adjacent to both endpoints of} \quad (4) \\ \text{some edge in } M.$$

In particular, suppose $n \geq 3\delta - 2$. Then if $|M| < \delta$ we have $|I| \geq \delta$. Since each $x \in I$ has at least δ neighbors to M , each $x \in I$ is adjacent to both endpoints of some edge in M . Since $|I| \geq \delta > |M|$, this contradicts (4). Therefore if $n \geq 3\delta - 2$ we have $|M| \geq \delta$ for all $G \in \mathcal{G}(n, \delta)$. We will first analyze the graphs where $|M| = \delta$ and then the graphs where $|M| > \delta$.

Case 1: Suppose that $|M| = \delta$ and $n > 3\delta$, so by (4) at most δ vertices in I are adjacent to both endpoints of some edge in M . Then there is at least one vertex in I that is adjacent to exactly one endpoint of each edge in M . However, this shows that *no* vertex in I can be adjacent to both endpoints of any edge in M , since any

vertex in I adjacent to both vertices of an edge in M would force one endpoint of M to have degree at least 2 into I and the other endpoint of M to have degree at least 1 into I (contradicting (3)). It follows that each vertex in I must be adjacent to each vertex in J , and so by (3) there are no edges between I and K .

Now suppose $k_1, k_2 \in K$ with $k_1 \sim k_2$. Then there exist distinct $j_1, j_2 \in J$ with $k_1 \sim_M j_1$ and $k_2 \sim_M j_2$. Letting i_1 and i_2 denote any two distinct vertices in I (and recalling that everything in I is adjacent to everything in J), $i_1 \sim j_1 \sim k_1 \sim k_2 \sim j_2 \sim i_2$ is an augmenting path of length 5, which contradicts the maximality of M . Therefore K is an independent set and so $K \cup I$ is an independent set. Since G has minimum degree δ , every vertex in $K \cup I$ is adjacent to every vertex in J , and so G must be the complete bipartite graph $K_{\delta, n-\delta}$ with some edges added to the size δ partition class.

We now show that adding any edge to the size δ partition class in $K_{\delta, n-\delta}$ will strictly decrease the number of H -colorings. Since H cannot contain K_{Δ}^{loop} (by (2)), there are two (possibly non-distinct) non-adjacent neighbors of a vertex in H with degree Δ . If any edge is added to the size δ partition class in $K_{\delta, n-\delta}$, then it is impossible for any H -coloring to color the endpoints of that edge with the non-adjacent vertices in H , but such a coloring is possible in $K_{\delta, n-\delta}$. Since any H -coloring of $K_{\delta, n-\delta}$ with an edge added is an H -coloring of $K_{\delta, n-\delta}$, this shows that the number of H -colorings strictly decreases whenever an edge is added to $K_{\delta, n-\delta}$.

In summary, we have shown that if G satisfies $n > 3\delta$, $|M| \leq \delta$, and $G \neq K_{\delta, n-\delta}$ then $\text{hom}(G, H) < \text{hom}(K_{\delta, n-\delta}, H)$.

Case 2: Now suppose that $|M| = k \geq \delta + 1$. We will show that for large enough n we have $\text{hom}(G, H) < \text{hom}(K_{\delta, n-\delta}, H)$, which will complete the proof.

Let $S(\delta, H)$ denote the vectors in $V(H)^\delta$ with the property that the elements of the vector have Δ common neighbors, and let $s(\delta, H) = |S(\delta, H)|$. (Note that $S(\delta, H) \neq \emptyset$, since if $v \in V(H)$ with $d(v) = \Delta$ then $(v, v, \dots, v) \in S(\delta, H)$.) We obtain a lower bound on $\text{hom}(K_{\delta, n-\delta}, H)$ by coloring the size δ partition class using an element of $S(\delta, H)$, and then independently coloring the vertices in the size $n - \delta$ partition class using the Δ common neighbors. This gives

$$\text{hom}(K_{\delta, n-\delta}, H) \geq s(\delta, H) \Delta^{n-\delta}.$$

We will show that for n large and $k \geq \delta + 1$, $\text{hom}(G, H) < s(\delta, H) \Delta^{n-\delta}$.

Our initial coloring scheme will be to color J arbitrarily first, then K , then I , keeping track of an upper bound on the number of choices we have for the color at each vertex. If a vertex in J is colored with $v \in V(H)$, its neighbor in M has at most $d(v)$ choices for a color. Since each vertex in I is adjacent to some vertex in $J \cup K$, there are at most Δ choices for the color of each vertex in I . This gives

$$\text{hom}(G, H) \leq \left(\sum_{v \in V(H)} d(v) \right)^k \Delta^{n-2k} = \Delta^n \left(\frac{\sum_{v \in V(H)} d(v)}{\Delta^2} \right)^k.$$

Recalling that H satisfies (2), if $k > \delta \log \Delta / \log \left(\frac{\Delta^2}{\sum_{v \in V(H)} d(v)} \right) = C_H \delta$ this upper bound is smaller than $\Delta^{n-\delta}$. So we may further assume that $\delta + 1 \leq k \leq C_H \delta$.

Let $I' \subset I$ be the set of vertices in I with neighbors exclusively in J , so by (3) we have $|I'| \geq n - 3k$. Since each vertex in I' has at least δ neighbors in J , we imagine each $x \in I'$ picking a subset of size δ from J (from among the $\binom{k}{\delta}$ possibilities). By the pigeonhole principle there is a set $J_1 \subset J$ with $|J_1| = \delta$ and at least $(n - 3k) / \binom{k}{\delta}$ vertices in I' adjacent to every vertex in J_1 . See Figure 2.




Figure 2: Vertices in I' adjacent to every vertex in J_1 .

We partition the H -colorings of G based on whether the colors on J_1 form a vector in $S(\delta, H)$ or not. If they do, then we next color $J \setminus J_1$, then K , and then I , giving at most

$$s(\delta, H) \cdot \left(\sum_{v \in V(H)} d(v) \right)^{k-\delta} \cdot \Delta^\delta \cdot \Delta^{n-2k}$$

H -colorings of G of this type.

If the colors on J_1 do not form a vector in $S(\delta, H)$, then we have at least $(n - 3k) / \binom{k}{\delta}$ vertices in I (namely those in I') that have at most $\Delta - 1$ choices for their color (here we're using that all edges are present between I' and J_1). Utilizing only this restriction, coloring $J \setminus J_1$, then K , then I gives at most

$$\left(\sum_{v \in V(H)} d(v) \right)^k \Delta^{n-2k} \left(\frac{\Delta - 1}{\Delta} \right)^{\frac{n-3k}{\binom{k}{\delta}}}$$

H -colorings of G of this type. Therefore, using $k \leq C_H \delta$ and $\binom{a}{b} \leq \left(\frac{ea}{b}\right)^b$, we have

$$\begin{aligned} \text{hom}(G, H) &\leq s(\delta, H) \cdot \left(\sum_{v \in V(H)} d(v) \right)^{k-\delta} \Delta^{n-2k+\delta} \\ &\quad + \left(\sum_{v \in V(H)} d(v) \right)^k \Delta^{n-2k} \left(\frac{\Delta-1}{\Delta} \right)^{\frac{n-3k}{\binom{k}{\delta}}} \\ &\leq s(\delta, H) \cdot \left(\frac{\sum_{v \in V(H)} d(v)}{\Delta^2} \right)^{k-\delta} \Delta^{n-\delta} \\ &\quad + \left(\frac{\sum_{v \in V(H)} d(v)}{\Delta^2} \right)^k \Delta^n \left(\frac{\Delta-1}{\Delta} \right)^{\frac{n-3C_H \delta}{(e \cdot C_H)^\delta}} \end{aligned}$$

so that

$$\text{hom}(G, H) \leq s(\delta, H) \Delta^{n-\delta} \left(\frac{\sum_{v \in V(H)} d(v)}{\Delta^2} \right)^{k-\delta} (1 + r_1(\delta, H))$$

where

$$r_1(\delta, H) = \frac{1}{s(\delta, H)} \left(\frac{\sum_{v \in V(H)} d(v)}{\Delta} \right)^\delta \left(\frac{\Delta-1}{\Delta} \right)^{\frac{n-3C_H \delta}{(e \cdot C_H)^\delta}}.$$

For $\delta + 1 \leq k \leq C_H \delta$ and $n \geq (c_H)^\delta$, this is smaller than $s(\delta, H) \Delta^{n-\delta}$.

We sharpen the bounds on n when all of the vertices of H with degree Δ have identical neighborhoods. (Notice that this only requires a new argument for the range $\delta + 1 \leq k \leq C_H \delta$.) Let $V_{=\Delta}(H)$ denote the set of degree Δ vertices in H , and so by assumption each vertex in $V_{=\Delta}(H)$ has the same Δ neighbors (and also $s(\delta, H) = |V_{=\Delta}(H)|^\delta$). Our new strategy is to find a set of δ vertices in J with large degree to I individually instead of finding those with a large common neighborhood in I .

Let $J = \{x_1, \dots, x_k\}$ and let a_t denote the number of edges from x_t to I for each t . Without loss of generality, assume $a_1 \geq a_2 \geq \dots \geq a_k$. Since I is an independent set of size $n - 2k$, there are at least $\delta(n - 2k)$ edges from I to $J \cup K$. Since the degree to I of each vertex in K is at most 1 and each a_t is at most $n - 2k$, we have

$$\delta(n - 2k) - k \leq \sum_{t=1}^k a_t \leq (k - \delta + 1)a_\delta + (\delta - 1)(n - 2k),$$

as $(n - 2k) - k$ is a lower bound on the number of edges from I to J , $a_\delta + a_{\delta+1} + \dots + a_k \leq (k - \delta + 1)a_\delta$ (by the ordering of the a_i 's), and $a_{\delta-1} \leq \dots \leq a_1 \leq n - 2k$.

This gives

$$a_\delta \geq \frac{n - 3k}{k - \delta + 1}.$$

Now set $J_2 = \{x_1, \dots, x_\delta\}$. We first upper bound the number of H -colorings of G that color each vertex in J_2 with a color from $V_{=\Delta}(H)$. By coloring $J \setminus J_2$ arbitrarily, then coloring K , then coloring I , we have at most

$$s(\delta, H) \left(\sum_{v \in V(H)} d(v) \right)^{k-\delta} \Delta^{n-2k+\delta} = s(\delta, H) \Delta^{n-\delta} \left(\frac{\sum_{v \in V(H)} d(v)}{\Delta^2} \right)^{k-\delta} \quad (5)$$

H -colorings of G of this type.

By similar means, we can put an upper bound on the number of H -colorings of G that have some vertex of J_2 colored from $V(H) \setminus V_{=\Delta}(H)$. Here, at least $\frac{n-3k}{k-\delta+1}$ vertices in I will have at most $\Delta - 1$ choices of a color for each coloring of $J \cup K$. Using $k \leq C_H \delta$, we have at most

$$\begin{aligned} & \left(\sum_{v \in V(H)} d(v) \right)^k \Delta^{n-2k} \left(\frac{\Delta - 1}{\Delta} \right)^{\frac{n-3k}{k-\delta+1}} \\ & \leq \Delta^{n-\delta} \left(\frac{\sum_{v \in V(H)} d(v)}{\Delta^2} \right)^k \Delta^\delta \left(\frac{\Delta - 1}{\Delta} \right)^{\frac{n-3C_H \delta}{C_H \delta - \delta + 1}} \end{aligned} \quad (6)$$

H -colorings of G of this type. Combining (5) and (6) we find that

$$\text{hom}(G, H) \leq s(\delta, H) \Delta^{n-\delta} \left(\frac{\sum_{v \in V(H)} d(v)}{\Delta^2} \right)^{k-\delta} (1 + r_2(\delta, H))$$

where

$$r_2(\delta, H) = \frac{1}{s(\delta, H)} \left(\frac{\sum_{v \in V(H)} d(v)}{\Delta} \right)^\delta \left(\frac{\Delta - 1}{\Delta} \right)^{\frac{n-3C_H \delta}{C_H \delta - \delta + 1}}$$

For $\delta + 1 \leq k \leq C_H \delta$ and $n > c_H \delta^2$, this is smaller than $s(\delta, H) \Delta^{n-\delta}$.

3 Proof of Theorems 1.5 and 1.9 ($\delta = 1$)

While Theorem 1.9 is more general than Theorem 1.5, for clarity of presentation we begin with the proof of Theorem 1.5 and later show how to modify the proof to obtain Theorem 1.9. Recall that we will assume that $|V(H)| = q$, and furthermore we will assume that G is edge-min-critical, so G has no edge between two vertices of degree larger than one. (This will give us the inequalities desired; we will address the




Figure 3: A graph G that is the disjoint union of stars; here we may take $n_1 = 3$, $n_2 = 2$, $n_3 = 3$, and $n_4 = 5$.

uniqueness statements in the theorems separately.) In particular, G is the disjoint union of stars, so we can write $G = \cup_i K_{1,n_i-1}$, where $\sum_i n_i = n$. See Figure 3.

Since the stars K_{1,n_i-1} are disjoint and can therefore be colored independently, we have

$$\text{hom}(G, H) = \prod_i \text{hom}(K_{1,n_i-1}, H) = \prod_i \text{hom}(K_{1,n_i-1}, H)^{\frac{n_i}{n_i}}.$$

If x is an integer value in $[2, n]$ that maximizes $\text{hom}(K_{1,x-1}, H)^{\frac{1}{x}}$, then

$$\text{hom}(G, H) \leq \prod_i \text{hom}(K_{1,x-1}, H)^{\frac{n_i}{x}} = \text{hom}(K_{1,x-1}, H)^{\frac{n}{x}}, \quad (7)$$

with equality occurring (when $x|n$) for $\frac{n}{x}K_{1,x-1}$. Because of this, it will be useful to know the integer value(s) of $x \geq 2$ that maximize $\text{hom}(K_{1,x-1}, H)^{\frac{1}{x}}$.

First we derive a formula for $\text{hom}(K_{1,x-1}, H)^{\frac{1}{x}}$ for each integer $x \geq 2$. Notice that all H -colorings of $K_{1,x-1}$ can be obtained by coloring the center of the star with any $v \in V(H)$ and then coloring the leaves (independently) with any neighbor of v , so

$$\text{hom}(K_{1,x-1}, H) = \sum_{v \in V(H)} d(v)^{x-1}. \quad (8)$$

Now (8) holds for each integer $x \geq 2$, and so it will be useful to know the maximum value of

$$\left(\sum_{v \in V(H)} d(v)^{x-1} \right)^{\frac{1}{x}} \quad (9)$$

over all integers $x \geq 2$. In fact, we will study (9) in a slightly more general setting; for the remainder of this proof we will analyze (9) over all *real* numbers $x \geq 2$.

Recall that we are assuming that H has no isolated vertices, so for all $v \in V(H)$ we have $1 \leq d(v) \leq \Delta$. Since there exists a $w \in V(H)$ with $d(w) = \Delta$, we have

$$\left(\sum_{v \in V(H)} d(v)^{x-1} \right)^{\frac{1}{x}} \rightarrow \Delta \quad \text{as } x \rightarrow \infty. \quad (10)$$

To obtain more information, for a fixed real $x \geq 2$ let $a = a(x, H) \in \mathbb{R}$ be such that

$$d(v_1)^{x-1} + \dots + d(v_q)^{x-1} = a^x.$$

Since $1 \leq d(v) \leq \Delta$ for all $v \in V(H)$, for any $\varepsilon > 0$ we have

$$d(v_1)^{x-1+\varepsilon} + \dots + d(v_q)^{x-1+\varepsilon} \leq \Delta^\varepsilon (d(v_1)^{x-1} + \dots + d(v_q)^{x-1}) = \Delta^\varepsilon a^x, \quad (11)$$

with strict inequality if $d(v_i) < \Delta$ for some i . Therefore for any $\varepsilon > 0$ (11) gives

$$a > \Delta \quad \Longrightarrow \quad \left(\sum_{v \in V(H)} d(v)^{x-1+\varepsilon} \right)^{\frac{1}{x+\varepsilon}} < \left(\sum_{v \in V(H)} d(v)^{x-1} \right)^{\frac{1}{x}}. \quad (12)$$

If $a = \Delta$ and $d(v_i) = \Delta$ for all i , then (11) gives $q\Delta^{x-1} = \Delta^x$ so $H = K_q^{\text{loop}}$. If $a = \Delta$ and $d(v_i) < \Delta$ for some i , then for any $\varepsilon > 0$ (11) gives

$$\left(\sum_{v \in V(H)} d(v)^{x-1+\varepsilon} \right)^{\frac{1}{x+\varepsilon}} < \Delta = \left(\sum_{v \in V(H)} d(v)^{x-1} \right)^{\frac{1}{x}}. \quad (13)$$

Finally, for any $\varepsilon > 0$, if $a < \Delta$ then $\Delta^\varepsilon a^x < \Delta^{x+\varepsilon}$, and so (11) gives

$$a < \Delta \quad \Longrightarrow \quad \left(\sum_{v \in V(H)} d(v)^{x-1+\varepsilon} \right)^{\frac{1}{x+\varepsilon}} < \Delta. \quad (14)$$

This already provides a substantial amount of information, fully analyzing the graphs H where $\sum_{v \in V(H)} d(v) \geq \Delta^2$ (here we focus on $x = 2$ and so the condition on H means $a = a(2, H) \geq \Delta$). Indeed, for $H \neq K_q^{\text{loop}}$, (12) and (13) applied at $x = 2$ imply

$$\left(\sum_{v \in V(H)} d(v)^{y-1} \right)^{\frac{1}{y}} < \left(\sum_{v \in V(H)} d(v) \right)^{\frac{1}{2}}$$

for any $y > 2$, which implies $\text{hom}(K_{1,x-1}, H)^{\frac{1}{x}} < \text{hom}(K_{1,1}, H)^{\frac{1}{2}}$ for any integer $x > 2$.

For the graphs H satisfying $\sum_{v \in V(H)} d(v) < \Delta^2$, we may already obtain a statement for large n (using (10) and also (14) at $x = 2$), but with an additional argument we can obtain a statement for all n . We need the following lemma, whose proof we give after first using the lemma to complete the proof of Theorem 1.5.

Lemma 3.1. *The function $\left(\sum_{v \in V(H)} d(v)^{x-1} \right)^{\frac{1}{x}}$ has at most one local maximum or minimum.*

If we assume Lemma 3.1, then (10), (12), (13), and (14) show that for $H \neq K_q^{\text{loop}}$ the function $\left(\sum_{v \in V(H)} d(v)^{x-1}\right)^{\frac{1}{x}}$ is either decreasing to Δ on $(2, \infty)$, increasing to Δ on $(2, \infty)$, or decreasing on $(2, x_0)$ and increasing to Δ on (x_0, ∞) for some $x_0 > 2$. See Figure 4 for the possible behaviors of $\left(\sum_{v \in V(H)} d(v)^{x-1}\right)^{\frac{1}{x}}$. So if $H \neq K_q^{\text{loop}}$, then (7) shows that Theorem 1.5 holds for any edge-min-critical $G \in \mathcal{G}(n, 1)$. This implies that the upper bounds given in Theorem 1.5 hold for any $G \in \mathcal{G}(n, 1)$.




Figure 4: The possible behaviors of the function $\left(\sum_{v \in V(H)} d(v)^{x-1}\right)^{\frac{1}{x}}$ for $H \neq K_{\Delta}^{\text{loop}}$.

Finally, we need to argue that for $H \neq K_q^{\text{loop}}$, the edge-min-critical graphs in $\mathcal{G}(n, 1)$ which achieve equality are the only possible graphs in $\mathcal{G}(n, 1)$ which achieve equality. It suffices to consider the addition of a single edge to one of the graphs achieving equality and showing that the number of H -colorings decreases in this case.

By considering the neighbors of a vertex $v \in V(H)$ with $d(v) = \Delta$, adding any edge to a disjoint union of stars strictly lowers the number of H -colorings unless H contains K_{Δ}^{loop} (a slight modification of the argument given in Case 1 of Section 2 will work, realizing that we need to consider both edges joining vertices in the same component and also edges joining vertices in different components). If H does contain K_{Δ}^{loop} and $H \neq K_{\Delta}^{\text{loop}}$, then H contains some other component and furthermore H must satisfy $\sum_{v \in V(H)} d(v) \geq \Delta^2$. Since $\frac{n}{2}K_2$ is the unique edge-min-critical graph achieving equality for this H and H has at least 2 components, adding any edge to $\frac{n}{2}K_2$ (which will necessarily join together two components of $\frac{n}{2}K_2$) will lower the number of H -colorings in this case as well. We leave the details to the reader. This completes the proof of Theorem 1.5.

Proof of Lemma 3.1: This lemma is a corollary of the following proposition about L^p norms, which is a special case of Lemma 1.11.5 in [18] (or, equivalently, Lemma 2 in Terence Tao's blog post 245C, Notes 1: Interpolation of L^p spaces). Recall that we assume H has no isolated vertices.

Proposition 3.2. *Define a measure μ on $V(H)$ by $\mu(v) = \frac{1}{d(v)}$, and let $g : V(H) \rightarrow \mathbb{R}$ be given by $g(v) = d(v)$. Then the function defined by $\frac{1}{x} \mapsto \|g\|_{L^x(V(H))} = \left(\sum_{v \in V(H)} d(v)^{x-1}\right)^{\frac{1}{x}}$ is log-convex for $x \in (2, \infty)$.*

Recall that a log-convex function can have at most one local maximum or local minimum. The composition of the reciprocal map and the map given in Proposition 3.2 is the function defined by $x \mapsto \left(\sum_{v \in V(H)} d(v)^{x-1}\right)^{\frac{1}{x}}$. Since the reciprocal is strictly monotone and therefore preserves local extremal values, Lemma 3.1 follows. \square

Lastly, we prove Theorem 1.9. Notice that we can still delete any edge from a graph $G \in \mathcal{G}(n, \delta, D)$ and remain in $\mathcal{G}(n, \delta, D)$ as long as the edge deletion does not lower the minimum degree. Therefore the proof of Theorem 1.5 also proves the inequality in Theorem 1.9 by restricting the function $\left(\sum_{v \in V(H)} d(v)^{x-1}\right)^{\frac{1}{x}}$ to values in $[2, D+1]$.

Having dealt with the inequality, we need to argue that the edge-min-critical graphs in $\mathcal{G}(n, 1, D)$ which achieve equality are the only possible graphs in $\mathcal{G}(n, 1, D)$ which achieve equality. This follows from similar arguments to those in the proof of Theorem 1.5 and we again leave the details to the reader.

4 Proof of Theorem 1.2

Recall that we will assume that $|V(H)| = q$, and we begin with a few remarks about the number of H -colorings of a cycle C_k . Let A denote the adjacency matrix of H . Then for $k \geq 3$, $\text{hom}(C_k, H) = \text{Tr } A^k$; indeed, the diagonal entry (ii) in A^k counts the number of H -colorings of the path on $k+1$ vertices P_{k+1} that color both endpoints with color i , and by identifying the endpoints we obtain a coloring of C_k with one fixed vertex having color i . Therefore if $\lambda_1, \lambda_2, \dots, \lambda_q$ are the eigenvalues of A , then

$$\text{hom}(C_k, H) = \lambda_1^k + \dots + \lambda_q^k. \quad (15)$$

It is possible to obtain results using ideas based on Proposition 3.2 (with some additional observations); we provide an alternate proof. Without loss of generality, assume that $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_q$. Notice that $\lambda_1 > 0$ and $\lambda_1 \geq |\lambda_q|$; this follows from the Perron-Frobenius theorem, but is also immediate as otherwise (15) would imply that $\text{hom}(C_k, H) < 0$ for large odd k .

We first address the inequality in Theorem 1.2, and deal with the cases of equality at the end. Suppose $k \geq 4$ is even and let $b = b(k, H) \geq \lambda_1$ be such that

$$\lambda_1^k + \lambda_2^k + \cdots + \lambda_q^k = b^k.$$

Then

$$\lambda_1^{k+2} + \cdots + \lambda_q^{k+2} \leq \lambda_1^2(\lambda_1^k + \cdots + \lambda_q^k) \leq b^2(\lambda_1^k + \cdots + \lambda_q^k) = b^{k+2},$$

with equality only for $b = \lambda_1$, so

$$(\lambda_1^{k+2} + \cdots + \lambda_q^{k+2})^{\frac{1}{k+2}} \leq (\lambda_1^k + \cdots + \lambda_q^k)^{\frac{1}{k}}, \quad (16)$$

which implies that $\text{hom}(C_k, H)^{\frac{1}{k}} \leq \text{hom}(C_4, H)^{\frac{1}{4}}$ for even $k \geq 6$.

Suppose next that $k \geq 5$ is odd, and so as above we have $\lambda_1^{k-1} + \cdots + \lambda_q^{k-1} = b^{k-1}$. Then

$$\lambda_1^k + \cdots + \lambda_q^k \leq |\lambda_1| \lambda_1^{k-1} + \cdots + |\lambda_q| \lambda_q^{k-1} \leq \lambda_1 b^{k-1} \leq b^k, \quad (17)$$

with equality only for $b = \lambda_1$, which implies that $\text{hom}(C_k, H)^{\frac{1}{k}} \leq \text{hom}(C_{k-1}, H)^{\frac{1}{k-1}}$.

Summarizing the above, the function $\text{hom}(C_k, H)^{\frac{1}{k}}$ is non-increasing from every even $k \geq 4$ to both $k+1$ and $k+2$ and so, for $k \geq 5$, $\text{hom}(C_k, H)^{\frac{1}{k}} \leq \text{hom}(C_4, H)^{\frac{1}{4}}$. Therefore for all $k \geq 3$,

$$\text{hom}(C_k, H)^{\frac{1}{k}} \leq \max\{\text{hom}(C_3, H)^{\frac{1}{3}}, \text{hom}(C_4, H)^{\frac{1}{4}}\}.$$

Now, if G is any 2-regular graph, then G is the disjoint union of cycles C_{k_i} . So if $\text{hom}(C_4, H)^{\frac{1}{4}} \geq \text{hom}(C_3, H)^{\frac{1}{3}}$,

$$\text{hom}(G, H) = \prod_i \text{hom}(C_{k_i}, H) \leq \prod_i \text{hom}(C_4, H)^{\frac{k_i}{4}} = \text{hom}(C_4, H)^{\frac{n}{4}},$$

with a similar statement holding if $\text{hom}(C_3, H)^{\frac{1}{3}} \geq \text{hom}(C_4, H)^{\frac{1}{4}}$.

Finally, we deal with the cases of equality. There is equality in (16) and (17) only when $b = \lambda_1$ (and so $\lambda_2 = \cdots = \lambda_q = 0$). Recall that A is symmetric and so has distinct eigenvectors associated to each λ_i , so in this case A has rank 1 and therefore all rows of A are scalar multiples of any other row. If any entry $A_{(ij)} = 0$, then some column and row of A is the 0 vector, which corresponds to an isolated vertex in H . Since we assume H has no isolated vertices, A must be the matrix of all 1's and so $H = K_q^{\text{loop}}$. Therefore, for $H \neq K_q^{\text{loop}}$ we have $\text{hom}(C_k, H)^{\frac{1}{k}} < \max\{\text{hom}(C_3, H)^{\frac{1}{3}}, \text{hom}(C_4, H)^{\frac{1}{4}}\}$ whenever $k \geq 5$. The statement about equality is now evident.

5 Proof of Theorem 1.6 ($\delta = 2$)

5.1 Preliminary remarks

We first gather together a number of observations that we will use in the proof. The following lemma is easily adapted from Lemma 3.2 in [6]. Recall that a graph with minimum degree δ is edge-min-critical if deleting any edge reduces the minimum degree of the resulting graph.

Lemma 5.1. *Let $G \in \mathcal{G}(n, 2)$ be an edge-min-critical graph. Either*

1. *G is a disjoint union of cycles, or*
2. *$V(G)$ may be partitioned into $Y_1 \cup Y_2$, with $1 \leq |Y_1| \leq n - 3$ in such a way that Y_1 induces a path, Y_2 induces a graph with minimum degree 2, each endpoint of the path induced by Y_1 has exactly one edge to Y_2 , the endpoints of these two edges to Y_2 are either the same or non-adjacent, and there are no other edges from Y_1 to Y_2 (see Figure 5).*

In the statement given in [6], the graph is assumed to be connected and *min-critical*, i.e. both edge-min-critical and *vertex-min-critical*. Being vertex-min-critical — meaning the deletion of any vertex reduces the minimum degree — is only used to show that $|Y_1| \geq 2$; the result holds for all edge-min-critical graphs by relaxing to $|Y_1| \geq 1$. To obtain the non-connected version, simply apply the connected version to each component. If some component is not a cycle, then augment the Y_2 obtained with the vertices in every other component.




Figure 5: A example of a path on 5 vertices ($|Y_1| = 5$) given in Lemma 5.1.

Corollary 5.2. *Let $G \in \mathcal{G}(n, 2)$ be an edge-min-critical graph. Then G may be constructed via the following iterative procedure:*

- *Start with a non-empty collection of disjoint cycles;*
- *Next, iteratively add a collection paths on $k \geq 2$ vertices which connect to existing vertices of the graph only at the endpoints of the path;*
- *Finally, add a collection of paths on 1 vertex (in other words, isolated vertices), each of which connects to exactly two distinct existing vertices of the graph, all at the same time.*

We emphasize that when we add a path on k vertices to a graph (via the construction in Corollary 5.2), we add k *new* vertices to the graph that induce a path in the augmented graph, so that the endpoints of the path are each adjacent to one existing vertex and no other vertices of the path are adjacent to existing vertices. Figure 5 may be viewed as an example of adding a path on 5 vertices (Y_1) to the graph on vertex set Y_2 .

Proof. By Lemma 5.1, paths on $k \geq 1$ vertices can be removed inductively until a collection of disjoint cycles remain, and so we may construct any graph G starting with the cycles. Reversing this, we may iteratively add paths on $k \geq 1$ vertices to produce G . The content of this corollary is that G may be constructed by adding all paths on $k \geq 2$ vertices before the paths on 1 vertex, and the paths on 1 vertex may all be added at the same time.

Why is this possible? Adding a path on 1 vertex creates a vertex of degree 2 adjacent to two vertices of degree at least 3. Since G is edge-min-critical, no future path on k vertices will be adjacent to the vertex of degree 2. \square

Lemma 5.1 is enough to prove the case when $\max\{\text{hom}(C_3, H)^{\frac{1}{3}}, \text{hom}(C_4, H)^{\frac{1}{4}}\} \geq \Delta$ (without a characterization of uniqueness); we will provide the details of this in the next section. A graph H which satisfies

$$\max\{\text{hom}(C_3, H)^{\frac{1}{3}}, \text{hom}(C_4, H)^{\frac{1}{4}}\} < \Delta \tag{18}$$

requires a few more observations.

Lemma 5.3. *For any two vertices u, v of H (not necessarily distinct), there are at most Δ^2 H -colorings of P_4 that map the initial vertex of the path to u and the terminal vertex to v . If H does not contain K_{Δ}^{loop} or $K_{\Delta, \Delta}$ as a component, then there are strictly less than Δ^2 such H -colorings.*

Proof. The first statement is obvious, since P_4 is connected and the maximum degree of H is Δ . Suppose there are Δ^2 extensions to an H -coloring of P_4 . Let P_4 have vertices w_1 through w_4 and edges $w_1 \sim w_2$, $w_2 \sim w_3$, and $w_3 \sim w_4$, and let w_1 and w_4 be given colors u and v in H , respectively. We color w_2 first (with color v_2) and then w_3 (with color v_3), conditioning on whether u is looped or not.

Suppose that u is unlooped in H . Clearly $d(u) = \Delta$ and each neighbor of u also has degree Δ . Since some of the Δ^2 extensions map w_3 to u , it must be the case that $v \sim_H u$. Furthermore, as w_2 maps to v_2 (so necessarily $v_2 \sim_H u$), every neighbor of v_2 must be adjacent to v . Since v_2 can be any neighbor of u , this implies that $K_{\Delta, \Delta}$ is the component of H containing u and v .

A similar analysis for looped u shows that K_{Δ}^{loop} is the component of H containing u and v . \square

Corollary 5.4. *Suppose that H satisfies (18) and let $k \geq 4$. For any two vertices u, v of H (not necessarily distinct), there are strictly fewer than Δ^{k-2} H -colorings of P_k that map the initial vertex of the path to u and the terminal vertex to v .*

Proof. Notice that $\text{hom}(C_4, K_{\Delta, \Delta})^{\frac{1}{4}} > \Delta$ and $\text{hom}(C_4, K_{\Delta}^{\text{loop}})^{\frac{1}{4}} = \Delta$. Color, beginning from one endpoint, until there are two uncolored vertices left. Then apply Lemma 5.3. \square

We can strengthen Corollary 5.4 when k is large.

Lemma 5.5. *Suppose that H satisfies (18). Then there exists a constant l_H (depending on H) such that if $k \geq l_H$ and the endpoints of P_k are mapped to H , then there are strictly fewer than $\frac{1}{|V(H)|^2} \Delta^{k-4}$ extensions to an H -coloring of P_k .*

Proof. Notice that a path must be mapped to a connected component of H ; focus on that component. If A is the adjacency matrix of that component, then the number of H -colorings of P_k with endpoints colored i and j is $A_{(ij)}^k$. If λ_1 denotes the largest eigenvalue of A , then by the Perron-Frobenius Theorem there exists a strictly positive vector \mathbf{x} such that $A^k \mathbf{x} = \lambda_1^k \mathbf{x}$ for all $k \geq 1$. By considering the row of A containing $\max_{i,j} A_{(ij)}^k$, we see that there is a constant c such that $\max_{i,j} A_{(ij)}^k \leq c \lambda_1^k$ (we can take $c = \max_j x_j / \min_j x_j$, where $\mathbf{x} = (x_j)$). Since $\lambda_1 \leq (\sum_i \lambda_i^4)^{\frac{1}{4}} = \text{hom}(C_4, H)^{\frac{1}{4}} < \Delta$ implies $\lambda_1 < \Delta$, this proves the lemma. \square

5.2 The proof

We are now ready to prove Theorem 1.6. We assume that G is edge-min-critical until we discuss the cases of equality in the upper bound. First, suppose that H satisfies

$$\max\{\text{hom}(C_3, H)^{\frac{1}{3}}, \text{hom}(C_4, H)^{\frac{1}{4}}\} \geq \Delta. \quad (19)$$

Using induction on n , we will show that for any $G \in \mathcal{G}(n, 2)$,

$$\text{hom}(G, H) \leq \max\{\text{hom}(C_3, H)^{\frac{n}{3}}, \text{hom}(C_4, H)^{\frac{n}{4}}\}.$$

The base case $n = 3$ is trivial.

For the inductive step, assume first that $\text{hom}(C_3, H)^{\frac{1}{3}} \leq \text{hom}(C_4, H)^{\frac{1}{4}}$. If all components of G are cycles, then we are finished by Theorem 1.2. If some component of G is not a cycle, then by Lemma 5.1 we can partition $V(G)$ into $Y_1 \cup Y_2$, with $1 \leq |Y_1| \leq n - 3$ and Y_1 connected to Y_2 . We imagine first coloring Y_2 and then extending this to Y_1 . By induction, there are at most $\text{hom}(C_4, H)^{\frac{n-|Y_1|}{4}}$ H -colorings of Y_2 . But since Y_1 is connected to Y_2 , for every fixed H -coloring of Y_2 , each vertex in Y_1 has at most Δ choices for a color. Therefore,

$$\text{hom}(G, H) \leq \Delta^{|Y_1|} \text{hom}(C_4, H)^{\frac{n-|Y_1|}{4}} \leq \text{hom}(C_4, H)^{\frac{n}{4}}. \quad (20)$$

The case when $\text{hom}(C_3, H)^{\frac{1}{3}} \geq \text{hom}(C_4, H)^{\frac{1}{4}}$ is similar.

With the upper bound established in this case, we turn to the cases of equality. First suppose that H satisfies $\max\{\text{hom}(C_3, H)^{\frac{1}{3}}, \text{hom}(C_4, H)^{\frac{1}{4}}\} > \Delta$. Then (20) is strict, which implies that equality can only be obtained for the disjoint union of cycles and hence Theorem 1.2 provides the cases of equality among edge-min-critical graphs.

Now suppose $\max\{\text{hom}(C_3, H)^{\frac{1}{3}}, \text{hom}(C_4, H)^{\frac{1}{4}}\} = \Delta$ (which implies that H cannot have $K_{\Delta, \Delta}$ as a component). Notice that equality is achieved for any G with $\text{hom}(G, H) = \Delta^n$, and suppose that $H \neq K_{\Delta}^{\text{loop}}$ (so since $\text{hom}(C_4, H)^{\frac{1}{4}} \leq \Delta$, H cannot contain K_{Δ}^{loop} as a component). By Theorem 1.2, the construction of G in Corollary 5.2 must start with disjoint copies of C_3 and/or C_4 . Corollary 5.4 implies that only paths on 1 vertex may be added to these cycles, and to achieve the bound of Δ^n , *every* coloring of these cycles must provide Δ choices for the color of the vertex in the path on 1 vertex. We outline the possible situations which occur when adding a path on 1 vertex to the cycles in Figure 5.2; the vertex labeled v must have Δ choices for a color regardless of how the adjacent cycles are colored.




Figure 6: The possible situations which occur when a path on 1 vertex (labeled v) is added to the cycles.

We will prove that in the situation of Case 5 from Figure 5.2, having Δ choices for the color on v for each coloring of C_4 forces H to contain K_{Δ}^{loop} as a component.

Suppose that the neighbors of v have colors i and j . We can assume that $i \neq j$, since C_4 can map its partition classes to the endpoints of any edge in H . Since v has Δ possibilities for its color, necessarily i and j must each have Δ neighbors and furthermore those Δ neighbors must be simultaneously neighbors of both i and j . In particular, since i and j are on adjacent vertices of C_4 , we have $i \sim_H j$ and so i and j must be possible colors for v . This means that i and j must be looped and that all neighbors of i are also neighbors of j . But if k is any other neighbor of i , then a similar argument (replacing j by k) shows that k is looped and is adjacent to all other neighbors of i . Therefore the component containing i is K_{Δ}^{loop} , which contradicts our assumption that K_{Δ}^{loop} is not a component of H .

A routine analysis of the other cases shows that having Δ choices for the color of the vertex in a path on 1 vertex always forces H to contain K_{Δ}^{loop} as a component. Therefore, equality can only occur when G is a disjoint union of cycles, so Theorem 1.2 again characterizes the cases of equality among edge-min-critical graphs.

Finally we need to show that edge-min-critical graphs are the only graphs achieving equality. Arguing as in Case 1 of Section 2, adding any edge to a C_4 will strictly lower the number of H -colorings unless H contains K_{Δ}^{loop} . The cases of adding an edge between two disjoint cycles is similar, and so adding any edge to a graph achieving equality will strictly lower the number of H -colorings unless H is the disjoint union of some number of fully looped complete graphs. If H is of this form and $H \neq K_{\Delta}^{\text{loop}}$, then $\text{hom}(C_3, H)^{\frac{1}{3}} > \text{hom}(C_4, H)^{\frac{1}{4}}$, and so in fact adding any edge to $\frac{n}{3}C_3$ will strictly lower the number of H -colorings since the disjoint copies of C_3 can be colored using different components of H , but the copies of C_3 joined by an edge must all be colored by a single component of H .

Now suppose that H satisfies

$$\max\{\text{hom}(C_3, H)^{\frac{1}{3}}, \text{hom}(C_4, H)^{\frac{1}{4}}\} < \Delta. \quad (21)$$

Recall that from Theorem 1.2 we have

$$\text{hom}(C_k, H)^{1/k} \leq \max\{\text{hom}(C_3, H)^{1/3}, \text{hom}(C_4, H)^{1/4}\},$$

for all $k \geq 3$, which we bound (for simplicity) by

$$\text{hom}(C_k, H) \leq (\Delta^4 - 1)^{k/4} \quad \text{for } k \geq 3. \quad (22)$$

As in the proof of Theorem 1.7, we will let $S(2, H)$ denotes the vectors in $V(H)^2$ with the property that the elements of the vector have Δ common neighbors, and $s(2, H) = |S(2, H)|$. Notice that $\text{hom}(K_{2, n-2}) \geq s(2, H)\Delta^{n-2}$.

Suppose first that G is edge-min-critical. We will utilize the construction of G from Corollary 5.2 to produce all H -colorings of G by coloring the disjoint cycles first and then coloring the paths.

If there are more than t vertices in the disjoint cycles, then by (22) we have $\text{hom}(G, H) \leq (\Delta^4 - 1)^{t/4} \Delta^{n-t}$. Therefore, we may assume that there are at most c_1 vertices in disjoint cycles. (All constants in the remainder of this proof will depend on H but will be independent of n .)

After coloring the cycles, we look at the paths that are added iteratively. If any path has length longer than some constant l , then by Lemma 5.5 we have $\text{hom}(G, H) < \Delta^{n-2}$. Since a path of length k , for $2 \leq k \leq l$, has at most $\Delta^k - 1$ extensions to an H -coloring by Corollary 5.4, if there are at least c_2 such paths then

$$\text{hom}(G, H) < \left(\prod_{i=1}^{c_2} (\Delta^{k_i} - 1) \right) \Delta^{n - \sum_{i=1}^{c_2} k_i} < \Delta^{n-2}.$$

So, we may assume that the decomposition of G from Corollary 5.2 has fewer than c_3 vertices in either disjoint cycles or paths on k vertices, with $2 \leq k \leq l$, and no paths on k vertices, with $k > l$. Therefore, the decomposition has at least $n - c_3$ vertices in paths on 1 vertex. Furthermore, each path on 1 vertex must be attached to two of the at most c_3 vertices composing the disjoint cycles and the paths on at least 2 vertices that are added. By the pigeonhole principle there exists a $c_4 > 0$ and two vertices in G with at least $c_4 n$ paths on 1 vertex joining them.

We have shown that every edge-min-critical graph G which does not have two vertices with at least $c_4 n$ paths on 1 vertex joining them has $\text{hom}(G, H) < \Delta^{n-2}$, and by Lemma 5.5 we have the same bound on $\text{hom}(G, H)$ if G has a path on k vertices when $k > l$. We now deal with the remaining edge-min-critical graphs G .

Let w_1 and w_2 denote the vertices in G joined by at least $c_4 n$ paths on 1 vertex. Recall from Section 2 that $S(2, H)$ is the set of vectors in $V(H)^2$ with the property that the elements of the vector have Δ common neighbors, and $s(2, H) = |S(2, H)|$. Suppose first that the colors on w_1 and w_2 are an element of $S(2, H)$. If G is different from $K_{2, n-2}$, then w_1 , w_2 , and the at least $c_4 n$ paths on 1 vertex between them do not form all of G . But then G must contain either a cycle which does not include w_1 or w_2 , or a path on k vertices (for $2 \leq k \leq l_H$) from w_i to w_j for some $i, j \in \{1, 2\}$. By first coloring w_1 and w_2 , then any remaining disjoint cycles, and finally the remaining vertices, Corollary 5.4 and (22) imply that there exists a $c_5 < 1$ such that there are at most

$$s(2, H) c_5 \Delta^{n-2} \tag{23}$$

H -colorings of G of this type.

Now suppose that the colors on w_1 and w_2 are not an element of $S(2, H)$. By first coloring w_1 and w_2 , then any remaining disjoint cycles, and finally the remaining vertices, we have at most

$$|V(H)|^2 \Delta^{n-c_4 n-2} (\Delta - 1)^{c_4 n} \tag{24}$$

H -colorings of G of this type.

Combining (23) and (24) gives

$$\text{hom}(G, H) \leq |V(H)|^2 \Delta^{n-2} \left(\frac{\Delta - 1}{\Delta} \right)^{c_4 n} + s(2, H) c_5 \Delta^{n-2} < s(2, H) \Delta^{n-2},$$

with the last inequality holding for large enough n .

We have shown that the only edge-min-critical graph which achieves equality is $K_{2, n-2}$. Arguing as in Case 1 of Section 2 shows that adding any edges to $K_{2, n-2}$ produces a graph G with $\text{hom}(G, H) < \text{hom}(K_{2, n-2}, H)$, which completes the proof.

6 Concluding Remarks

We now briefly describe how to generalize the results in this paper to weighted H -colorings. Suppose that $\Lambda = \{\lambda_v : v \in V(H)\}$ is a set of positive weights indexed by $V(H)$. Let

$$d_\Lambda(v) = \sum_{w \sim_H v} \lambda_w$$

(noting that a loop on v will include one λ_v term in the sum) and

$$\Delta_\Lambda = \max_{v \in V(H)} d_\Lambda(v).$$

Furthermore, for $i, j \in V(H)$, let A_Λ be the matrix with ij entry $\sqrt{\lambda_i \lambda_j} \mathbf{1}_{i \sim_H j}$. Then the results of Theorems 1.2, 1.5, 1.6, and 1.7 hold for weighted H -colorings by changing $d(v)$ to $d_\Lambda(v)$, Δ to Δ_Λ , A to A_Λ , and $\text{hom}(G, H)$ to $Z_\Lambda(G, H)$. For example, the condition $\sum_{v \in V(H)} d(v) < \Delta^2$ is replaced by $\sum_{v \in V(H)} \lambda_v d_\Lambda(v) < (\Delta_\Lambda)^2$. The generalization is straightforward and we leave the details to the interested reader.

The proofs of Theorems 1.5 and 1.6 depend heavily on analyzing the structure of edge-min-critical graphs. For $\delta \geq 3$, there is no helpful structural characterization of these graphs. A nice answer to the following question might help prove Conjecture 1.4 for other values of δ .

Question 6.1. *What can be said about the structure of edge-min-critical graphs in $\mathcal{G}(n, \delta)$ for $\delta \geq 3$?*

In light of Theorem 1.9, it would be interesting to consider a maximum degree condition D in addition to a minimal degree condition δ when $\delta > 1$. Again writing $\mathcal{G}(n, \delta, D)$ for the set of graphs with minimum degree δ and maximum degree at most D , if $D < n - \delta$ then $K_{\delta, n-\delta} \notin \mathcal{G}(n, \delta, D)$. It is possible that, as in the $\delta = 1$ case, the graph $K_{\delta, n-\delta}$ is replaced by $\frac{n}{\delta+D} K_{\delta, D}$ in $\mathcal{G}(n, \delta, D)$. This leads to the following natural question.

Question 6.2. Fix H . Is it true that for any $G \in \mathcal{G}(n, \delta, D)$

$$\text{hom}(G, H) \leq \max\{\text{hom}(K_{\delta+1}, H)^{\frac{n}{\delta+1}}, \text{hom}(K_{\delta,\delta}, H)^{\frac{n}{2\delta}}, \text{hom}(K_{\delta,D}, H)^{\frac{n}{\delta+D}}\}?$$

Acknowledgement: The author is grateful to Igor Rivin for suggesting the use of Proposition 3.2 and to David Galvin for a number of helpful discussions.

References

- [1] J. Alexander, J. Cutler, and T. Mink, *Independent sets in graphs with given minimum degree*, Electron. J. Combin. **19(3)** (2012), #P37.
- [2] J. Cutler and A. J. Radcliffe, *Extremal graphs for homomorphisms*, J. Graph Theory **67** (2011), 261-284.
- [3] ———, *Extremal graphs for homomorphisms II*, to appear in J. Graph Theory, DOI: 10.1002/jgt.21749.
- [4] ———, *The maximum number of complete graphs in a graph given maximum degree*, arXiv:1306.1803.
- [5] J. Engbers and D. Galvin, *H-coloring bipartite graphs*, J. Combin. Theory Ser. B **102** (2012), 726-742.
- [6] ———, *Counting independent sets of a fixed size in graphs with given minimum degree*, Manuscript.
- [7] D. Galvin, *Two problems on independent sets in graphs*, Discrete Math. **311** (2011), 2105-2112.
- [8] ———, *Maximizing H-colorings of regular graphs*, J. Graph Theory **73** (2013), 66-84.
- [9] ———, *Counting colorings of a regular graph*, arXiv:1205.2718.
- [10] D. Galvin, F. Martinelli, K. Ramanan, and P. Tetali, *The multi-state hard core model on a regular tree*, SIAM J. of Disc. Math **25** (2011), 894-916.
- [11] D. Galvin and P. Tetali, *On weighted graph homomorphisms*, Graphs, Morphisms, and Statistical Physics, DIMACS Ser. in Discrete Math. Theoret. Comput. Sci. **63** (2004), 97-104.
- [12] D. Galvin and Y. Zhao, *The number of independent sets in a graph with small maximum degree*, Graphs Combin. **27** (2011), 177-186.
- [13] J. Kahn, *An entropy approach to the hard-core model on bipartite graphs*, Combin. Probab. and Comput. **10** (2001), 219-237.
- [14] ———, *Entropy, independent sets and antichains: a new approach to Dedekind's problem*, Proc. Amer. Math. Soc. **130** (2002), no. 2, 371-378.
- [15] F. Lazebnik, *On the greatest number of 2 and 3 colorings of a (v, e) -graph*, J. Graph Theory **13** (1989), no. 2, 203-214.
- [16] P.-S. Loh, O. Pikhurko, and B. Sudakov, *Maximizing the number of q-colorings*, Proc. Lon. Math. Soc. **101** (2010), 655-696.
- [17] K. Ramanan, A. Sengupta, I. Ziedins, and P. Mitra, *Markov random field models of multicasting in tree networks*, Adv. in Appl. Probab. **34** (2002), 58-84.

- [18] T. Tao, *An epsilon of room, I: real analysis: pages from year three of a mathematical blog*, Graduate studies in mathematics, vol. 117, American Mathematical Society, Providence, RI, 2009.
- [19] Y. Zhao, *The number of independent sets in a regular graph*, *Combin. Probab. and Comput.* **19** (2010), 315-320.
- [20] ———, *The bipartite swapping trick on graph homomorphisms*, *SIAM J. Disc. Math* **25** (2011), 660-680.