

Marquette University

e-Publications@Marquette

Biomedical Sciences Faculty Research and
Publications

Biomedical Sciences, Department of

9-1-1993

Evidence for a Second Type of Fibril Branch Point in Fibrin Polymer Networks, the Trimolecular Junction

Michael W. Mosesson
Blood Center of Wisconsin

James P. DiOrio
Sinai Samaritan Medical Center

Kevin R. Siebenlist
Marquette University, kevin.siebenlist@marquette.edu

Joseph S. Wall
Sinai Samaritan Medical Center

James F. Hainfeld
University of Wisconsin Medical School

Follow this and additional works at: https://epublications.marquette.edu/biomedsci_fac



Part of the [Neurosciences Commons](#)

Recommended Citation

Mosesson, Michael W.; DiOrio, James P.; Siebenlist, Kevin R.; Wall, Joseph S.; and Hainfeld, James F., "Evidence for a Second Type of Fibril Branch Point in Fibrin Polymer Networks, the Trimolecular Junction" (1993). *Biomedical Sciences Faculty Research and Publications*. 233.
https://epublications.marquette.edu/biomedsci_fac/233

Marquette University

e-Publications@Marquette

Biomedical Sciences Faculty Research and Publications/College of Health Sciences

This paper is NOT THE PUBLISHED VERSION.

Access the published version via the link in the citation below.

Blood, Vol. 82, No. 5 (September 1993): 1517-1521. [DOI](#). This article is © American Society of Hematology and permission has been granted for this version to appear in [e-Publications@Marquette](#). American Society of Hematology does not grant permission for this article to be further copied/distributed or hosted elsewhere without express permission from American Society of Hematology.

Evidence for a Second Type of Fibril Branch Point in Fibrin Polymer Networks, the Trimolecular Junction

Michael W. Mosesson

Sinai Samaritan Medical Center, University of Wisconsin Medical School, Milwaukee

James P. DiOrio

Sinai Samaritan Medical Center, University of Wisconsin Medical School, Milwaukee

Kevin R. Siebenlist

Sinai Samaritan Medical Center, University of Wisconsin Medical School, Milwaukee

Joseph S. Wall

Sinai Samaritan Medical Center, University of Wisconsin Medical School, Milwaukee

James F. Hainfeld

Sinai Samaritan Medical Center, University of Wisconsin Medical School, Milwaukee

Abstract

Fibrin molecules polymerize to double-stranded fibrils by intermolecular end-to-middle domain pairing of complementary polymerization sites, accompanied by fibril branching to form a clot network. Mass/length measurements on scanning transmission electron microscopic images of fibrils comprising branch points showed two types of junctions. Tetramolecular junctions occur when two fibrils converge, creating a third branch with twice the mass/length of its constituents. Newly recognized trimolecular junctions have three fibril branches of equal mass/length, and occur when an extraneous fibrin molecule initiates branching in a propagating fibril by bridging across two unpaired complementary polymerization sites. When trimolecular junctions predominate, clots exhibit nearly perfect elasticity.

FBRINOGEN MOLECULES are symmetrical, elongated (46 ± 2 nm) trinodular structures¹ consisting of a central disulfide-bonded domain (the "E" domain) containing the amino terminal regions of its three pairs of chains ($A\alpha$, $B\beta$, γ), connected through coiled-coil regions to each of its two outer "D" domains.^{2,3} Fibrin polymerization begins after thrombin cleavage of fibrinopeptide A from the amino terminus of the two $A\alpha$ chains of fibrinogen molecules, exposing two "A" polymerization sites in the central E domain.^{4,5,6} Each "A" site then interacts with an available complementary site, "a," in the outer D domain of another fibrin molecule,^{7,8,9} forming double-stranded fibrils composed of a half-staggered overlapping arrangement of molecules^{10,11,12,13,14,15,16,17} (Fig 1). Under physiological conditions, fibrils undergo branching and extensive lateral association, producing a predominantly thick fiber clot network^{15,18,19,20,21} that provides a structural framework for the intravascular thrombus.




Fig 1.. Schematic diagram illustrating the half-staggered arrangement of fibrin molecules (elongated bars) to form double-stranded fibrils. Branching occurs at tetramolecular or trimolecular junctions as described in the text.

Two extreme forms of fibrin polymer networks are known, namely, fine and coarse.²² Fine clots consist of a branching thin fibril network, are produced at high ionic strength ($I \geq 0.4$) and $\text{pH} > 8$,^{16,17} and exhibit nearly perfect elasticity.^{23,24} Coarse clots form readily at physiological ionic strength ($I = 0.15$) and $\text{pH} 7.4$, and consist mainly of a branching thick fiber network.^{15,18,19,20,21,22,25,26,27} However, intermediate structures containing fine clot networks are observed under coarse clot buffer conditions

depending on fibrin concentration and/or the kinetics of fibrin assembly,^{17, 18, 19, 28} and are favored by the presence in the clotting milieu of certain protein components of blood such as circulating catabolic fibrinogen derivatives (eg, fraction I-9) lacking car-boxy terminal segments of A α chains,^{29, 30, 31, 32, 33} IgG, thrombospondin, histidine-rich glycoprotein, or albumin.^{34, 35, 36, 37}

Fibrin branching in fine or coarse polymer networks has long been assumed to occur solely by the process of convergence and lateral association of fibrils to form branch points.^{15, 18, 19, 38, 39, 40} We term these well-recognized junctions “tetramolecular branch points” to distinguish them from a newly proposed type of fibril network junction, the “trimolecular branch point.”²⁸ Evidence for two types of branch junctions is currently based on fibril width measurements on low-resolution transmission electron microscopic (TEM) images of critical point dried fibrin networks.²⁸ Tetramolecular junctions, which form by coalescence of two double-stranded fibrils, have a third four-stranded fibril twice the width of its constituents (Fig 1). In contrast, all three fibrils contributing to a trimolecular junction are of equal width.²⁸

Although the morphometric evidence for a trimolecular junction is attractive, its existence as a distinct type of branch point is not intuitively evident, and this junction has not, even in recent reports,⁴⁰ been considered as a significant matrix element for modelling fibrin clot assembly. Therefore, to provide persuasive proof of its existence and at the same time confirm the mechanism of its formation, we made mass measurements on clot network structures that had been imaged by high-resolution scanning transmission electron microscope (STEM).^{1, 41, 42}

MATERIALS AND METHODS

Formation of fine clots.

Human plasma fibrinogen fractions I-2 and I-9 were prepared as previously described.^{29, 32} Clots for TEM examination were formed as 50- μ L droplets on Parafilm (Amer. Con Co, Greenwich, CT) from fibrinogen fraction I-2 (25 μ g/mL) in 400 mmol/L NaCl, 10 mmol/L HEPES, 5 mmol/L EDTA, pH 8.5 buffer (I = 0.4), or from fibrinogen fraction I-9 (25 μ g/mL) in 290 mmol/L NaCl, 10 mmol/L HEPES, pH 7.0 buffer (I = 0.3), at an α -thrombin concentration of 1 U/mL, and incubated for 2 hours at room temperature. Grid specimens were picked up on ultrathin carbon films and washed with 0.15 mol/L ammonium acetate solution. The specimens were subjected to progressive alcohol dehydration, Freon treatment, drying over a vapor barrier of Freon, and rotary shadowing with platinum-carbon, and were then examined in a Philips 400 electron microscope (Philips Instruments Co, Roselle, IL).

For STEM experiments, fibrinogen fraction I-2 or fraction I-9 solutions were dialyzed against 400 mmol/L NaCl, 10 mmol/L HEPES, 5 mmol/L EDTA, or 1 mmol/L CaCl₂, pH 8.5 buffer (I = 0.4), deposited as 50- μ L droplets (16 to 50 μ g/mL) on Parafilm, and converted to fibrin at an α -thrombin concentration of 1 U/mL. After incubation of clotting mixtures at room temperature for 60 to 150 minutes, grid specimens were obtained by passing the fibrin droplet back and forth several times through a pipette before injecting a sample into a droplet of buffer on the grid surface or by passing the grid directly through the fibrin droplet. After 1 minute, the specimen on the grid was washed 8 to 10 times with 50 mmol/L ammonium acetate solution, quickly frozen in liquid nitrogen, freeze-dried, transferred under vacuum to the microscope stage, and imaged by using a 40-kV probe focussed at 0.25 nm.

Mass measurements on fibrin fibrils.

Mass determinations on STEM images were based on electron scattering measurements and were performed off-line as previously described.^{1, 41, 42} A “fiber” program was used for determining mass/length of fibrils, a “circle” program for estimating the mass of trimolecular complexes composed of two D and one E domain (D · E · D) complexes within fibrils, or a “user-defined boundary” program for measuring the mass at a fibril branch junction.^{41, 42} Tobacco mosaic virus particles were used as a mass calibration standard.^{43, 44}

The predicted mass/length of fibrils.

Plasma subfractions containing mostly intact fibrinogen molecules (ie, fraction I-2) have a mean molecular weight of $340 \pm 15 \times 10^3$, whereas subfractions containing molecules from which COOH-terminal portions of A α chains have been cleaved (eg, fraction I-9) have a mean molecular weight of $300 \pm 15 \times 10^3$.¹ The mean molecular weight of fraction I-2 fibrin molecules is 334×10^3 , about 2% lower than the precursor fibrinogen due to loss of both fibrinopeptides; that of fraction I-9 fibrin is 294×10^3 . The predicted mass/length of double- and four-stranded fibrin fibrils is calculated using the values for fibrin monomer and an assumed length of 46 nm for a fibrin molecule.¹

RESULTS AND DISCUSSION

Rotary-shadowed TEM images of fine clot networks formed from intact fibrinogen molecules (fraction I-2) or from fibrinogen molecules lacking carboxy terminal regions of A α chains (fraction I-9) (not shown) showed a highly branched fibril network (Fig 2). Most branch junctions, as judged from fibril widths, were trimolecular branch points (arrows), although tetramolecular branch points were also found (double-shaft arrows) at a lower frequency. Thus, these observations indicate that trimolecular branches are the predominant junctional components in fine clot matrices. Multi-stranded fibers were found commonly (Fig 2B), although they were not as thick or as predominant as in coarse clots. These fibers often possessed protruding thin fibrils forming an arm of a branch junction (arrowheads), but because the fibers were multistranded, the origin of such fibrils as an element of either a trimolecular or tetramolecular branch point could not be ascertained.




Fig 2.. TEM images of fibrin. Arrows indicate fibril junctions having branches of the same width (trimolecular junctions); double-shaft arrows indicate tetramolecular junctions. Arrowheads indicate fibrils protruding from a fiber bundle. Bar = 100 nm.

STEM images of unstained freeze-dried fine fibrin network fibers showed many fibrils and frequent branches (Fig 3). Fibrils with widths corresponding to double-stranded fibrils that had been formed from intact fibrinogen molecules had mass/length values of 14.4 Kd/nm (EDTA-containing buffer) and 14.8 Kd/nm (CaCl₂-containing buffer) (Table 1), respectively, both values corresponding closely to the predicted value of 14.5 Kd/nm. As had been observed in rotary-shadowed images, these fibrils were characterized by the presence of oblong to triangular dense structures, 8 to 16 nm across and 19 ± 2.7 nm in length along long fibril axes, regularly spaced ~22.5 nm apart; this interval corresponds to one-half the length of a fibrinogen molecule, and reflects the half-staggered arrangement of fibrin molecules within the fibril. The mass of these structures (eg, the circle in Fig 3A), was 300 ± 38 Kd (integration radius [r_i] = 12 nm; n = 48), a value corresponding to a D · E · D complex.¹ The apices of consecutive triangular-shaped D · E · D complexes commonly were reversed along the fibril (connected arrows, Fig 3A, B, and D), consistent with the fibril twisting that is known to take place.^{16, 17, 28} A schematic drawing of these fibrils is presented in Fig 4.




Fig 3.. STEM images of fibrils and fibril junctions. Fibrin was formed from fibrinogen fraction 1-2 solutions in 400 mmol/L NaCl, 10 mmol/L HEPES, 1 mmol/L CaCl₂, pH 8.5 buffer (I = 0.4) (A through C), or in 400 mmol/L NaCl, 10 mmol/L HEPES, 5 mmol/L EDTA, pH 8.5 buffer (I = 0.4) (D and E). A “fiber” program (eg, rectangle in A) was used for determining fibril mass/length, a “circle” program (eg, circle in A) was used for estimating the mass of D · E · D complexes within fibrils, or a “user-defined boundary” program (eg, irregular enclosed shape in E) was used for measuring the mass at a junctional nidus.^{20, 21} Bar = 100 nm.

Table 1. Fibril Mass/Length

Buffer Condition	Empty Cell	Fibrinogen Preparation	Double-Stranded Fibrils	Trimolecular Branch Fibrils	Four-Stranded Fibrils
0.4 mol/L NaCl, 10 mmol/L HEPES, pH 8.5 (I = 0.4)	5 mmol/L EDTA,	Fraction I-2*	14.4 ± 1.5	14.9 ± 1.2	26.4 ± 2.3
			(n 73)	(n = 12)	(n = 8)

0.4 mol/L NaCl, 10 mmol/L HEPES, pH 8.5 (I = 0.4)	1 mmol/L CaCl ₂ ,	Fraction I-2*	14.8 ± 2.1 (n = 38)	14.8 ± 2.3 (n = 15)	29.9 ± 4.2 (n = 8)
0.29 mol/L NaCl, 10 mmol/L HEPES, pH 7.0 (I = 0.3)		Fraction I-9†	13.8 ± 1.6 (n = 112)	13.1 ± 1.4 (n = 14)	24.2 ± 3.4 (n = 8)

Values are in kilodaltons per nanometer.

*Predicted mass/length for double-stranded fibril, 14.5 Kd/nm; four-stranded fibril, 29.0 Kd/nm.

†Predicted mass/length for double-stranded fibril, 12.8 Kd/nm; four-stranded fibril, 25.6 Kd/nm.




Fig 4.. An illustrated model of fibrin fibrils and fibril branch junctions. Fibrin molecules are represented by a small black oval (central E domain) and two larger ovals (outer D domains) that are connected through the respective E domain by either a solid or dashed line. The D domains of each molecule have the same fill pattern. Fibril assembly occurs by end-to-middle domain interactions between complementary polymerization sites in the E (“A” site) and D domains (“a” site), forming double-stranded, twisting fibrils. Branching occurs at tetramolecular (1) or trimolecular junctions (2). The three D · E · D complexes comprising the nidus of a trimolecular junction are enclosed by a dashed circle. An incipient trimolecular fibril junction is depicted in the lower part of the diagram.

Mass measurements at branch junctions containing a fibril with twice the width of its two constituent thin fibrils indicated that the wider fibril also had twice the mass/length, thus identifying them as tetramolecular junctions (double-shafted arrows, Fig 3A and D). Similarly, junctions consisting of three thin fibrils of equal width and equal mass/length (Table 1) could be unambiguously identified by either criterion as trimolecular junctions (arrows, Fig 3B, C, E). In addition, the mass at the nidus of trimolecular junctions (eg, irregular shape, Fig 3E) was 898 ± 139 Kd ($n = 7$), a value corresponding to the three D · E · D complexes that constitute such a junction (cf, Fig 4). Taken together, these observations indicate that trimolecular branches are initiated by extension of the D domain of a fibrin molecule away from a propagating fibril at an incipient branch junction, concomitant with bridging across the unpaired complementary polymerization sites by an extraneous fibrin molecule, as originally suggested somewhat less concisely.²⁸ To our knowledge, this mechanism of branching has not yet been observed in other types of biologic and synthetic polymer networks.⁴⁵ It is also interesting to note that this type of branching was recently proposed as a feature of early fibrin polymerization, in the context of an intermediate mechanism for subsequent lateral fibril associations or for unordered fibrin aggregation.⁴⁶

Fibrin from fibrinogen fraction I-9 molecules have a diminished tendency to form thick fibers,^{29, 30} consistent with evidence suggesting that a second polymerization site ("b") contributing to lateral fibril association is located in the car-boxy-terminal segments of A α chains.³³ This fibrin formed a fine matrix consisting predominantly of trimolecular branch structures (not shown), but at a lower pH (7.0) and ionic strength (0.3) than did intact fibrin. STEM mass measurements of fibril D · E · D nodules yielded a value of 280 \pm 39 Kd (r_f = 12 nm; n = 39). The lower mass for these complexes, compared with those in fraction I-2 fibrin ($P < .02$), corresponds to the lower mass found in the vicinity of the E domains of fraction I-9 molecules due to absence of car-boxy-terminal portions of A α chains.¹ Mean thin fibril widths were the same as for fraction I-2 fibrin, as was the 22.5 nm spacing of D · E · D nodules. Their mass/length value (13.8 Kd/nm) was lower than that of intact fibrin fibrils ($P < .02$) and close to the predicted value of 12.8 Kd/nm for this type of fibrin. Similarly, four-stranded fibrils had mass/length values twice that of double-stranded fibrils (Table 1).

The finding that each of the three fibrils constituting trimolecular branch points was double-stranded by mass/length criteria, corroborates previous conclusions based on fibril width measurements,²⁸ provides proof of the existence of these junctions as structural elements of the fibrin network, and is a means for distinguishing them unambiguously from tetramolecular junctions. In view of the close correspondence between fibril widths and mass/length measurements, we conclude that fibril width measurements alone are sufficient for identifying fibril branch junctions. Using this criterion, we note that negatively contrasted TEM images reported by Hantgan et al^{18, 19} and critical point dried TEM images reported by Müller et al¹⁶ and Mosesson et al¹⁷ present a number of excellent examples of trimolecular branch junctions formed under coarse or fine clot buffer conditions. In contrast, the number of thin fibrils comprising multistranded fibers can be accurately determined only by STEM mass measurements, because as yet there is no simple correlation between fiber width and the number of constituent fibrils.

It is likely that trimolecular branch junctions play an important role in determining the rheologic properties of fibrin clots that help to define their physiologic function in relation to other clot elements. Lateral fibril associations vastly predominate in coarse clots in which thick fiber bundles are the rule, a composition that evidently accounts for the fiber or junctional slippage and realignments that occur in coarse clot networks under stress conditions leading to irrecoverable deformation.^{23, 24} On the other hand, trimolecular branch points predominate in fine clot networks, which exhibit nearly perfect elasticity,²³ a property that we attribute to their presence. Their contributions to the properties of coarse clot networks in relation to the other elements of network structure remain to be explored.

ACKNOWLEDGMENT

We thank William Semrad and Karen Mickey Higgins for graphics/photography services.

REFERENCES

- 1 Mosesson MW, Hainfeld JF, Haschemeyer R, Wall JS. **Identification and mass analysis of human fibrinogen molecules and their domains by scanning transmission electron microscopy.** *J Mol Biol*, 153 (1981), p. 695
- 2 Doolittle RF. **The structure and evolution of vertebrate fibrinogen.** *Ann NY Acad Sci*, 408 (1983), p. 13
- 3 Henschen A, Lottspeich F, Kehl M, Southan C. **Covalent structure of fibrinogen.** *Ann NY Acad Sci*, 408 (1983), p. 28
- 4 Scheraga HA, Laskowski M Jr. **The fibrinogen-fibrin conversion.** *Adv Protein Chem*, 12 (1957), p. 1

- 5 Doolittle RF. **Structural aspects of the fibrinogen to fibrin conversion.** *Adv Protein Chem*, 27 (1973), p. 1
- 6 Kudryk B, Collen D, Woods KR, Blombäck B. **Evidence for localization of polymerization sites in fibrinogen.** *J Biol Chem*, 249 (1974), p. 3322
- 7 Kudryk B, Reuterby J, Blombäck B. **Adsorption of plasmic fragment D to thrombin modified fibrinogen-sepharose.** *Thromb Res*, 2 (1973), p. 297
- 8 Olexa SA, Budzynski AZ. **Evidence for four different polymerization sites involved in human fibrin formation.** *Proc Natl Acad Sci USA*, 77 (1980), p. 1374
- 9 Blombäck B, Hessel B, Hogg D, Therkildsen L. **A two-step fibrinogen-fibrin transition in blood coagulation.** *Nature*, 257 (1978), p. 501
- 10 Ferry JD. **The mechanism of polymerization of fibrinogen.** *Proc Natl Acad Sci USA*, 38 (1952), p. 566
- 11 Stryer L, Cohen C, Langridge R. **Axial period of fibrinogen and fibrin.** *Nature*, 197 (1963), p. 793
- 12 Krakow W, Endres GF, Siegel BM, Scheraga HA. **An electron microscopic investigation of the polymerization of bovine fibrin monomer.** *J Mol Biol*, 71 (1972), p. 95
- 13 Fowler WE, Hantgan RR, Hermans J, Erickson HP. **Structure of the fibrin protofibril.** *Proc Natl Acad Sci USA*, 78 (1981), p. 4872
- 14 Williams RC. **Morphology of fibrinogen monomers and of fibrin polymers.** *Ann NY Acad Sci*, 408 (1983), p. 180
- 15 Hantgan R, Hermans J. **Assembly of fibrin: A light scattering study.** *J Biol Chem*, 254 (1979), p. 11272
- 16 Müller MF, Ris H, Ferry JD. **Electron microscopy of fine fibrin clots and coarse fibrin films.** *J Mol Biol*, 174 (1984), p. 369
- 17 Mosesson MW, DiOrio JP, Müller MF, Shainoff JR, Siebenlist KR, Amrani DL, Homandberg GA, Soria J, Soria C, Samama M. **Studies on the ultrastructure of fibrin lacking fibrinopeptide B (β -fibrin).** *Blood*, 69 (1987), p. 1073
- 18 Hantgan R, Fowler W, Erickson H, Hermans J. **Fibrin assembly: A comparison of electron microscopic and light scattering results.** *Thromb Haemost*, 44 (1980), p. 119
- 19 Hantgan RR, McDonagh J, Hermans J. **Fibrin assembly.** *Ann NY Acad Sci*, 408 (1983), p. 344
- 20 Hewat EA, Tranqui L, Wade RH. **Electron microscope structural study of modified fibrin and a related modified fibrinogen aggregate.** *J Mol Biol*, 170 (1983), p. 203
- 21 Carr MJ Jr, Shen LL, Hermans J. **Mass-length ratio of fibrin fibres from gel permeation and light scattering.** *Biopolymers*, 17 (1977), p. 1
- 22 Ferry JD, Morrison PR. **Preparation and properties of serum and plasma proteins. VIII. The conversion of human fibrinogen to fibrin under various conditions.** *J Am Chem Soc*, 69 (1947), p. 388
- 23 Ferry JD. **Structure and rheology of fibrin networks.** Kramer O (Ed.), *Biological and Synthetic Polymer Networks*, Elsevier, New York, NY (1988), p. 41
- 24 Bale MD, Müller MF, Ferry JD. **Rheological studies of creep and creep recovery of unligated fibrin clots: Comparison of clots prepared with thrombin and anicrod.** *Biopolymers*, 24 (1985), p. 461
- 25 Wolfe JK, Waugh DF. **Relations between enzymatic and association reactions in the development of bovine fibrin clot structure.** *Arch Biochem Biophys*, 211 (1981), p. 125
- 26 Shah GA, Ferguson IA, Dhall TZ, Dhall DP. **Polydispersion in the diameter of fibres in fibrin networks: Consequences on the measurement of mass-length ratio by permeability and turbidity.** *Biopolymers*, 21 (1982), p. 1037
- 27 Shah GA, Nair CH, Dhall DP. **Physiological studies on fibrin network structure.** *Thromb Res*, 40 (1985), p. 181
- 28 Mosesson MW, Siebenlist KR, Amrani DL, DiOrio JP. **Identification of covalently linked trimeric and tetrameric D domains in crosslinked fibrin.** *Proc Natl Acad Sci USA*, 86 (1989), p. 1113

- 29 Mosesson MW, Sherry S. **The preparation and properties of human fibrinogen of relatively high solubility.** *Biochemistry*, 5 (1966), p. 2829
- 30 Mosesson MW, Alkjaersig N, Sweet B, Sherry S. **Human fibrinogen of relatively high solubility. Comparative biophysical, biochemical and biological studies with fibrinogen of lower solubility.** *Biochemistry*, 6 (1967), p. 3279
- 31 Mills D, Karpatkin S. **Heterogeneity of human fibrinogen: Possible relation to proteolysis by thrombin and plasmin as studied by SDS-polyacrylamide gel electrophoresis.** *Biochem Biophys Res Commun*, 40 (1970), p. 206
- 32 Mosesson MW, Finlayson JS, Umfleet RA, Galanakis D. **Heterogeneities of human fibrinogen I. Structural and related studies of plasma fibrinogens which are high solubility catabolic intermediates.** *J Biol Chem*, 247 (1972), p. 5210
- 33 Hasegawa N, Sasaki S. **Location of the binding site "b" for lateral polymerization of fibrin.** *Thromb Res*, 57 (1990), p. 183
- 34 Gabriel DA, Smith LA, Folds JD, Davis L, Canelosi SE. **The influence of immunoglobulin (IgG) on the assembly of fibrin gels.** *J Lab Clin Med*, 101 (1983), p. 545
- 35 Bale MD, Mosher DF. **Effects of thrombospondin on fibrin polymerization and structure.** *J Biol Chem*, 261 (1986), p. 862
- 36 Leung LLK. **Interaction of histidine-rich glycoprotein with fibrinogen and fibrin.** *J Clin Invest*, 77 (1986), p. 1305
- 37 Galanakis DK, Lane BP, Simon SR. **Albumin modulates lateral assembly of fibrin polymers: Evidence of enhanced fine fibril formation and of unique synergism with fibrinogen.** *Biochemistry*, 26 (1987), p. 2389
- 38 Hermans J, McDonagh J. **Fibrin: Structure and interactions.** *Semin Thromb Haemost*, 8 (1982), p. 11
- 39 Erickson HP, Fowler WE. **Electron microscopy of fibrinogen, its plasmic fragments and small polymers.** *Ann NY Acad Sci*, 408 (1983), p. 146
- 40 Weisel JW, Nagaswami C. **Computer modeling of fibrin polymerization kinetics correlated with electron microscopic and turbidity observations: Clot structure and assembly are kinetically controlled.** *Biophys J*, 63 (1992), p. 111
- 41 Wall JS, Hainfeld JF. **Mass mapping with the scanning transmission electron microscope.** *Ann Rev Biophys Biophys Chem*, 15 (1986), p. 355
- 42 Hainfeld JF, Wall JS, Desmond EJ. **A small computer system for micrograph analysis.** *Ultramicroscopy*, 8 (1982), p. 263
- 43 Wall JS. **Biological scanning transmission electron microscopy.** Hren JJ, Goldstein JI, Joy DC (Eds.), *Introduction to Analytical Electron Microscopy*, Plenum, New York, NY (1979), p. 333
- 44 Wall JS. **Limits on visibility of single heavy atoms in the scanning transmission electron microscope—An experimental study.** *Chem Scripta*, 15 (1979), p. 271
- 45 Kramer O. *Biological and Synthetic Polymer Networks*, Elsevier, New York, NY (1988)
- 46 Hunziker EB, Straub PW, Haeberli A. **A new concept of fibrin formation based upon the linear growth of interlacing and branching polymers and molecular alignment into interlocked single-stranded segments.** *J Biol Chem*, 265 (1990), p. 7455