Marquette University

e-Publications@Marquette

Psychology Faculty Research and Publications

Psychology, Department of

4-2017

Clarifying the Relationship Between Trichotillomania and Anxiety

Jennifer R. Alexander
Marquette University

David C. Houghton
Texas A&M University - College Station

Michael P. Twohig
Utah State University

Martin E. Franklin
University of Pennsylvania

Stephen M. Saunders
Marquette University, stephen.saunders@marquette.edu

See next page for additional authors

Follow this and additional works at: https://epublications.marquette.edu/psych_fac

Part of the [Psychology Commons](#)

Recommended Citation

Alexander, Jennifer R.; Houghton, David C.; Twohig, Michael P.; Franklin, Martin E.; Saunders, Stephen M.; Neal-Barnett, Angela M.; Compton, Scott N.; and Woods, Douglas W., "Clarifying the Relationship Between Trichotillomania and Anxiety" (2017). *Psychology Faculty Research and Publications*. 282.
https://epublications.marquette.edu/psych_fac/282

Authors

Jennifer R. Alexander, David C. Houghton, Michael P. Twohig, Martin E. Franklin, Stephen M. Saunders, Angela M. Neal-Barnett, Scott N. Compton, and Douglas W. Woods

Marquette University

e-Publications@Marquette

Psychology Faculty Research and Publications/College of Arts and Sciences

This paper is NOT THE PUBLISHED VERSION; but the author's final, peer-reviewed manuscript. The published version may be accessed by following the link in the citation below.

Journal of Obsessive-Compulsive and Related Disorders, Vol. 13 (April, 2017): 30-34. [DOI](#). This article is © Elsevier and permission has been granted for this version to appear in [e-Publications@Marquette](#). Elsevier does not grant permission for this article to be further copied/distributed or hosted elsewhere without the express permission from Elsevier.

Clarifying the Relationship Between Trichotillomania and Anxiety

[Jennifer R. Alexander](#)

Department of Psychology, Marquette University, Milwaukee, WI

[David C. Houghton](#)

Department of Psychology, Texas A&M University, College Station, TX

[Michael P. Twohig](#)

Department of Psychology, Utah State University, Logan, UT

[Martin E. Franklin](#)

Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA

[Stephen M. Saunders](#)

Department of Psychology, Marquette University, Milwaukee, WI

[Angela M. Neal-Barnett](#)

Department of Psychology, Kent State University, Kent, OH

[Scott N. Compton](#)

Department of Psychiatry and Behavioral Sciences, Duke University School of Medicine, NC

Douglas W. Woods

Department of Psychology, Marquette University, Milwaukee, WI

Abstract

Although research has consistently linked unidimensional anxiety with [Trichotillomania](#) (TTM) severity, the relationships between TTM severity and anxiety dimensions (e.g., cognitive and somatic anxiety) are unknown. This knowledge gap limits current TTM conceptualization and treatment. The present study examined these relationships with data collected from ninety-one adults who participated in a [randomized clinical trial](#) for TTM treatment. Based on prior research, it was hypothesized that TTM severity would be related to the cognitive anxiety dimension and that psychological inflexibility would mediate the association. Hypotheses were not made regarding the relationship between TTM severity and [somatic anxiety](#). Regression analyses indicated that only cognitive dimensions of anxiety predicted TTM severity and that psychological inflexibility mediated this relationship. Implications for the conceptualization and treatment of TTM are discussed.

Keywords

Trichotillomania, Cognitive anxiety, Somatic anxiety, Psychological inflexibility

1. Clarifying the relationship between Trichotillomania and anxiety

[Trichotillomania](#) (TTM; also referred to as Hair Pulling Disorder) is a debilitating condition defined by recurrent hair pulling despite repeated attempts to stop ([American Psychiatric Association \[APA\], 2013](#)). Based on research suggesting that negative affect, particularly anxiety, contributes to disorder duration and severity ([Christenson et al., 1991](#), [Christenson and Mansueto, 1999](#), [Hajcak et al., 2006](#), [Mansueto et al., 1997](#); [Roberts, O'Connor, & Bélanger, 2013](#); [Stanley, Borden, Bell, & Wagner, 1994](#)), various TTM treatment models augment traditionally effective behavioral treatments (e.g., Habit Reversal Therapy; [Azrin & Nunn, 1973](#)) with anxiety-targeting techniques. However, different TTM treatment models target different anxiety components. For instance, comprehensive behavioral treatment models tend to target physiological symptoms of anxiety via techniques like deep breathing ([Azrin, Nunn, & Frantz, 1980](#)); cognitive-behavioral treatment (CBT) models target [anxiogenic](#) cognitions and feelings via techniques like cognitive restructuring ([Lerner, Franklin, Meadows, Hembree, & Foa, 1999](#)); and third-wave behavioral treatment models target meta-cognitions about experiencing anxiety via techniques like [mindfulness](#) ([Keuthen et al., 2010](#), [Woods and Twohig, 2008](#)). Unfortunately, there is currently little empirical basis for targeting any one component of anxiety over another, as the specific nature of the relationship between anxiety and TTM is unknown. Developing a better understanding of how the various components of anxiety relate to TTM is necessary for developing a better understanding of the disorder and its treatment.

TTM research has predominantly treated anxiety as a unidimensional construct (e.g., [Diefenbach, Mouton-Odum, & Stanley, 2002](#); [Diefenbach, Tolin, Hannan, Crocetto, & Worhunsky, 2005](#); [Diefenbach, Tolin, Meunier, & Worhunsky, 2008](#); [Duke, Bodzin, Tavers, Geffken, & Storch, 2009](#); [Houghton et al., 2014](#); [Shusterman, Feld, Baer, & Keuthen, 2009](#); [Stanley et al., 1994](#)). However, anxiety researchers recognize anxiety as a multidimensional

construct consisting of cognitive (sometimes referred to as subjective anxiety; e.g., fear and worry) and [somatic](#) (e.g., physiological arousal and panic) dimensions ([Clark and Watson, 1991](#), [Nitschke et al., 2001](#), [Ree et al., 2008](#)). Conceptualizing anxiety in such a way is clinically meaningful, as research suggests that cognitive and [somatic anxiety](#) symptoms may respond best to different treatments ([Norton and Johnson, 1983](#), [Ree et al., 2008](#), [Schwartz et al., 1978](#), [Tamaren et al., 1985](#)). Specifically, some research suggests that cognitive anxiety may be most responsive to treatments like cognitive restructuring and meditation, while somatic anxiety may be most responsive to diaphragmatic breathing exercises and [progressive muscle relaxation](#) ([Norton and Johnson, 1983](#), [Ree et al., 2008](#), [Schwartz et al., 1978](#), [Tamaren et al., 1985](#)). Accordingly, to develop treatments that effectively target anxiety associated with TTM, it is important to understand the relationships between the different dimensions of anxiety and TTM severity.

The few studies that have examined the relationships between TTM severity and the cognitive and somatic anxiety dimensions have not been conclusive. Some evidence suggests that TTM severity is related to aspects of cognitive anxiety, such as worry ([Hajcak et al., 2006](#)), and other evidence suggests TTM is related to fear of negative evaluation ([Norberg, Wetterneck, Woods, & Conelea, 2007](#)). Still, other evidence suggests that while TTM-affected persons demonstrate higher levels of cognitive and somatic anxiety than unaffected persons, TTM severity may not be related to either anxiety dimension ([Wetterneck, Lee, Flessner, Leonard, & Woods, 2016](#)). However, this study was limited by a small sample size that may have precluded power to detect such relationships. Overall, research appears to suggest that TTM severity may be related to cognitive aspects of anxiety, but it is unclear whether TTM severity is related to somatic anxiety.

In addition to understanding how particular aspects of anxiety are related to TTM symptoms, researchers have also examined how anxiety symptoms are perceived and acted upon by persons with TTM. To this end, a growing body of research focuses on the role of psychological inflexibility in TTM. Psychological inflexibility refers to the propensity to prioritize the mitigation of undesirable private experiences over the pursuit of adaptive goals ([Bond et al., 2011](#)). Generally, the most frequently discussed form of psychological inflexibility is experiential avoidance, which refers to the propensity to engage in behaviors that facilitate disengagement from undesirable experiences ([Bond et al., 2011](#)). Studies that have examined the role of psychological inflexibility in TTM have found that psychological inflexibility mediates the relationships between hair pulling severity and unidimensional measures of anxiety ([Houghton et al., 2014](#)) as well as between hair pulling severity and certain cognitive aspects of anxiety (i.e., apprehension/anxious thoughts about being negatively evaluated by others; [Norberg et al., 2007](#)). However, it is unclear whether psychological inflexibility mediates the relationship between TTM severity and the broader cognitive anxiety dimension. Further, no study has examined whether psychological inflexibility mediates the relationship between TTM severity and somatic aspects of anxiety.

Since the introduction of “psychological inflexibility” as a construct, it has been implied that different psychiatric disorders are characterized by different manifestations of psychological inflexibility. That is, the manner in which psychological inflexibility manifests in [social anxiety disorder](#) may differ from the manner in which psychological inflexibility manifests in TTM.

Accordingly, it has been proposed that using disorder-specific measures of psychological inflexibility may yield more precise information about the manner in which psychological inflexibility is involved in that disorder ([Houghton et al., 2014](#), [MacKenzie and Kocovski, 2010](#), [Sandoz et al., 2013](#)). Although such a measure has been created to measure psychological inflexibility descriptive of TTM, the Acceptance and Action Questionnaire-Trichotillomania (AAQ-TTM; [Houghton et al., 2014](#)), no study has examined whether TTM-specific psychological inflexibility mediates the relationships between TTM severity and the cognitive and somatic dimensions of anxiety.

The current study was intended to explore the anxiety-TTM relationship. Using the same data set utilized by [Houghton et al. \(2014\)](#), the current study examined whether TTM severity (assessed via self-report and clinician rating) is differentially related to different anxiety dimensions (i.e., cognitive and somatic). In addition, the current study examined whether TTM-specific psychological inflexibility mediated the relationship between these anxiety dimensions and TTM severity.

Based on the findings of [Hajcak et al. \(2006\)](#) and [Norberg et al. \(2007\)](#), we hypothesized that the cognitive domain of anxiety would be significantly related to TTM severity. In addition, consistent with findings from [Norberg et al. \(2007\)](#), we hypothesized that TTM-specific psychological inflexibility would mediate the relationship between cognitive anxiety and TTM severity. Based on prior research showing no relationship between somatic forms of anxiety and TTM ([Wetterneck et al., 2016](#)), we did not anticipate finding an association between TTM severity and the somatic dimension of anxiety in the current study.

2. Method

2.1. Participants

Data were collected as part of a [randomized clinical trial](#) for TTM treatment. Information about the sample and recruitment can be found in [Houghton et al. \(2016\)](#).

Ninety-one participants (92% female; 8% male) were included in the current study. Seventy-six participants identified as European American (84%), eleven identified as African American (12%), one identified as Asian American (1%), and three identified as multiracial or did not disclose their race (3%). In addition, one participant identified as Hispanic or Latino (1%). Participants had a mean age of 35.04 years ($SD=12.68$).

2.2. Measures

2.2.1. Cognitive and somatic anxiety

The [Beck Anxiety Inventory](#) (BAI; [Beck, Epstein, Brown, & Steer, 1988](#)) is a 21-item self-report measure of clinical anxiety that assesses both [somatic](#) and cognitive symptoms of anxiety. Each item on the BAI is ranked on a 0-3 scale. Thus, raw scores range from 0–63, with higher scores indicating higher levels of anxiety. Although the BAI total score is frequently used in research, several studies suggest that the BAI consists of multiple factors. Two studies ([Beck and Steer, 1991](#), [Steer et al., 1993](#)) exploring the factor structure of the BAI in clinically

anxious samples concluded that the BAI consists of four factors: subjective anxiety, neurophysiological anxiety, autonomic anxiety, and panic anxiety. These same researchers hypothesized that raw scores on the subjective anxiety factor assessed cognitive anxiety and raw scores on the neurophysiological, autonomic, and panic anxiety factors assessed [somatic anxiety](#) (Steer, Rissmiller, Ranieri, & Beck, 1993).

We calculated all four of the BAI factors observed by [Steer, Ranieri et al. \(1993\)](#) using data from the current sample. To examine whether the [Steer, Ranieri et al. \(1993\)](#) BAI factors were adequate in the current sample, we analyzed these factors' internal consistencies. This analysis indicated that the subjective anxiety ($\alpha=.86$), neurophysiological anxiety ($\alpha=.88$), and autonomic anxiety ($\alpha=.80$) factors each demonstrated acceptable internal consistency. The panic anxiety factor demonstrated poor internal consistency ($\alpha=.54$). Therefore, only the [Steer, Ranieri et al.'s \(1993\)](#) subjective, neurophysiological, and autonomic anxiety factors were analyzed in the current study. As proposed by [Steer, Rissimiller et al. \(1993\)](#), the subjective anxiety factor measured cognitive anxiety and the neurophysiological and autonomic anxiety factors measured somatic anxiety. Correlations between cognitive anxiety and the two forms of somatic anxiety (i.e., neurophysiological and autonomic) are presented in [Table 1](#).

Table 1. Pearson's Correlations between BAI Factors.

Factors	1	2	3
1.Cognitive Anxiety	-		
2.Neurophysiological Anxiety	.61**	-	
3.Autonomic Anxiety	.54**	.51**	-

** $p<.001$.

2.2.2. TTM severity

The *Massachusetts General Hospital Hairpulling Scale* (MGH-HS; [Keuthen et al., 1995](#)) is a seven-item self-report measure of hair pulling and hair pulling urge severity. Total scores on the MGH-HS range from 0–28, and higher scores indicate greater hair pulling severity. Research suggests the MGH-HS is reliable and valid in TTM samples ([Keuthen et al., 2007](#), [O'Sullivan et al., 1995](#)).

The *NIMH-Trichotillomania Severity Scale* (NIMH-TSS; [Swedo, Rapoport, Leonard, Lenane, & Cheslow, 1989](#)) is a five-item clinician rated measure of TTM severity. Total scores range from 0–25, with higher scores indicating greater TTM severity. Research suggests the NIMH-TSS demonstrates acceptable reliability and validity in TTM samples ([Diefenbach et al., 2005](#), [Swedo et al., 1989](#)).

2.2.3. TTM-specific psychological inflexibility

Based on the revised version of the Acceptance and Action Questionnaire (AAQ-II; [Bond et al., 2011](#)), which measures general psychological inflexibility, the Acceptance and Action Questionnaire *Trichotillomania* (AAQ-TTM; [Houghton et al., 2014](#)) is a nine-item self-report measure of TTM-specific psychological inflexibility. That is, the AAQ-TTM measures

respondents' tendencies to engage in TTM-related behaviors (e.g., hair pulling), at the expense of engaging in more adaptive behaviors, when experiencing unwanted experiences (e.g., urges to hair pull). Higher scores indicate greater TTM-specific psychological flexibility (i.e., greater ability to resist hair pulling when distressed by thoughts/urges related to hair pulling), while lower scores indicate greater TTM-specific psychological inflexibility (i.e., greater tendency to engage in hair pulling when experiencing thoughts/urges related to hair pulling). Preliminary research suggests that the AAQ-TTM has satisfactory validity and reliability in clinical TTM samples ([Houghton et al., 2014](#)).

2.3. Procedure

Information about the procedures used in the [randomized, controlled trial](#) can be found in [Houghton et al. \(2016\)](#). Only data from the baseline visit were used. During the baseline visit, participants completed the BAI, MGH-HS, and AAQ-TTM and were evaluated with the NIMH-TSS.

3. Results

Correlational analyses were used to examine the relationship between each of the anxiety factors, hair pulling severity, and TTM-specific psychological inflexibility (see [Table 2](#) for correlation coefficients). Results indicated that only cognitive anxiety factor was significantly correlated with TTM severity. Cognitive anxiety and the two forms of [somatic anxiety](#) (i.e., neurophysiological and autonomic) were correlated with TTM-specific psychological inflexibility.

Table 2. Pearson's Correlations between BAI Factors and Measures of TTM Severity and Psychological Inflexibility.

	MGH-HS	NIMH-TSS	AAQ-TTM
1. Cognitive Anxiety	.23*	.22*	-.45**
2. Neurophysiological Anxiety	.11	.13	-.30**
3. Autonomic Anxiety	.03	.11	-.13

Note. AAQ-TTM = Acceptance and Action Questionnaire [Trichotillomania](#); MGH-HS = Massachusetts General Hospital Hairpulling Scale; NIMH-TSS = National Institute of Mental Health Trichotillomania Severity Scale.

* $p < .05$,

** $p \leq .001$.

Because the cognitive anxiety factor was the only anxiety factor significantly correlated with TTM severity, only this relationship was explored further. The [Baron and Kenny \(1986\)](#) approach was used to examine whether TTM-specific psychological inflexibility mediated the relationship between cognitive anxiety and TTM severity. Results indicated that TTM-specific psychological inflexibility fully mediated the relationships between cognitive anxiety and both

self-reported TTM severity and clinician-rated TTM severity. The mediation analyses are summarized in [Fig. 1](#), [Fig. 2](#).

Fig. 1. Regression coefficients for the relationship between cognitive anxiety and MGH-HS as mediated by TTM-Specific experiential avoidance. MGH-HS = Massachusetts General Hospital Hairpulling Scale. $*p < .05$, $**p \leq .001$.

Fig. 2. Regression coefficients for the relationship between cognitive anxiety and NIMH-TSS as mediated by TTM-Specific experiential avoidance. NIMH-TSS = National Institute of Mental Health [Trichotillomania](#) Severity Scale. $*p < .05$, $**p \leq .001$.

4. Discussion

Research demonstrates that anxiety is associated with TTM, but the nuances of this relationship are relatively unknown. Accordingly, we explored the relationship between dimensions of anxiety (i.e., cognitive and [somatic](#) anxiety) and TTM severity. We also explored whether TTM-specific psychological flexibility mediated the relationships between anxiety dimensions and TTM severity. Based on research suggesting that hair pulling severity is related to cognitive aspects of anxiety ([Hajcak et al., 2006](#), [Norberg et al., 2007](#)) and that psychological inflexibility mediates this relationship ([Norberg et al., 2007](#)), we hypothesized that cognitive anxiety would be related to TTM severity and that TTM-specific psychological inflexibility would mediate this relationship. We also conducted analyses looking at whether somatic domains of anxiety were related to TTM severity.

The current results confirmed the hypothesis that greater cognitive anxiety was associated with greater TTM severity, whereas the two dimensions of [somatic anxiety](#) analyzed in the current study (i.e., neurophysiological and autonomic symptoms) were not associated with TTM severity. Further, the current results demonstrated that TTM-specific psychological inflexibility fully mediated the relationship between cognitive anxiety and TTM severity, suggesting that when an individual with TTM experiences cognitive anxiety, his or her pulling severity is primarily a function of his or her TTM-specific psychological flexibility. TTM-affected individuals with greater intolerance of thoughts and urges related to hair pulling will be significantly more likely to pull when cognitive anxiety is present; conversely, those who are more tolerable of thoughts and urges related to hair pulling will be able to resist pulling in the presence of cognitive anxiety.

Such findings have significant treatment implications. The current results suggest that reducing TTM-affected individuals' cognitive anxiety may lead to reductions in hair pulling severity. Accordingly, such results appear to provide support for supplementing traditional TTM [behavioral therapy](#) with techniques that target cognitive anxiety (e.g., cognitive restructuring and meditation; [Ree et al., 2008](#); [Schwartz et al., 1978](#); [Tamaren et al., 1985](#)). However, given that the current results suggest that cognitive and somatic anxiety are highly related, it is possible that TTM-affected individuals will also derive some therapeutic benefit from techniques that target somatic anxiety. Moreover, as results suggest TTM-specific psychological inflexibility mediates the relationship between cognitive anxiety and TTM severity, the current study supports the use of TTM treatment programs that target TTM-specific psychological inflexibility, such as ACT-enhanced behavior therapy (AEBT; [Woods & Twohig, 2008](#)), or dialectical behavior therapy-enhanced behavior therapy ([Keuthen et al., 2010](#)).

The current study did have limitations. One methodological limitation was the relatively small sample size. Nevertheless, given that TTM is a low-prevalence condition, this was a relatively large sample that was also well-characterized. An additional theoretical limitation was that the current study only considered the relationship between anxiety and TTM severity. Future research may also benefit from examining the relationship between other forms of negative affect and TTM severity.

In addition to providing further support for findings from [Haicak et al. \(2006\)](#) and [Norberg et al. \(2007\)](#), the current study suggests that TTM severity is differentially related to anxiety dimensions. Future research should continue to explore the relationship between anxiety dimensions and other variables relevant to TTM (e.g., disorder duration, quality of life, TTM disorder-related impairment, comorbidity, etc.). Future research may also benefit from examining the relationship between TTM severity and variables related to the experience of anxiety (e.g., anxiety sensitivity, affect regulation, and neural activation).

References

- 1 American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: Author.
- 2 N.H. Azrin, R.G. Nunn **Habit-reversal: A method of eliminating nervous habits and tics** *Behaviour Research and Therapy*, 11 (4) (1973), pp. 619-628, [10.1016/0005-7967\(73\)90119-8](#)
- 3 N.H. Azrin, R.G. Nunn, S.E. Frantz **Treatment of hairpulling (trichotillomania): A comparative study of habit reversal and negative practice training** *Journal of Behavior Therapy and Experimental Psychiatry*, 11 (1) (1980), pp. 13-20, [10.1016/0005-7916\(80\)90045-2](#)
- 4 R.M. Baron, D.A. Kenny **The moderator–mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations** *Journal of Personality and Social Psychology*, 51 (6) (1986), p. 1173, [10.1037/0022-3514.51.6.1173](#)
- 5 A.T. Beck, N. Epstein, G. Brown, R.A. Steer **An inventory for measuring clinical anxiety: Psychometric properties** *Journal of Consulting and Clinical Psychology*, 56 (6) (1988), pp. 893-897, [10.1037/0022-006X.56.6.893](#)
- 6 A.T. Beck, R.A. Steer **Relationship between the Beck Anxiety Inventory and the Hamilton Anxiety Rating Scale with anxious outpatients** *Journal of Anxiety Disorders*, 5 (3) (1991), pp. 213-223, [10.1016/0887-6185\(91\)90002-B](#)
- 7 F.W. Bond, S.C. Hayes, R.A. Baer, K.M. Carpenter, N. Guenole, H.K. Orcutt, R.D. Zettle **Preliminary psychometric properties of the Acceptance and Action Questionnaire–II: A revised measure of psychological inflexibility and experiential avoidance** *Behavior Therapy*, 42 (4) (2011), pp. 676-688, [10.1016/j.beth.2011.03.007](#)
- 8 G.A. Christenson, T.B. Mackenzie, J.E. Mitchell **Characteristics of 60 adult chronic hair pullers** *American Journal of Psychiatry*, 148 (1991), pp. 365-370, [10.1176/ajp.148.3.365](#)
- 9 G.A. Christenson, C.S. Mansueto **Trichotillomania: Descriptive characteristics and phenomenology** D. Stein, G. Christenson, E. Hollander (Eds.), *Trichotillomania*, American Psychiatric Press, Washington, DC (1999), pp. 1-41

- 10 L.A. Clark, D. Watson **Tripartite model of anxiety and depression: Psychometric evidence and taxonomic implications** *Journal of Abnormal Psychology*, 100 (3) (1991), p. 316, [10.1037/0021-843X.100.3.316](https://doi.org/10.1037/0021-843X.100.3.316)
- 11 G.J. Diefenbach, S. Mouton-Odum, M.A. Stanley **Affective correlates of trichotillomania** *Behaviour Research and Therapy*, 40 (11) (2002), pp. 1305-1315, [10.1016/S0005-7967\(02\)00006-2](https://doi.org/10.1016/S0005-7967(02)00006-2)
- 12 G.J. Diefenbach, D.F. Tolin, J. Crocetto, N. Maltby, S. Hannan **Assessment of trichotillomania: A psychometric evaluation of hair-pulling scales** *Journal of Psychopathology and Behavioral Assessment*, 27 (3) (2005), pp. 169-178, [10.1007/s10862-005-0633-7](https://doi.org/10.1007/s10862-005-0633-7)
- 13 G.J. Diefenbach, D.F. Tolin, S. Hannan, J. Crocetto, P. Worhunsky **Trichotillomania: Impact on psychosocial functioning and quality of life** *Behaviour Research and Therapy*, 43 (7) (2005), pp. 869-884, [10.1016/j.brat.2004.06.010](https://doi.org/10.1016/j.brat.2004.06.010)
- 14 G.J. Diefenbach, D.F. Tolin, S. Meunier, P. Worhunsky **Emotion regulation and trichotillomania: A comparison of clinical and nonclinical hair pulling** *Journal of Behavior Therapy and Experimental Psychiatry*, 39 (1) (2008), pp. 32-41, [10.1016/j.jbtep.2006.09.002](https://doi.org/10.1016/j.jbtep.2006.09.002)
- 15 D.C. Duke, D.K. Bodzin, P. Tavares, G.R. Geffken, E.A. Storch **The phenomenology of hairpulling in a community sample** *Journal of Anxiety Disorders*, 23 (8) (2009), pp. 1118-1125, [10.1016/j.janxdis.2009.07.015](https://doi.org/10.1016/j.janxdis.2009.07.015)
- 16 G. Hajcak, M.E. Franklin, R.F. Simons, N.J. Keuthen **Hairpulling and skin picking in relation to affective distress and obsessive-compulsive symptoms** *Journal of Psychopathology and Behavioral Assessment*, 28 (3) (2006), pp. 177-185, [10.1007/s10862-005-9001-x](https://doi.org/10.1007/s10862-005-9001-x)
- 17 D.C. Houghton, S.N. Compton, M.P. Twohig, S.M. Saunders, M.E. Franklin, A.M. Neal-Barnett, D.W. Woods **Measuring the role of psychological inflexibility in trichotillomania** *Psychiatry Research*, 220 (1) (2014), pp. 356-361, [10.1016/j.psychres.2014.08.003](https://doi.org/10.1016/j.psychres.2014.08.003)
- 18 D.C. Houghton, J. Maas, M.P. Twohig, S.M. Saunders, S.N. Compton, A.M. Neal-Barnett, D.W. Woods **Comorbidity and quality of life in adults with hair pulling disorder** *Psychiatry Research*, 239 (2016), pp. 12-19, [10.1016/j.psychres.2016.02.063](https://doi.org/10.1016/j.psychres.2016.02.063)
- 19 N.J. Keuthen, C.A. Flessner, D.W. Woods, M.E. Franklin, D.J. Stein, S.E. Cashin **Factor analysis of the Massachusetts General Hospital Hairpulling Scale** *Journal of Psychosomatic Research*, 62 (6) (2007), pp. 707-709, [10.1016/j.jpsychores.2006.12.003](https://doi.org/10.1016/j.jpsychores.2006.12.003)
- 20 N.J. Keuthen, R.L. O'Sullivan, J.N. Ricciardi, D. Shera, C.R. Savage, A.S. Borgmann, L. Baer **The Massachusetts General Hospital (MGH) Hairpulling Scale: 1.**

Development and factor analyses *Psychotherapy and Psychosomatics*, 64 (3-4) (1995), pp. 141-145, [10.1159/000289003](https://doi.org/10.1159/000289003)

- 21 N.J. Keuthen, B.O. Rothbaum, S.S. Welch, C. Taylor, M. Falkenstein, M. Heekin, M.A. Jenike **Pilot trial of dialectical behavior therapy-enhanced habit reversal for trichotillomania** *Depression and Anxiety*, 27 (10) (2010), pp. 953-959, [10.1002/da.20732](https://doi.org/10.1002/da.20732)
- 22 J. Lerner, M.E. Franklin, E.A. Meadows, E. Hembree, E.B. Foa **Effectiveness of a cognitive behavioral treatment program for trichotillomania: An uncontrolled evaluation** *Behavior Therapy*, 29 (1) (1999), pp. 157-171, [10.1016/S0005-7894\(98\)80036-1](https://doi.org/10.1016/S0005-7894(98)80036-1)
- 23 M.B. MacKenzie, N.L. Kocovski **Self-reported acceptance of social anxiety symptoms: Development and validation of the Social Anxiety—Acceptance and Action Questionnaire** *International Journal of Behavioral Consultation and Therapy*, 6 (3) (2010), pp. 214-231, [10.1037/h0100909](https://doi.org/10.1037/h0100909)
- 24 C.S. Mansueto, R.M.T. Stemberger, A.M. Thomas, R.G. Golomb **Trichotillomania: A comprehensive behavioral model** *Clinical Psychology Review*, 17 (5) (1997), pp. 567-577, [10.1016/S0272-7358\(97\)00028-7](https://doi.org/10.1016/S0272-7358(97)00028-7)
- 25 J.B. Nitschke, W. Heller, J.C. Imig, R.P. McDonald, G.A. Miller **Distinguishing dimensions of anxiety and depression** *Cognitive Therapy and Research*, 25 (1) (2001), pp. 1-22, [10.1023/A:1026485530405](https://doi.org/10.1023/A:1026485530405)
- 26 M.M. Norberg, C.T. Wetterneck, D.W. Woods, C.A. Conelea **Experiential avoidance as a mediator of relationships between cognitions and hair-pulling severity** *Behavior Modification*, 31 (4) (2007), pp. 367-381, [10.1177/0145445506297343](https://doi.org/10.1177/0145445506297343)
- 27 G.R. Norton, W.E. Johnson **A comparison of two relaxation procedures for reducing cognitive and somatic anxiety** *Journal of Behavior Therapy and Experimental Psychiatry*, 14 (3) (1983), pp. 209-214, [10.1016/0005-7916\(83\)90050-2](https://doi.org/10.1016/0005-7916(83)90050-2)
- 28 R.L. O'Sullivan, N.J. Keuthen, C.F. Hayday, J.N. Ricciardi, M.L. Buttolph, M.A. Jenike, L. Baer **The Massachusetts General Hospital (MGH) Hairpulling Scale: 2. Reliability and validity** *Psychotherapy and Psychosomatics*, 64 (3-4) (1995), pp. 146-148, [10.1159/000289003](https://doi.org/10.1159/000289003)
- 29 M.J. Ree, D. French, C. MacLeod, V. Locke **Distinguishing cognitive and somatic dimensions of state and trait anxiety: Development and validation of the State-Trait Inventory for Cognitive and Somatic Anxiety (STICSA)** *Behavioural and Cognitive Psychotherapy*, 36 (03) (2008), pp. 313-332, [10.1017/S1352465808004232](https://doi.org/10.1017/S1352465808004232)
- 30 S. Roberts, K. O'Connor, C. Bélanger **Emotion regulation and other psychological models for body-focused repetitive behaviors** *Clinical Psychology Review*, 33 (6) (2013), pp. 745-762, [10.1016/j.cpr.2013.05.004](https://doi.org/10.1016/j.cpr.2013.05.004)

- 31 E.K. Sandoz, K.G. Wilson, R.M. Merwin, K.K. Kellum **Assessment of body image flexibility: The Body Image-Acceptance and Action Questionnaire** Journal of Contextual Behavioral Science, 2 (1) (2013), pp. 39-48, [10.1016/j.jcbs.2013.03.002](https://doi.org/10.1016/j.jcbs.2013.03.002)
- 32 G.E. Schwartz, R.J. Davidson, D.J. Goleman **Patterning of cognitive and somatic processes in the self-regulation of anxiety: Effects of meditation versus exercise** Psychosomatic Medicine, 40 (4) (1978), pp. 321-328, [10.1097/00006842-197806000-00004](https://doi.org/10.1097/00006842-197806000-00004)
- 33 A. Shusterman, L. Feld, L. Baer, N. Keuthen **Affective regulation in trichotillomania: Evidence from a large-scale internet survey** Behaviour Research and Therapy, 47 (8) (2009), pp. 637-644, [10.1016/j.brat.2009.04.004](https://doi.org/10.1016/j.brat.2009.04.004)
- 34 M.A. Stanley, J.W. Borden, G.E. Bell, A.L. Wagner **Nonclinical hair pulling: Phenomenology and related psychopathology** Journal of Anxiety Disorders, 8 (2) (1994), pp. 119-130, [10.1016/0887-6185\(94\)90010-8](https://doi.org/10.1016/0887-6185(94)90010-8)
- 35 R.A. Steer, W.F. Ranieri, A.T. Beck, D.A. Clark **Further evidence for the validity of the Beck Anxiety Inventory with psychiatric outpatients** Journal of Anxiety Disorders, 7 (3) (1993), pp. 195-205, [10.1016/0887-6185\(93\)90002-3](https://doi.org/10.1016/0887-6185(93)90002-3)
- 36 R.A. Steer, D.J. Rissmiller, W.F. Ranieri, A.T. Beck **Structure of the computer-assisted Beck Anxiety Inventory with psychiatric inpatients** Journal of Personality Assessment, 60 (3) (1993), pp. 532-542, [10.1207/s15327752jpa6003_10](https://doi.org/10.1207/s15327752jpa6003_10)
- 37 S.E. Swedo, J.L. Rapoport, H. Leonard, M. Lenane, D. Cheslow **Obsessive-compulsive disorder in children and adolescents: Clinical phenomenology of 70 consecutive cases** Archives of General Psychiatry, 46 (4) (1989), pp. 335-341, [10.1001/archpsyc.1989.01810040041007](https://doi.org/10.1001/archpsyc.1989.01810040041007)
- 38 A.J. Tamaren, R.M. Carney, T.W. Allen **Predictive validity of the cognitive vs. somatic anxiety distinction** The Pavlovian Journal of Biological Science: Official Journal of the Pavlovian, 20 (4) (1985), pp. 177-180, [10.1007/BF03003656](https://doi.org/10.1007/BF03003656)
- 39 C.T. Wetterneck, E.B. Lee, C.A. Flessner, R.C. Leonard, D.W. Woods **Personality characteristics and experiential avoidance in Trichotillomania: Results from an age and gender matched sample** Journal of Obsessive-Compulsive and Related Disorders, 8 (2016), pp. 64-69, [10.1016/j.jocrd.2015.12.003](https://doi.org/10.1016/j.jocrd.2015.12.003)
- 40 D.W. Woods, M.P. Twohig **Trichotillomania: An ACT-enhanced behavior therapy approach therapist guide** Oxford University Press, New York (2008)