Marquette University

e-Publications@Marquette

Chemistry Faculty Research and Publications

Chemistry, Department of

12-15-2014

Mono-*N*-acyl-2,6-diaminopimelic Acid Derivatives: Analysis by Electromigration and Spectroscopic Methods and Examination of Enzyme Inhibitory Activity

Jan Hlaváček

Academy of Sciences of the Czech Republic

Miloslava Vítovcová

Academy of Sciences of the Czech Republic

Petra Sázelová

Academy of Sciences of the Czech Republic

Jan Pícha

Academy of Sciences of the Czech Republic

Václav Vaněk

Academy of Sciences of the Czech Republic

See next page for additional authors

Follow this and additional works at: https://epublications.marquette.edu/chem_fac

Part of the [Chemistry Commons](#)

Recommended Citation

Hlaváček, Jan; Vítovcová, Miloslava; Sázelová, Petra; Pícha, Jan; Vaněk, Václav; Buděšínský, Miloš; Jiráček, Jiří; Gillner, Danuta M.; Holz, Richard C.; Mikšík, Ivan; and Kašička, Václav, "Mono-*N*-acyl-2,6-diaminopimelic Acid Derivatives: Analysis by Electromigration and Spectroscopic Methods and Examination of Enzyme Inhibitory Activity" (2014). *Chemistry Faculty Research and Publications*. 355.

https://epublications.marquette.edu/chem_fac/355

Authors

Jan Hlaváček, Miloslava Vítovcová, Petra Sázelová, Jan Pícha, Václav Vaněk, Miloš Buděšínský, Jiří Jiráček, Danuta M. Gillner, Richard C. Holz, Ivan Mikšík, and Václav Kašíčka

Marquette University

e-Publications@Marquette

Chemistry Faculty Research and Publications/College of Arts and Sciences

This paper is NOT THE PUBLISHED VERSION; but the author's final, peer-reviewed manuscript. The published version may be accessed by following the link in the citation below.

Analytical Biochemistry, Vol. 467 (December 15, 2014): 4-13. [DOI](#). This article is © Elsevier and permission has been granted for this version to appear in [e-Publications@Marquette](#). Elsevier does not grant permission for this article to be further copied/distributed or hosted elsewhere without the express permission from Elsevier.

Mono-N-Acyl-2,6-Diaminopimelic Acid Derivatives: Analysis by Electromigration And Spectroscopic Methods and Examination of Enzyme Inhibitory Activity

Jan Hlaváček

Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic, 166 10 Prague 6, Czech Republic

Miloslava Vítovcová

Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic, 166 10 Prague 6, Czech Republic

Petra Sázelová

Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic, 166 10 Prague 6, Czech Republic

Jan Pícha

Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic, 166 10 Prague 6, Czech Republic

Václav Vaněk

Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic, 166 10 Prague 6, Czech Republic

Miloš Buděšínský

Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic, 166 10 Prague 6, Czech Republic

Jiří Jiráček

Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic, 166 10 Prague 6, Czech Republic

Danuta M. Gillner

Department of Chemistry and Biochemistry, Loyola University Chicago, Chicago, IL
Department of Chemistry, Silesian University of Technology, 44-100 Gliwice, Poland

Richard C. Holz

Department of Chemistry and Biochemistry, Loyola University Chicago, Chicago, IL

Ivan Mikšík

Institute of Physiology, Academy of Sciences of the Czech Republic, 142 20 Prague 4, Czech Republic

Václav Kašička

Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic, 166 10 Prague 6, Czech Republic

Abstract

Thirteen mono-*N*-acyl derivatives of 2,6-diaminopimelic acid (DAP)—new [potential](#) inhibitors of the dapE-encoded *N*-succinyl-L,L-diaminopimelic acid desuccinylase (DapE; EC 3.5.1.18)—were analyzed and characterized by [infrared](#) (IR) and [nuclear magnetic resonance](#) (NMR) spectroscopies and two capillary electromigration methods: [capillary zone electrophoresis](#) (CZE) and [micellar electrokinetic chromatography](#) (MEKC). Structural features of DAP derivatives were characterized by [IR and NMR spectroscopies](#), whereas CZE and MEKC were applied to evaluate their purity and to investigate their electromigration properties. Effective [electrophoretic mobilities](#) of these compounds were determined by CZE in acidic and alkaline background electrolytes (BGEs) and by MEKC in acidic and alkaline BGEs containing a pseudostationary phase of anionic [detergent sodium dodecyl sulfate](#) (SDS) or cationic detergent [cetyltrimethylammonium bromide](#) (CTAB). The best separation of DAP derivatives, including [diastereomers](#) of some of them, was achieved by MEKC in an acidic BGE (500 mM [acetic acid](#) [pH 2.54] and 60 mM SDS). All DAP derivatives were examined for their ability to inhibit [catalytic activity](#) of DapE from [Haemophilus influenzae](#) (HiDapE) and ArgE from [Escherichia coli](#) (EcArgE). None of these DAP derivatives worked as an effective inhibitor of HiDapE, but one derivative—*N*-fumaryl, Me-ester-DAP—was found to be a moderate inhibitor of EcArgE, thereby providing a promising lead structure for further studies on ArgE inhibitors.

Keywords

2,6-Diaminopimelic acid derivatives, Capillary zone electrophoresis, Micellar electrokinetic chromatography, Enzyme inhibition, IR spectroscopy, NMR spectroscopy

The [World Health](#) Organization reported that bacterial infections represent a significant and permanently growing medical problem around the world due to the increasing number of disease-causing [microbes](#) that have become resistant to currently available [antibiotics](#) [1], [2], [3]. In fact, several pathogenic [bacteria](#), some of which were [thought](#) to have been eradicated, have made a significant resurgence due to bacterial [resistance](#) to [antibiotics](#) [4], [5]. For example, resistance to [tetracyclines](#) increased from 0% in 1948 to 98% in 1998 [4], [5], and tuberculosis is currently one of the leading causes of death in adults by an infectious disease worldwide, which is significant given that [death rates](#) due to tuberculosis had declined to near

imperceptible levels in industrial nations [6], [7]. According to the Centers for Disease Control and Prevention (USA), several [bacterial strains](#) currently exhibit [multidrug resistance](#), with more than 60% of hospital-acquired infections in the United States alone caused by the so-called ESKAPE pathogens (*Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, and *Enterobacter* sp.). These findings have stimulated an intensive search for new potent antimicrobial agents that are active against drug-resistant bacterial strains [5], [8].

Inhibitors of cell wall biosynthesis—vancomycin and [\$\beta\$ -lactams](#), to [name](#) a couple—have proven to be very potent [antibiotics](#), evidence that interfering with cell wall synthesis has deleterious effects on bacterial [cell survival](#). Enzymes that are targeted by these [antibiotics](#) tend to be present in all [bacteria](#) and are highly similar in structure and function. Unfortunately, only two new [antibacterial drugs](#) have emerged since 1962. Because every antibiotic has a finite [lifetime](#) given that [resistance](#) will ultimately occur, particularly if the same enzymes are repeatedly targeted, [development](#) of new classes of inhibitors that target previously untargeted cellular enzymes is essential to retain control of infectious disease [5], [8]. As such, [antimicrobial peptides](#) have been introduced as an innate defense system and shown to provide protection against a wide variety of [microorganisms](#) [9], [10], [11], [12]. Another approach targets microbial enzymes, especially those catalyzing [metabolic](#) processes exclusive to bacteria, as [potential](#) targets for potent and selective antibiotics [8].

Recently, we described a series of inhibitors for the bacterial enzyme *N*^α-acetyl-l-ornithine deacetylase (ArgE),² which catalyzes the conversion of *N*^α-acetyl-l-ornithine to l-ornithine in the fifth step of the [biosynthetic pathway](#) for [arginine](#), a necessary step for [bacterial growth](#) [13]. Based on [bacterial genetic](#) information, the *meso*-diaminopimelate (mDAP)/lysine biosynthetic pathway (Fig. 1) offers several [potential](#) antibacterial targets that have yet to be explored [14], [15], [16], [17]. One of the products of this pathway, [lysine](#) (10), is required in [protein synthesis](#) and is also used in the [peptidoglycan](#) layer of gram-positive [bacteria](#) cell walls. A second product, *meso*-diaminopimelic acid (mDAP) (9), is an essential component of the [peptidoglycan](#) cell wall in [gram-negative bacteria](#), providing a link between [polysaccharide](#) strands. It has been shown that deletion of the [gene](#) encoding for one of the enzymes in the mDAP/lysine biosynthetic pathway, the DapE-encoded *N*-succinyl-l,l-diaminopimelic acid desuccinylase (DapE; EC 3.5.1.18) [18], is [lethal](#) to *Helicobacter pylori* and *Mycobacterium smegmatis* [19], [20]. Even in the presence of lysine-supplemented media, *H. pylori* was unable to grow. Therefore, DapE is essential for [cell growth and proliferation](#) and is a part of a biosynthetic pathway that is the only source of [lysine](#) in bacteria. Because there are no similar biosynthetic pathways in [mammals](#), inhibitors that target one or more of the enzymes in the mDAP/lysine biosynthetic pathway are hypothesized to exhibit selective toxicity only against bacteria, providing a previously undescribed class of [antimicrobial agents](#) [14], [21].

Fig.1. Biosynthetic reactions leading from [aspartate](#) to [lysine](#): l-aspartate (1); *N*-succinyl-l,l-diaminopimelate (7); l,l-diaminopimelate (8); d,l-diaminopimelate (9); l-lysine (10).

With the aim of developing new [antimicrobial agents](#) that target DapE enzymes, a methodology for the quick and efficient preparation of mono-*N*-acylated 2,6-diaminopimelic acid (DAP) derivatives was recently

reported [22]. These compounds were designed as competitive inhibitors of the mono-*N*-succinyl-DAP [hydrolysis](#) reaction catalyzed by DapE, in which the structure of the N-linked [succinate](#) moiety was altered to inhibit the enzyme's ability to cleave the adjacent [amide](#) bond. The syntheses of a series of 13 DAP derivatives ([Table 1](#)) bearing different N-linked acyl side chains comprising (i) a hydrophobic aliphatic or aromatic moiety (**A1–A6**), (ii) an aliphatic moiety terminated by a carboxyl group (**C1–C4**), (iii) an aliphatic moiety terminated by an [ester](#) group (**E1–E3**), and their [purification](#) and characterization by [high-performance liquid chromatography](#) (HPLC), [elemental analysis](#), and [electrospray ionization–mass spectrometry](#) (ESI–MS) were described previously [22].

Table 1. [Name](#), indication, structural formula, and relative [molecular mass](#) (M_r) of 2,6-diaminopimelic acid and its mono-*N*-acyl derivatives.

Compound name and indication	Structural formula	M_r
2,6-Diaminopimelic acid		190.2
DAP		
<i>N</i> -Acetyl-DAP		232.1
A1		
<i>N</i> -Isobutyryl-DAP		260.3
A2		
<i>N</i> -Butyryl-DAP		260.3
A3		
<i>N</i> -Pivalyl DAP		274.3
A4		
<i>N</i> -Valeryl-DAP		274.3

A5		
N-Benzoyl-DAP		294.2
A6		
N-Glutaryl-DAP		304.3
C1		
N-3-Me-glutaryl-DAP		318.3
C2		
N-3,3-di-Me-glutaryl-DAP		332.4
C3		
N-2,3-Tetrafluoro-succinyl-DAP		362.2
C4		
N-Fumaryl, Me-ester-DAP		302.3
E1		
N-Malonyl, Et-ester-DAP		304.3
E2		

N-2-Me-succinyl, Me-ester-DAP		318.3
E3		

We report here the analytical characterization of these DAP derivatives in order to confirm their structural properties using [infrared](#) (IR) and ^1H and ^{13}C [nuclear magnetic resonance](#) (NMR) spectroscopies. We also performed qualitative and quantitative analysis of each of these compounds by two capillary electromigration methods: [capillary zone electrophoresis](#) (CZE) and [micellar electrokinetic chromatography](#) (MEKC). These two techniques are excellent tools for separation of both ionogenic and electroneutral compounds [23], [24], [25] and are frequently used for analysis and characterization of amino acids and their derivatives [26], [27], [28]. In addition, these DAP derivatives were characterized by their effective [electrophoretic mobilities](#) determined by CZE in acidic and alkaline classical or isoelectric background electrolytes (BGEs) or by MEKC in acidic and alkaline BGEs containing a micellar pseudostationary phase constituted by the anionic [detergent sodium dodecyl sulfate](#) (SDS) or the cationic detergent [cetyltrimethylammonium bromide](#) (CTAB). Finally, all of these DAP derivatives were tested for their [inhibition](#) activity toward both DapE and ArgE.

Materials and methods

Chemicals

All chemicals used were of analytical reagent grade. Tris(hydroxymethyl)aminomethane (Tris) was purchased from Serva (Heidelberg, Germany). [Phosphoric acid](#) (H_3PO_4), [acetic acid](#) (AcOH), and [sodium hydroxide](#) (NaOH) were obtained from Lachema (Brno, Czech Republic). [Iminodiacetic acid](#) (IDA) was obtained from Bachem (Bubendorf, Switzerland). SDS and CTAB were supplied by Fluka (Buchs, Switzerland). [Acetonitrile](#) (ACN) was obtained from Sigma–Aldrich (Prague, Czech Republic) and [methanol](#) (MeOH) from Penta (Chrudim, Czech Republic).

IR and NMR spectroscopies

[Fourier transform](#) (FT)-IR [spectra](#) (KBr , ν_{max} cm^{-1}) were recorded on a Nicolet 6700 FT-IR instrument (Madison, WI, USA). ^1H and ^{13}C NMR spectra were obtained on a Bruker AVANCE-600 NMR spectrometer (Bruker BioSpin, Rheinstetten, Germany) using a 5-mm CPTCI cryoprobe (^1H at 600.13 MHz, ^{13}C at 150.9 MHz) in [dimethyl sulfoxide](#) ($\text{DMSO}-d_6$) at 300 K. A series of [proton](#) 1D (one-dimensional), 2D–COSY (two-dimensional correlation spectroscopy), [2D–TOCSY](#) (total correlation spectroscopy) (spin-lock time 90 ms), and 2D–ROESY (rotating frame NOE [nuclear Overhauser effect] spectroscopy) (mixing time 300 ms) spectra were recorded at 27 °C for complete structural assignment of the observed [proton](#) signals. The structural assignment of the ^{13}C signals was achieved by combining 1D–APT (attached proton test), 2D– ^1H , ^{13}C –HSQC (heteronuclear single quantum coherence), and 2D– ^1H , ^{13}C –HMBC (heteronuclear multiple bond coherence) spectra.

Capillary electromigration methods

CZE and MEKC were performed on a homemade manually operated apparatus for [high-performance capillary electrophoresis](#) [29] equipped with an ultraviolet (UV) [absorption](#) photometric detector operating at 206 nm. Internally untreated fused silica capillary with outer [polyimide](#) coating (Polymicro Technologies, Phoenix, AZ, USA) with inner diameter of 50 μm , outer diameter of 375 μm , total [length](#) of 400 mm, and effective length (from injection to detector) of 290 mm was used for all separations. [Data acquisition](#) and evaluation was carried out by [Chromatography](#) and [ElectrophoresisStation Clarity](#) (DataApex, Prague, Czech Republic). The analyzed DAP derivatives were dissolved in the BGE or water in the concentration range 0.5 to 1.1 mg/ml and were analyzed as [cations](#) in acidic BGEs or as [anions](#) in alkaline BGEs. The BGEs were passed through a 0.45- μm Millipore filter (Bedford, MA, USA). The nanoliter [sample volumes](#) were introduced into the capillary

hydrodynamically by pneumatically induced [pressure](#) (600–1000 Pa) for a period of 3 to 20 s. The applied separation [voltage](#) was +12 kV or –12 kV (anode or [cathode](#) at the injection capillary end), whereas the [electric current](#) was in the range 7.1 to 45.0 μ A at an [ambient temperature](#) of 22 to 26 °C.

Determination of IC₅₀

The DapE from *Haemophilus influenzae* (HiDapE) was purified as described previously [18]. IC₅₀ values were determined using 3 mM *N*-succinyl-2,6-diaminopimelic acid (mixture of d,d-l,l isoforms, 1:1) as the substrate. All [kinetic](#) experiments were performed in 50 mM Chelex-100-treated sodium [phosphate buffer](#) at [pH](#) 7.5 and 25 °C. [Catalytic activities](#) were determined by monitoring the initial rates of [amide](#) bond cleavage at 225 nm ($\epsilon = 698 \text{ M}^{-1} \text{ cm}^{-1}$) in 50 mM phosphate buffer (pH 7.0) on a Shimadzu UV-2600 [UV–Vis](#) (ultraviolet–visible) spectrometer at 25 °C in the presence of two equivalents of Zn(II). Measurements were performed in triplicate at a minimum of five inhibitor concentrations. [Enzyme activities](#) were expressed as units per milligram, where 1 unit is defined as the amount of enzyme that cleaves 1 μ mol of dd,l,l-*N*-succinyl-2,6-diaminopimelic acid at 25 °C in 1 min. These data provide a [dose–response curve](#) that allows for the concentration required for 50% [inhibition](#) (IC₅₀) of DapE to be extracted.

The ArgE from *Escherichia coli* (EcArgE) was purified as described previously [30]. IC₅₀ values were determined spectrophotometrically using 2 mM *N* ^{α} -acetyl-l-ornithine (L-NAO) as the substrate. The rate of L-NAO [deacetylation](#) was monitored as a decrease in absorbance at 214 nm using a Shimadzu UV-2600 [UV–Vis](#) spectrophotometer at 25 °C. All kinetic experiments were performed in 50 mM Chelex-100-treated [sodium phosphate](#) buffer at pH 7.5 and 25 °C. Initial rates of L-NAO [deacetylation](#) were determined in triplicate at a minimum of five inhibitor concentrations, providing a dose–response curve from which the concentration required for 50% inhibition (IC₅₀) of ArgE could be extracted. Catalytic activities were determined with an [error](#) of $\pm 10\%$.

Results and discussion

IR and NMR spectroscopies

The structural features of the DAP derivatives were characterized by [IR and NMR spectroscopies](#). IR (KBr, ν_{max} cm^{-1}) [spectra](#) of each of the DAP derivatives reveal characteristic bands corresponding to the functional groups present in these compounds (see [Table SI-1 in the online supplementary material](#))—that is, broad bands in the range $\nu(3427\text{--}3050 \text{ cm}^{-1})$ due to [amino groups](#) and/or carboxylic OH groups, relatively sharp bands in the $\nu(1735\text{--}1726 \text{ cm}^{-1})$ range due to carboxylic C=O groups, and bands in the $\nu(1670\text{--}1630 \text{ cm}^{-1})$ and $\nu(1550\text{--}1522 \text{ cm}^{-1})$ ranges due to [amide I](#) and [amide II](#) groups, respectively. The [IR spectra](#) also show distinct C=O and/or C—F bands at $\nu \sim 1200 \text{ cm}^{-1}$ due to [trifluoroacetate counterions](#), which originate from the HPLC [purification](#) of DAP derivatives where 0.05% (v/v) [trifluoroacetic acid](#) was used as an acid modifier of the [mobile phase](#).

The ¹H and ¹³C [chemical shift](#) data for each DAP derivative were assigned according to the following general formula:

where **R** indicates the various acyl substituents (see [Tables SI-2 and SI-3 in supplementary material](#)). Signals due to minor [isomers](#) (if resolved from a major isomer) are in italics. The series of DAP derivatives examined were prepared from an enantiomeric/diastereomeric mixture of *RS,RS*-2,6-diaminopimelic acid, and the mixtures of two diastereoisomeric forms were also detected. Due to the additional chiral center on the acyl substituent **R** in compounds **C2** and **E3**, ratios of four [isomers](#) without any ascribed priority were estimated.

Capillary electromigration methods

Selection of experimental conditions

The DAP derivatives examined ([Table 1](#)) are typical amphoteric compounds containing both acidic (carboxylic) and basic (amino) groups with different acyl substituents **R** on one of the two amino groups. Hence, the experimental conditions for CZE and MEKC analyses and separations of DAP derivatives were selected based on the general rules for suitable [capillary electrophoresis](#) separation conditions for amino acids and [oligopeptides](#) [31], [32], [33], [34], [35] while also taking into account the specific properties of the DAP derivatives. The selection of BGEs includes the type and concentration of [buffer](#) components and [pH](#), and it also takes into account the [solubility](#) and [chemical stability](#) of each compound. Effective [mobility](#) of weak electrolytes and [ampholytes](#) strongly depends on pH; therefore, pH is one of the most important parameters influencing separation efficiency and selectivity of the BGEs.

For CZE analysis and separation of amphoteric DAP derivatives, either acidic or alkaline BGEs are suitable. Therefore, the following BGEs were used: (i) acidic BGEs (pH 2.18–2.54) in which the amino groups are fully protonated and the carboxylic groups are only partially dissociated, resulting in cationic migration of the DAP derivatives; and (ii) basic BGEs (pH 8.20–8.64) in which the amino groups are partially protonated and the carboxylic groups are fully dissociated, resulting in anionic migration of the DAP derivatives. BGE compositions were also selected with respect to their buffering capacity in order to maintain proper pH in the zone of migrating analytes (i.e., pH of the BGE was close to the pK_a value of the buffering constituent of the BGE). With the addition of anionic (SDS) or cationic (CTAB) [detergents](#), the same BGEs were also employed for the separation of DAP derivatives by MEKC. The complete composition and pH of the BGEs used, as well as the separation [voltages](#) and [electric currents](#) employed in the CZE and MEKC analyses and separations of DAP derivatives, are presented in [Table 2](#).

Table 2. Background electrolytes, separation [voltage](#) (U_{sep}), and [electric current](#) (I) used in CZE and MEKC methods.

BGE number	BGE composition	pH	Method	U_{sep} (kV)	I (μ A)
BGE I	500 mM AcOH	2.54	CZE	12.0	7.1
BGE II	100 mM Tris, 50 mM AcOH	8.20	CZE	12.0	23.3
BGE III	20 mM Tris, 5 mM H_3PO_4 , 50 mM SDS	8.64	MEKC	12.0	18.6
BGE IV	200 mM iminodiacetic acid	2.28	CZE	12.0	35.0
BGE V	50 mM Tris, 100 mM H_3PO_4	2.18	CZE	12.0	45.0
BGE VI	500 mM AcOH, 60 mM SDS	2.54	MEKC	–12.0	24.0
BGE VII	100 mM Tris, 50 mM AcOH, 5 mM CTAB	8.20	MEKC	–12.0	23.0
BGE VIII	100 mM Tris, 50 mM AcOH, 10 mM CTAB	8.20	MEKC	–12.0	23.5

Determination of purity of DAP derivatives

DAP derivatives were synthesized for the first time; their standard preparations were not available, and for that reason their purity degree could be quantified only relatively by the relative corrected peak area, P_{CA} [36], defined by the following relation:

$$(1) P_{CA(i)}(\%) = \frac{A_{c(i)}}{\sum_{i=1}^n A_{c(i)}} \cdot 100$$

where $A_{c(i)}$ is the corrected peak area of the i -th component of the given preparation and n is the number of components of this preparation. Corrected peak area is the peak area normalized (divided) by the migration time of this peak. This way, defined purity degree is only an approximate measure of analyte purity given that molar [absorption](#) coefficients of the individual components may differ. Nevertheless, with respect to the

expected structural similarity of the admixtures and main synthetic product, it is reasonable to assume similar values of their molar absorption coefficients and, thus, also a relatively good estimation of the purity degree of the main product.

Analyses and purity degree determinations were performed by CZE in the acidic BGE I and alkaline BGE II and by MEKC in the alkaline BGE III. The P_{CA} for each DAP derivative is summarized in [Table 3](#). For all analytes, high degrees of purity were observed with P_{CA} values in the range 95.0 to ~100%. For comparison purposes, CE analyses for the DAP derivatives **E1** and **C3** in the electrolyte systems BGE I, BGE II, and BGE III are discussed further. The derivative **E1** was found to be highly pure ($P_{CA} \sim 100\%$) in all three BGEs ([Figs. 2A–C](#) and [Table 3](#)). For the DAP derivative **C3**, one nonidentified admixture (peak x) with lower [electrophoretic mobility](#) than that of the analyte **C3** was found in BGE I and BGE II, and its purity was found to be 94.8% in BGE I and 94.1% in BGE II ([Figs. 2D and 2E](#) and [Table 3](#)). However, MEKC performed in BGE III revealed three nonidentified minor admixtures—x1, x2, and x3—and its purity was found to be 90.8% ([Fig. 2F](#) and [Table 3](#)). These and other examples show that significant differences in purity were detected, depending on the method (CZE or MEKC) and BGE used. CZE analyses on analyte **C1** in BGEs I and II suggest high purity ($P_{CA} \sim 100\%$), but MEKC in BGE III revealed two minor admixtures x1 and x2 and purity of 94.5% (see [Figs. SI-1A–C in supplementary material](#)). The analyte **C4** exhibits high purity ($P_{CA} \sim 100\%$) when analyzed as a [cation](#) by CZE in acidic BGE I, but CZE in alkaline BGE II and MEKC in alkaline BGE III with SDS pseudostationary phase revealed one or two minor admixtures indicating lower purity—97.1 and 95.0%, respectively ([Figs. SI-1D–F](#)). Thus, MEKC in BGE III with SDS seems to be more efficient for analysis of DAP derivatives than CZE because it was able to separate admixtures, which remained unrevealed by CZE analyses in BGEs I and II.

Table 3. Purity degree (P_{CA}) of DAP and its mono-*N*-acyl-DAP derivatives estimated by CZE in BGE I and BGE II and by MEKC in BGE III.

P_{CA} (%)			
Compound	BGE I	BGE II	BGE III
	pH 2.54	pH 8.20	pH 8.64
DAP	~100	~100	~100
A1	94.5	93.6	93.7
A2	95.8	95.4	96.1
A3	96.6	97.1	97.0
A4	94.3	95.0	95.8
A5	95.6	96.5	95.6
A6	95.2	95.4	96.2
C1	~100	~100	94.5
C2	96.9	98.4	93.2
C3	94.8	94.1	90.8
C4	~100	97.1	95.0
E1	~100	~100	~100
E2	94.3	91.4	95.1
E3	94.4	93.4	95.9

Note. For [names](#) and structures of the compounds, see [Table 1](#). For composition of the BGEs, see [Table 2](#).

Fig.2. Analyses of DAP derivatives **E1** and **C3** by CZE in BGE I and BGE II and by MEKC in BGE III: (A) **E1** by CZE in BGE I; (B) **E1** by CZE in BGE II; (C) **E1** by MEKC in BGE III; (D) **C3** by CZE in BGE I; (E) **C3** by CZE in BGE II; (F) **C3** by MEKC in BGE III. [Sample](#) injection: [hydrodynamic](#), 10 mbar $\times$ 5 s, x, x1, x2, x3, nonidentified admixtures. For DAP derivative [names](#) and structures, see [Table 1](#). The BGE composition and other experimental conditions are given in [Table 2](#) and in “Materials and Methods”.

Characterization of DAP derivatives

DAP derivatives were also characterized by their effective electrophoretic mobilities in eight BGEs. The effective [electrophoretic mobility](#) at the [temperature](#) inside the capillary, $m_{\text{eff},T}$, was determined from the CZE and MEKC experimental data using the Eq. (2):

$$(2) m_{\text{eff},T} = \left(\frac{1}{t_{\text{mig}}} - \frac{1}{t_{\text{eof}}} \right) \cdot \frac{L_{\text{eff}} \cdot L_{\text{tot}}}{U_{\text{sep}}}$$

where t_{mig} is the migration time of the given analyte (s), t_{eof} is the migration time of the [electroosmotic flow marker](#) (s), L_{eff} and L_{tot} are the effective and total capillary [lengths](#) (m), respectively, and U_{sep} is the applied separation voltage (V). CZE and MEKC experiments were performed at different [ambient temperatures](#) (22–26 °C) and variable input power (Joule heat). To compare the observed mobilities, the actual mobility at temperature T inside the capillary was recalculated to a reference mobility at 25 °C, $m_{\text{eff},25}$, using Eq. (3):

$$(3) m_{\text{eff},25} = m_{\text{eff},T} [1 - 0.020(T - 25)]$$

The actual temperature inside the capillary for each BGE was obtained based on the experimentally measured dependence of temperature increase inside the capillary on the input power (Joule heat) per unit capillary length as described previously [\[37\]](#).

The values for effective electrophoretic mobilities of DAP derivatives in eight BGEs at 25 °C are presented in [Table 4](#). These data exhibit a strong dependence of mobilities of DAP derivatives on pH and reflect the presence of the particular ionogenic groups and the ratio of effective charge versus size (M_r) of these compounds. Cationic effective electrophoretic mobilities determined by CZE in the acidic BGE I (0.5 M AcOH, pH 2.54) were relatively low at approximately 5.0 MU (MU = mobility unit = $1 \times 10^{-9} \text{ m}^2 \text{ V}^{-1} \text{ s}^{-1}$) due to compensation of the positive charge of the amino group of DAP derivatives by the partially negative charges of their two or three carboxyl groups. Interestingly, in the **C4** DAP derivative, four [fluorine atoms](#) in the side chain significantly increased the [acidity](#) of the corresponding carboxyl group, resulting in a prevailing anionic form of this compound with negative effective mobility even in the acidic BGE I (pH 2.54). In CZE analyses in alkaline BGE II (100 mM [Tris](#) and 50 mM [acetic acid](#), pH 8.20), all analyzed DAP derivatives migrated as [anions](#). The derivatives **A1** to **A6** and **E1** to **E3** exhibited similar moderate negative effective mobilities in the range –13.6 to –15.7 MU due to similar ratios of their effective charges and [molecular masses](#). The derivatives **C1** to **C4** containing the *N*-acyl substituent with additional free carboxyl groups possessed an increased negative charge and exhibited enhanced anionic mobilities in the range –23.3 to –25.2 MU for compounds **C1** to **C3** and up to –30.4 MU for the most strongly negatively charged derivative **C4**.

Table 4. Effective [electrophoretic mobilities](#) at 25 °C of DAP derivatives determined by CZE in BGEs I, II, IV, and V and by MEKC in BGEs III, VI, VII, and VIII.

	$m_{\text{eff},25} \times 10^9 \text{ (m}^2 \text{ V}^{-1} \text{ s}^{-1}\text{)}$							
DAP derivative	BGE I	BGE II	BGE III	BGE IV	BGE V	BGE VI	BGE VII	BGE VIII
	pH 2.54	pH 8.20	pH 8.64	pH 2.28	pH 2.18	pH 2.54	pH 8.20	pH 8.20
A1	4.7	−15.7	−14.2	7.3	8.5 ^a	−9.2 ^a	−14.2	−13.0
					8.3 ^a	−9.1 ^a		
A2	5.0	−14.8	−13.6	7.0	7.8 ^a	−16.6 ^a	−11.9	−10.8
					7.7 ^a	−16.5 ^a		
A3	4.8	−14.6	−13.2	7.0	7.9	−17.8 ^a	−11.8	−10.0
						−17.6 ^a		
A4	5.0	−14.5	−12.7	6.9	7.9	−21.7	−9.3	−6.8
					7.8			
A5	4.7	−13.9	−12.2	6.7	7.6	−23.1 ^a	−8.7	−6.1
						−23.0 ^a		
A6	4.4	−13.9	−11.4	6.6	8.1	−25.7 ^a	−1.2 ^a	4.9
						−25.6 ^a	−0.7 ^a	
C1	4.1	−25.2	−23.2	6.3	7.0	−10.6	−15.3	−10.5
C2	4.3	−24.0	−23.6	6.3	7.4	−14.0 ^a	−12.3	−7.4
						−15.9 ^a		
C3	4.3	−23.3	−22.9	6.2	7.4	−20.6	−8.3	~0
C4	−2.4	−30.4	−29.6	~0	−10.3	−12.9	28.0	28.6
E1	3.5	−14.0	−11.1	6.0	7.3	−16.0 ^a	−9.4	−7.5
						−15.8 ^a		
E2	4.1	−14.3	−12.1	6.7	7.7 ^a	−19.2	−11.9	−9.4
					7.6 ^a			
E3	4.9	−13.6	−11.7	6.9	7.9	−19.2 ^a	−11.5	−9.1
						−18.9 ^a		

Note. For DAP derivative [names](#) and structures, see [Table 1](#). The BGE composition and other experimental conditions are given in [Table 2](#) and in Materials and Methods.

^aEffective mobility of resolved diastereomer of DAP derivative.

Similar electromigration [behavior](#) was observed in the MEKC analyses of DAP derivatives in BGE III (20 mM Tris, 5 mM H₃PO₄, and 50 mM SDS, pH 8.64), where derivatives **A1** to **A6** and **E1** to **E3** exhibited anionic mobilities in the range −11.1 to −14.2 MU and the mobilities of the compounds **C1** to **C4** achieved values of −22.9 to −29.6 MU. The small observed difference between mobilities in the micelle-free BGE II and those in the micelle-containing BGE III suggests that the DAP derivatives do not strongly interact with the SDS micellar pseudophase of BGE III.

In the CZE analyses of DAP derivatives in the isoelectric BGE IV (200 mM IDA, pH 2.28) and in the classical BGE V (100 mM H₃PO₄ and 50 mM Tris, pH 2.18), similar structure–mobility relationships of these compounds were observed as in the BGE I. Slightly higher cationic mobilities were observed in BGEs IV and V (~6.0–8.0 MU) than in BGE I (4.0–5.0 MU), which is due to the increased positive charge of the DAP derivatives at lower pH in BGEs IV and V (pH 2.28 and 2.18) than in BGE I (pH 2.54). The exception to this behavior is compound **C4**, which exhibited negative mobility even under strongly [acidic conditions](#) in BGEs IV and V.

An interaction between the positively charged DAP derivatives and the negatively charged SDS [micelles](#) in acidic BGE VI (0.5 M AcOH, pH 2.54) resulted in a negatively charged analyte–micelle complex with variable negative mobilities in the range –9.1 to –25.7 MU for compounds **A1** to **A6** and **E1** to **E3** and in the range –10.6 to –20.6 MU for compounds **C1** to **C4**. In this case, the differences in mobilities did not result from differences in the charge/size ratio but rather resulted from differences in the [hydrophobicities](#) of the analytes and from the binding [strength](#) of their complexes with SDS [micelles](#).

Negative effective charge and negative mobility were observed for nearly all DAP derivatives (with the exception of **A6** and **C4**) in alkaline media such as BGE VII (100 mM Tris, 50 mM AcOH, and 5 mM CTAB, pH 8.2) and BGE VIII (100 mM Tris, 50 mM AcOH, and 10 mM CTAB, pH 8.2), although diminished due to their weak interaction with positively charged CTAB micelles. Slightly lower absolute values of mobilities in BGE VIII as compared with BGE VII are apparently caused by the higher concentration of CTAB in BGE VIII (10 mM) versus BGE VII (5 mM). Reverse migration behavior was observed for DAP derivatives containing the aromatic group (**A6**) in BGE VIII and the acidic DAP derivative **C4** in BGE VII and BGE VIII. These two derivatives form strong complexes with cationic CTAB micelles, resulting in a positive charge and cationic migration with a low positive mobility (4.9 MU) for **A6** and a high positive mobility (28.0–28.6 MU) for **C4**. It is worth noting that in some cases the [diastereomers](#) of the DAP derivatives were partially or completely resolved in achiral media—by CZE in BGE V and by MEKC in BGE VI and BGE VII (see [Fig.3](#), [Fig.4](#) in the next section). Thus, these separations allowed determination of effective electrophoretic mobilities for both diastereomers (enantiomer pairs) of these compounds (see [Table 4](#)).

Fig.3. MEKC separation of partial mixtures of DAP derivatives in BGE VI: (A) mixture M1 (**A1–A3**); (B) mixture M2 (**A4–A6**); (C) mixture M3 (**E1–E3**); (D) mixture M4 (**C1–C4**). [Sample](#) injection: 10 mbar × 5 s, x, nonidentified admixtures. For DAP derivative [names](#) and structures, see [Table 1](#). The BGE composition and other experimental conditions are given in [Table 2](#) and in “Materials and Methods”.

Fig.4. MEKC separation of the 11 DAP derivatives in BGE VI. [Sample](#) injection: 10 mbar $\times$ 5 s, x, nonidentified admixtures. For DAP derivative [names](#) and structures, see [Table 1](#). The BGE composition and other experimental conditions are given in [Table 2](#) and in “Materials and Methods”.

Separation of DAP derivatives

From the eight BGEs tested, the largest differences in electrophoretic mobilities for DAP derivatives were found in BGE VI (0.5 M AcOH [pH 2.54] and 60 mM SDS). Consequently, the separation of mixtures of DAP derivatives was carried out by MEKC in this BGE. First, four partial mixtures of the analytes were constituted according to similarity of their *N*-acyl substituent ([Table 1](#)). The mixture M1 was constituted from compounds **A1** to **A3** with linear and branched C2 and C4 aliphatic [acyl groups](#); mixture M2 from compounds **A4** and **A5** with linear and branched C5 aliphatic acyl groups, respectively, and compound **A6** with an aromatic acyl group; mixture M3 from compounds **E1** to **E3** containing esterified carboxyl groups; and mixture M4 composed of compounds **C1** to **C4**, which contain free carboxylic groups. [Fig. 3](#) shows complete MEKC separations for all components of mixtures M1, M2, and M4, including a complete or partial separation of the diastereomers of **A1**, **A2**, **A3**, **A5**, **A6**, **C1**, and **C2**, whereas in mixture M3 a baseline separation of the diastereomers of **E1** was achieved but **E2** and **E3** and their diastereomers comigrated. The addition of organic modifiers (methanol or acetonitrile) to the BGE did not improve the separation of these two compounds and their diastereomers. Finally, using this optimized BGE, a mixture of all DAP derivatives, with the exception of comigrating derivatives **E2** and **E3**, was separated by MEKC within a short time of 11 min ([Fig. 4](#)).

Enzyme inhibition

All DAP derivatives were examined for their ability to inhibit the [catalytic activity](#) of *HiDapE* enzyme. Of the 13 compounds tested, only **A1** showed a weak [inhibition](#) of the catalytic activity with an IC_{50} value of 17 mM. These data suggest that the amino moiety of mono-*N*-succinyl-DAP forms an important interaction within the [active site](#) of DapE given that the acylated amino acids did not inhibit [enzymatic activity](#). Alternatively, the acylated versions of these amino acids may introduce a repulsive steric interaction because of bulky [alkyl groups](#), thereby preventing [enzyme binding](#). All of the DAP derivatives were also examined for their ability to inhibit *EcArgE*. Some of the DAP derivatives exhibited a weak inhibition toward *EcArgE*; the best inhibitor was **E1**, with an IC_{50} value of 480 μ M, thereby providing a promising lead structure for further studies on ArgE inhibitors. The other DapE inhibitors that showed a dose-dependent response to *EcArgE* were **A1**, **C1**, and **C3**, with IC_{50} values of 2.63, 2.11, and 1.28 mM, respectively, whereas the remaining DAP derivatives did not inhibit *EcArgE*. It implies that both DapE and ArgE active sites possess strict [substrate specificity](#) with regard to both functional groups and [stereochemistry](#).

Conclusions

The data presented here proved that [IR and NMR spectroscopies](#) and capillary electromigration methods, CZE and MEKC, represent powerful tools for structural analysis, purity determination, separation, and physicochemical characterization of amino acids and their derivatives. Structural features of mono-*N*-acyl derivatives of 2,6-DAP were characterized by IR and [NMR spectroscopies](#), whereas purity degree and effective [electrophoretic mobilities](#) of cationic and anionic forms of these compounds in acidic and alkaline classical or isoelectric BGEs with and without anionic (SDS) or cationic (CTAB) [detergents](#) were determined by CZE and MEKC methods. The best separation of DAP derivatives, including [diastereomers](#) of some of them, was achieved by MEKC in an acidic BGE (500 mM [acetic acid](#), [pH](#) 2.54) with micellar pseudophase constituted by anionic detergent, 60 mM SDS. None of the analyzed DAP derivatives worked as an effective inhibitor of [catalytic activity](#) of the DapE from *H. influenza*, but one derivative—*N*-fumaryl, Me-ester-DAP—was found to be a moderate inhibitor of ArgE from *E. coli*, thereby providing a promising lead structure for further studies on ArgE inhibitors.

Acknowledgments

This work was supported by the Czech Science Foundation ([P206/12/0453](#) and [13-17224S](#)), the Grant Agency of the Academy of Sciences of the Czech Republic ([IAA400550614](#)), the Academy of Sciences of the Czech Republic ([RVO 61388963](#)), and the National Institutes of Health ([R15 AI085559-01A1](#) to R.C.H.). S. Vašíčková is thanked for measurement of [IR spectra](#).

References

- [1] W.J. Hierholzer, J.S. Garner, A.B. Adams, D.E. Craven, D.W. Fleming, S.W. Forlenza, M.J. Gilchrist, D.A. Goldman, E.L. Larson, C.G. Mayhall, R.D. McCormick, R.L. Nichols, C.E. Edmiston, D.G. Maki, G. Pugliese, R.A. Weinstein, O.C. Tablan, F.C. Tenover, W.J. Martone, R.F. Gaynes, W.R. Jarvis, M.F. Favero **Recommendations for preventing the spread of vancomycin resistance: recommendations of the Hospital Infection Control Practices Advisory Committee (HICPAC)** *Am. J. Infect. Control*, 23 (1995), pp. 87-94
- [2] R.E.W. Hancock **Peptide antibiotics** *Lancet*, 349 (1997), pp. 418-422
- [3] R.A. Howe, K.E. Bowker, T.R. Walsh, T.G. Feest, A.P. MacGowan **Vancomycin-resistant *Staphylococcus aureus*** *Lancet*, 351 (1998), p. 602
- [4] S.B. Levy **The challenge of antibiotic resistance** *Sci. Am.*, 278 (1998), pp. 46-53
- [5] M. Teuber **Spread of antibiotic resistance with food-borne pathogens** *Cell. Mol. Life Sci.*, 56 (1999), pp. 755-763
- [6] P.J. Dolin, M.C. Raviglione, A. Kochi **Global tuberculosis incidence and mortality during 1990–2000** *Bull. WHO*, 72 (1994), pp. 213-220
- [7] M.C. Raviglione, D.E. Snider, A. Kochi **Global epidemiology of tuberculosis: morbidity and mortality of a worldwide epidemic** *JAMA*, 273 (1995), pp. 220-226
- [8] J.R. Miller, S. Dunham, I. Mochalkin, C. Banotai, M. Bowman, S. Buist, B. Dunkle, D. Hanna, H.J. Harwood, M.D. Huband, A. Karnovsky, M. Kuhn, C. Limberakis, J.Y. Liu, S. Mehrens, W.T. Mueller, L. Narasimhan, A. Ogden, J. Ohren, J.V.N.V. Prasad, J.A. Shelly, L. Skerlos, M. Sulavik, V.H. Thomas, S. VanderRoest, L.A. Wang, Z. G. Wang, A. Whitton, T. Zhu, C.K. Stover **A class of selective antibacterials derived from a protein kinase inhibitor pharmacophore** *Proc. Natl. Acad. Sci. U.S.A.*, 106 (2009), pp. 1737-1742
- [9] L. Monincová, M. Buděšínský, J. Slaninová, O. Hovorka, J. Cvačka, Z. Voburka, V. Fučík, L. Borovičková, L. Bednarová, J. Straka, V. Čerovský **Novel antimicrobial peptides from the venom of the eusocial bee *Halictus sexcinctus* (Hymenoptera: Halictidae) and their analogs** *Amino Acids*, 39 (2010), pp. 763-775
- [10] L. Monincová, J. Slaninová, V. Fučík, O. Hovorka, Z. Voburka, L. Bednarová, P. Maloň, J. Štokrová, V. Čerovský **Lasiocepsin, a novel cyclic antimicrobial peptide from the venom of eusocial bee *Lasioglossum laticeps* (Hymenoptera: Halictidae)** *Amino Acids*, 43 (2012), pp. 751-761
- [11] P. Niederhafner, L. Bednarová, M. Buděšínský, M. Šafařík, S. Ehala, J. Ježek, L. Borovičková, V. Fučík, V. Čerovský, J. Slaninová **Melectin MAPs: the influence of dendrimerization on antimicrobial and hemolytic activity** *Amino Acids*, 39 (2010), pp. 1553-1561
- [12] J.Y. Kim, S.C. Park, M.Y. Yoon, K.S. Hahm, Y. Park **C-terminal amidation of PMAP-23: translocation to the inner membrane of gram-negative bacteria** *Amino Acids*, 40 (2011), pp. 183-195
- [13] J. Hlaváček, J. Pícha, V. Vaněk, J. Jiráček, J. Slaninová, V. Fučík, M. Buděšínský, D. Gilner, R.C. Holz **Inhibitors of N^α-acetyl-L-ornithine deacetylase: synthesis, characterization, and analysis of their inhibitory potency** *Amino Acids*, 38 (2010), pp. 1155-1164
- [14] G. Scapin, J.S. Blanchard **Enzymology of bacterial lysine biosynthesis** *Adv. Enzymol.*, 72 (1998), pp. 279-324
- [15] T.L. Born, J.S. Blanchard **Structure/function studies on enzymes in the diaminopimelate pathway of bacterial cell wall biosynthesis** *Curr. Opin. Chem. Biol.*, 3 (1999), pp. 607-613
- [16] J.M. Girodeau, C. Agouridas, M. Masson, R. Pineau, F. Legoffic **The lysine pathway as a target for a new genera of synthetic antibacterial antibiotics** *J. Med. Chem.*, 29 (1986), pp. 1023-1030
- [17] D.M. Gillner, D.P. Becker, R.C. Holz **Lysine biosynthesis in bacteria: a metallodesuccinylase as a potential antimicrobial target** *J. Biol. Inorg. Chem.*, 18 (2013), pp. 155-163

- [18] D.L. Bienvenue, D.M. Gilner, R.S. Davis, B. Bennett, R.C. Holz **Substrate specificity, metal binding properties, and spectroscopic characterization of the DapE-encoded N-succinyl-L, L-diaminopimelic acid desuccinylase from *Haemophilus influenzae*** *Biochemistry*, 42 (2003), pp. 10756-10763
- [19] M. Karita, M.L. Etterbeek, M.H. Forsyth, M.K.R. Tummuru, M.J. Blaser **Characterization of *Helicobacter pylori* dapE and construction of a conditionally lethal dapE mutant** *Infect. Immun.*, 65 (1997), pp. 4158-4164
- [20] M.S. Pavelka, W.R. Jacobs **Biosynthesis of diaminopimelate, the precursor of lysine and a component of peptidoglycan, is an essential function of *Mycobacterium smegmatis*** *J. Bacteriol.*, 178 (1996), pp. 6496-6507
- [21] C.A. Hutton, M.A. Perugini, J.A. Gerrard **Inhibition of lysine biosynthesis: an evolving antibiotic strategy** *Mol. BioSyst.*, 3 (2007), pp. 458-465
- [22] V. Vaněk, J. Pícha, M. Buděšínský, M. Šanda, J. Jiráček, R.C. Holz, J. Hlaváček **Synthesis of N-succinyl-L, L-diaminopimelic acid mimetics via selective protection** *Protein Pept. Lett.*, 17 (2010), pp. 405-409
- [23] M. Geiger, A.L. Hogerton, M.T. Bowser **Capillary electrophoresis** *Anal. Chem.*, 84 (2012), pp. 577-596
- [24] M. Silva **Micellar electrokinetic chromatography: a practical overview of current methodological and instrumental advances** *Electrophoresis*, 32 (2011), pp. 149-165
- [25] M. Silva **Micellar electrokinetic chromatography: a review of methodological and instrumental innovations focusing on practical aspects** *Electrophoresis*, 34 (2013), pp. 141-158
- [26] V. Poinso, M.A. Carpena, J. Bouajila, P. Gavard, B. Feurer, F. Couderc **Recent advances in amino acid analysis by capillary electrophoresis** *Electrophoresis*, 33 (2012), pp. 14-35
- [27] V. Poinso, V. Ong-Meang, P. Gavard, F. Couderc **Recent advances in amino acid analysis by capillary electromigration methods, 2011–2013** *Electrophoresis*, 35 (2014), pp. 50-68
- [28] C. Desiderio, F. Iavarone, D.V. Rossetti, I. Messina, M. Castagnola **Capillary electrophoresis–mass spectrometry for the analysis of amino acids** *J. Sep. Sci.*, 33 (2010), pp. 2385-2393
- [29] V. Kašička, Z. Prusík, P. Sázelová, M. Chiari, I. Mikšík, Z. Deyl **External electric field control of electroosmotic flow in non-coated and coated fused-silica capillaries and its application for capillary electrophoretic separations of peptides** *J. Chromatogr. B*, 741 (2000), pp. 43-54
- [30] W.C. McGregor, S.I. Swierczek, B. Bennett, R.C. Holz **ArgE-encoded N-acetyl-L-ornithine deacetylase from *Escherichia coli* contains a dinuclear metalloactive site** *J. Am. Chem. Soc.*, 127 (2005), pp. 14100-14107
- [31] G.M. Janini, H.J. Issaq **Selection of buffers in capillary zone electrophoresis: application to peptide and protein analysis** *Chromatographia*, 53 (2001), pp. S18-S26
- [32] V. Kašička **Capillary electrophoresis of peptides** *Electrophoresis*, 20 (1999), pp. 3084-3105
- [33] G.K.E. Scriba, A. Psurek **Separation of peptides by capillary electrophoresis** P. Schmitt-Kopplin (Ed.), *Capillary Electrophoresis*, Humana, Totowa, NJ (2008), pp. 483-506
- [34] V. Kašička **Recent developments in CE and CEC of peptides [2009–2011]** *Electrophoresis*, 33 (2012), pp. 48-73
- [35] P.G. Righetti, R. Sebastiano, A. Citterio **Capillary electrophoresis and isoelectric focusing in peptide and protein analysis** *Proteomics*, 13 (2013), pp. 325-340
- [36] V. Šolínová, V. Kašička, D. Koval, T. Barth, A. Ciencialová, L. Žáková **Analysis of synthetic derivatives of peptide hormones by capillary zone electrophoresis and micellar electrokinetic chromatography with ultraviolet-absorption and laser-induced fluorescence detection** *J. Chromatogr. B*, 808 (2004), pp. 75-82
- [37] D. Koval, V. Kašička, J. Jiráček, M. Collinsova, T.A. Garrow **Determination of dissociation constant of phosphinate group in phosphinic pseudopeptides by capillary zone electrophoresis** *J. Chromatogr. B*, 770 (2002), pp. 145-154

⁴These authors contributed equally to this article.

²[Abbreviations used:](#) ArgE, *N*^α-acetyl-L-ornithine deacetylase; mDAP, *meso*-diaminopimelate or *meso*-diaminopimelic acid; DapE, dapE-encoded *N*-succinyl-L,L-diaminopimelic acid desuccinylase; DAP, 2,6-diaminopimelic acid; HPLC, high-performance liquid chromatography; IR, infrared; NMR, nuclear magnetic resonance; CZE, capillary zone electrophoresis; MEKC, micellar electrokinetic chromatography; BGE, background electrolyte; SDS, sodium dodecyl sulfate; CTAB, cetyltrimethylammonium bromide; Tris, tris(hydroxymethyl)aminomethane; H₃PO₄, phosphoric acid; AcOH, acetic acid; NaOH, sodium hydroxide; IDA, iminodiacetic acid; ACN, acetonitrile; MeOH, methanol; FT, Fourier transform; UV, ultraviolet; *Hi*DapE, DapE from *Haemophilus influenzae*; UV–Vis, ultraviolet–visible; *Ec*ArgE, ArgE from *Escherichia coli*; L-NAO, *N*^α-acetyl-L-ornithine; MU, mobility units.

Appendix A. Supplementary data

[Download Word document \(5MB\)](#)[Help with doc files](#)

Supplementary data 1.