Marquette University

e-Publications@Marquette

Biological Sciences Faculty Research and
Publications

Biological Sciences, Department of

11-1998

Force-velocity-power and Force-pCa Relationships of Human Soleus Fibers After 17 Days of Bed Rest

Jeffrey J. Widrick
Marquette University

Kris M. Norenberg
Marquette University

C. A. Blaser
Marquette University

M. Karhanek
Marquette University

Jennifer J. Sherwood
Marquette University

See next page for additional authors

Follow this and additional works at: https://epublications.marquette.edu/bio_fac

Part of the [Biology Commons](#)

Recommended Citation

Widrick, Jeffrey J.; Norenberg, Kris M.; Blaser, C. A.; Karhanek, M.; Sherwood, Jennifer J.; Trappe, Scott W.; Trappe, Todd A.; Costill, David L.; and Fitts, Robert H., "Force-velocity-power and Force-pCa Relationships of Human Soleus Fibers After 17 Days of Bed Rest" (1998). *Biological Sciences Faculty Research and Publications*. 448.

https://epublications.marquette.edu/bio_fac/448

Authors

Jeffrey J. Widrick, Kris M. Norenberg, C. A. Blaser, M. Karhanek, Jennifer J. Sherwood, Scott W. Trappe, Todd A. Trappe, David L. Costill, and Robert H. Fitts

Marquette University

e-Publications@Marquette

Department of Biology Faculty Research and Publications/College of Health Sciences

This paper is NOT THE PUBLISHED VERSION; but the author's final, peer-reviewed manuscript. The published version may be accessed by following the link in the citation below.

Journal of Applied Physiology, Vol. 85, No. 5 (November, 1998): 1949-1956. [DOI](#). This article is © American Physiological Society and permission has been granted for this version to appear in [e-Publications@Marquette](#). American Physiological Society does not grant permission for this article to be further copied/distributed or hosted elsewhere without the express permission from American Physiological Society.

Force-velocity-power and force-pCa relationships of human soleus fibers after 17 days of bed rest

J. J. Widrick

Department of Biology, Marquette University, Milwaukee, Wisconsin

K. M. Norenberg

Department of Biology, Marquette University, Milwaukee, Wisconsin

J. G. Romatowski

Department of Biology, Marquette University, Milwaukee, Wisconsin

C. A. Blaser

Department of Biology, Marquette University, Milwaukee, Wisconsin

M. Karhanek

Department of Biology, Marquette University, Milwaukee, Wisconsin

J. Sherwood

Department of Biology, Marquette University, Milwaukee, Wisconsin

S. W. Trappe

Human Performance Laboratory, Ball State University, Muncie, Indiana

T. A. Trappe

Human Performance Laboratory, Ball State University, Muncie, Indiana

D. L. Costill

Human Performance laborator, Ball State University, Muncie, Indiana

R. H. Fitts

Department of Biology, Marquette University, Milwaukee, Wisconsin

Abstract

Soleus muscle fibers from the rat display a reduction in peak power and Ca^{2+} sensitivity after hindlimb suspension. To examine human responses to non-weight bearing, we obtained soleus biopsies from eight adult men before and immediately after 17 days of bed rest (BR). Single chemically skinned fibers were mounted between a force transducer and a servo-controlled position motor and activated with maximal (isotonic properties) and/or submaximal (Ca^{2+} sensitivity) levels of free Ca^{2+} . Gel electrophoresis indicated that all pre- and post-BR fibers expressed type I myosin heavy chain. Post-BR fibers obtained from one subject displayed increases in peak power and Ca^{2+} sensitivity. In contrast, post-BR fibers obtained from the seven remaining subjects showed an average 11% reduction in peak power ($P < 0.05$), with each individual displaying a 7–27% reduction in this variable. Post-BR fibers from these subjects were smaller in diameter and produced 21% less force at the shortening velocity associated with peak power. However, the shortening velocity at peak power output was elevated 13% in the post-BR fibers, which partially compensated for their lower force. Post-BR fibers from these same seven subjects also displayed a reduced sensitivity to free Ca^{2+} ($P < 0.05$). These results indicate that the reduced functional capacity of human lower limb extensor muscles after BR may be in part caused by alterations in the cross-bridge mechanisms of contraction.

WEIGHT-BEARING ACTIVITY is recognized as an important environmental stimulus for maintaining normal neuromuscular function of the soleus and other antigravity extensor muscles. In the absence of hindlimb weight bearing, the rat soleus muscle has been shown to atrophy and to display a reduced peak tetanic tension and power output and an increased percentage of fast-twitch fibers (3, 10). Similar functional changes have been observed for maximally Ca^{2+} -activated single muscle fibers isolated from the rat soleus after hindlimb suspension (1, 11, 20). Additionally, the individual rat fibers displayed a reduced sensitivity to intracellular Ca^{2+} (1, 11, 22). These observations suggest that at least a portion of the change in whole-muscle function after non-weight bearing can be attributed to an altered cellular function coupled with molecular changes in the type of protein expressed.

In humans, voluntary torque of the lower limb extensor muscle group is impaired after spaceflight or prolonged bed rest (5, 15). However, it is not clear to what extent these changes in human neuromuscular performance are due to alterations in neural mechanisms of recruitment, to changes in muscle architecture and/or fiber type composition, and/or to alterations in cellular processes of contraction.

Recently, investigators have used single-fiber in vitro preparations to study human muscle function after non-weight bearing. Larsson et al. (19) found that 42 days of complete bed rest produced large decrements in the force production of single vastus lateralis fibers along with reductions in their unloaded shortening velocity (V_0). After 17 days of bed rest, our laboratory has observed a more modest atrophy of type I soleus fibers with an average shift in V_0 toward higher velocities (30).

Previous human studies have concentrated on isometric and unloaded contractions (19, 30). These are not the conditions present during movement, when muscle fibers must shorten while under an external load. Therefore, the first goal of this study was to determine whether the isotonic contractile properties of human muscle fibers were altered after a 17-day period of bed rest. Our second goal was to characterize the force-pCa relationship (where $pCa = -\log$ of the Ca^{2+} concentration) of human muscle fibers before and after bed rest to determine whether Ca^{2+} sensitivity was altered by non-weight bearing.

METHODS

This study was approved by the Institutional Review Boards at Marquette University, Ball State University, and the National Aeronautics and Space Administration (NASA) Ames Research Center. Eight men were selected from a pool of volunteers. Each subject underwent a medical examination and provided written informed consent before participation. Their mean age, height, and pre-bed-rest body mass were 43 ± 3 (SE) yr, 182 ± 2 cm, and 82.2 ± 4.3 kg, respectively. Each subject was assigned an identification number of 1–8 exactly as described in our previous work (30).

Subjects resided continuously at the Human Research Facility at NASA Ames Research Center (Moffett Field, CA) during a 14-day ambulatory control period, a 17-day 6° head-down-tilt bed-rest period, and an 8-day ambulatory recovery period. This duration of bed rest corresponded with the planned duration of the NASA Life and Microgravity Sciences Spacelab (LMS) space shuttle mission (shuttle transport system-78) scheduled to take place ~ 1 year after this study. Subjects underwent physiological evaluations during the control, bed-rest, and recovery periods as described previously (30). These evaluations were similar to that planned for the LMS mission and included determinations of maximal oxygen consumption by using a supine ergometer and evaluation of right ankle flexor torque by using a specially designed isokinetic dynamometer.

One to three days before the control period, subjects reported to the Human Research Facility, and a pre-bed-rest muscle sample was obtained from the left soleus by using the percutaneous needle-biopsy technique. At the conclusion of bed rest but before reambulation, a post-bed-rest sample was obtained from the right soleus. Pre- and post-bed-rest muscle samples were divided into several portions. The portion used for the present experiments was immediately placed in cold skinning solution [composition in mM: 125 K propionate, 20.0 imidazole (pH 7.0), 2.0 EGTA, 4.0 ATP, and 1.0 $MgCl_2$ and 50% glycerol vol/vol] and was shipped overnight at $4^\circ C$ to Marquette University where it was stored at $-20^\circ C$. All contractile experiments were conducted within 28 days of muscle sampling.

The total and free concentrations of each metal, ligand, and metal-ligand complex in the relaxing and activating solutions were calculated by using an iterative computer program (6). Stability constants used in these calculations (12) were adjusted for the temperature, pH, and ionic strength conditions of the present experiments. Each solution contained (in mM) 7 EGTA, 20 imidazole, 14.5 creatine phosphate, 1 free Mg^{2+} , and 4 free MgATP and sufficient KCl and KOH to produce a total ionic strength of 180 mM and a pH of 7.0. In addition, the relaxing and activating solutions had a free Ca^{2+} concentration of pCa 9.0 and pCa 4.5, respectively.

On the day of an experiment, a single fiber segment was isolated from the muscle bundle and transferred to a small chamber milled into a stainless steel plate. While submerged under relaxing solution, the fiber was mounted between a force transducer (Cambridge model 400, Cambridge Technology, Watertown, MA) and a direct current position motor (Cambridge model 300B, Cambridge Technology). The motor was controlled by a servomechanism similar to that described previously (18). The mounting procedure utilized 4-0 monofilament pins and 10-0 suture to fasten the fiber ends securely into small stainless steel troughs extending from the transducer and motor (31). The stainless steel plate was attached to the stage of an inverted microscope so that the fiber could be viewed at $\times 800$ during data collection. Sarcomere length was adjusted to $2.5 \mu m$ by using an

eyepiece micrometer. The length of the fiber (FL) suspended between the transducer and motor was measured with a micrometer that advanced the plate across the field of view. A Polaroid photograph was taken of the fiber while it was briefly suspended in air (<5 s) by using an Olympus PM-10AD photomicrographic system ($\times 300$). Fiber diameter was determined from the average fiber width measured at three points along the photograph. The fiber diameter in micrometers was measured by using a photograph of an American Optical micrometer (10 μm per division) produced with the same photomicrographic system as the fiber photograph ($\times 300$). Fiber cross-sectional area (CSA) was calculated from fiber diameter under the assumption that the fiber forms a circular cross section when suspended in air (23).

Vertical and horizontal displacements of the stainless steel plate enabled the mounted fiber to be rapidly transferred into an adjacent chamber filled with activating solution. Output from the force transducer and position motor were displayed on a digital oscilloscope before being amplified and interfaced to a personal computer via a LabMaster data-acquisition board. Custom software performed online analysis and stored data to disk. Relaxing and activating solutions were maintained at 15°C during all experiments.

The force-velocity relationship was determined from a series of isotonic releases as described in Fig. 1. P_o was defined as the peak force obtained during the experiment. $V_{\max}$ was defined as the intercept of the force-velocity curve with the velocity axis. Peak absolute power was calculated from P_o , $V_{\max}$, and a/P_o , the parameter that specifies the curvature of the force-velocity relationship (where a is a constant with dimensions of force). For graphical purposes, composite force-velocity-power relationships were plotted by using the mean parameters.

Fig. 1. Force-velocity data obtained from a representative pre-bed-rest type I soleus fiber. *A*: force and position records for 3 consecutive isotonic steps. Fiber attained peak isometric force (P_o) at time point 1. At time point 2, fiber was allowed to shorten as indicated by upward deflection of position record and rapid drop in force. Between time points 2 and 3, rate of fiber shortening was controlled by a servomechanism (18) so that fiber contracted isotonicly for 150 ms. A second and third isotonic step were applied at time points 3 and 4, respectively. At time point 5, fiber was slackened by 20% of its original fiber length, causing force to drop to baseline. Fiber was then returned to relaxing solution and reextended to its original length. *B*: fiber force and shortening velocity were determined over the last one-half of each isotonic step ($\bullet$). Entire procedure was repeated so that each fiber underwent a total of 15–18 randomized isotonic loads. $\circ$, Data points collected during subsequent isotonic contractions; solid line, Hill equation (17) fit to data by using an iterative curve-fitting algorithm. Force-velocity parameters for this fiber were $P_o = 0.95$ mN, parameter that specifies the curvature of the force-velocity relationship (a/P_o) = 0.052, and intercept of force-velocity curve with velocity axis ($V_{\max}$) 0.41 fiber lengths (FL)/s.

A subgroup of fibers were activated with a series of solutions that had free Ca^{2+} concentrations ranging from pCa 6.8 to pCa 5.0. These solutions were made by mixing appropriate volumes of the activating (pCa 4.5) and relaxing (pCa 9.0) solutions. Peak force of each contraction was recorded and expressed as a fraction of the peak force obtained during maximal Ca^{2+} activation. Results for a representative fiber are plotted as in Fig. 2A. The first and approximately every fourth subsequent contraction was performed at pCa 4.5. Hill plots were used to

determine the Ca^{2+} concentration associated with Ca^{2+} activation threshold and with half-maximal activation as described in Fig.2B.

Fig. 2. Representative pCa-force data obtained from a single pre-bed-rest type I soleus fiber (where pCa is $-\log$ of Ca^{2+} concentration). A: pCa-force relationship for this fiber. Peak force for each contraction was expressed as a fraction of maximal Ca^{2+} -activated force, i.e., submaximal force/force at pCa 4.5 (P_r). B: Hill plots of data illustrated in A. Least squares regression lines were fit to data points $<50\%$ of maximal Ca^{2+} activated force ($r^2 = 0.99$; slope = 2.47) and data points $>50\%$ of maximal Ca^{2+} activated force ($r^2 = 0.99$; slope = 1.29). Activation threshold was defined as Ca^{2+} concentration where $\log P_r / (1 - P_r) = -2.5$. Half-maximal activation was calculated as mean intercept of least squares regression lines with the line $y = 0$. In this example, activation threshold was pCa 7.04 with half-maximal activation occurring at pCa 6.02.

After the contractile measurements, the fiber was removed from the transducer and motor, solubilized in 10 μl of 1% SDS sample buffer, and stored at -80°C . The myosin heavy chain (MHC) content of each individual fiber was determined by 5% PAGE. A portion of each fiber was also run on a 12% gel to determine myosin light chain and thin-filament regulatory protein composition. A description of these procedures and representative 5 and 12% gels have been presented in a previous paper (30).

Fibers expressing solely type I MHC comprised $\geq 87\%$ of the fibers randomly isolated for study from either the pre- or the post-bed-rest soleus samples (30). Consequently, these were the only fibers included in the present analysis. Pre- and post-bed-rest means were analyzed by using a two-tailed t -test. Statistical significance was accepted at $P < 0.05$. All data are presented as means $\pm$ SE.

RESULTS

Isotonic contractile properties were studied on 201 pre- and 134 post-bed-rest fibers, all of which expressed type I MHC on 5% SDS-PAGE gels. On average, the post-bed-rest type I soleus fibers were smaller in diameter (pre-bed rest: $95 \pm 1 \mu\text{m}$, post-bed rest: $90 \pm 1 \mu\text{m}$; $P < 0.05$), produced less peak force (pre-bed rest: $1.01 \pm 0.02 \text{ mN}$, post-bed rest: $0.89 \pm 0.02 \text{ mN}$; $P < 0.05$), and had a greater V_{max} (pre-bed rest: $0.52 \pm 0.02 \text{ FL/s}$, post-bed rest: $0.71 \pm 0.05 \text{ FL/s}$; $P < 0.05$) than did the pre-bed-rest fiber population. For the eight subjects in this study, peak power was similar before and after bed rest (pre-bed rest: $13.8 \pm 0.4 \mu\text{N} \cdot \text{FL} \cdot \text{s}^{-1}$, post-bed rest: $13.6 \pm 0.6 \mu\text{N} \cdot \text{FL} \cdot \text{s}^{-1}$; $P > 0.05$).

Figure 3A demonstrates that, over the course of this study, fibers obtained from seven of the eight subjects displayed reductions in peak power. These reductions averaged 7–27% on an individual-subject basis. In contrast to the majority of subjects, the average peak power of fibers obtained from *subject 1* increased from 12.8 ± 1.1 (31 fibers) to $22.5 \pm 2.1 \mu\text{N} \cdot \text{FL} \cdot \text{s}^{-1}$ (16 fibers). This 76% relative change in peak power was over 2 SDs greater than the average change for all eight subjects combined.

Fig. 3. Mean peak power and Ca^{2+} sensitivity for individual subjects. **A:** peak power ($\mu\text{N} \cdot \text{FL} \cdot \text{s}^{-1}$). **B:** half-maximal activation ($\mu\text{mol Ca}^{2+}$). In **A** and **B**, mean pre-bed-rest (x -axis) value for each subject has been plotted vs. mean post-bed-rest (y -axis) value. Error bars, ± 1 SE. Nos. indicate subject identification nos. Solid line, line of identity. Note that *subject 1* was the only subject who displayed an increase in average fiber peak power and an increase in Ca^{2+} sensitivity after bed rest.

Peak power rose for fibers obtained from *subject 1* because their average P_o increased by 11% and their $V_{\max}$ more than doubled. These responses differed substantially from those observed for fibers from the other seven subjects. Representative pre- and post-bed-rest fibers obtained from one of these individuals (*subject 7*) are illustrated in Fig. 4. Comparison of the two fibers reveals that $V_{\max}$ was 50% greater (0.42 vs. 0.28 FL/s), whereas P_o was 35% lower (0.81 vs. 1.25 mN), for the post-bed-rest fiber. For this particular fiber, the parameter a/P_o was virtually identical pre- vs. post-bed rest so that the relative force associated with peak power output was 20 and 19% of P_o , respectively. At 20% of P_o , the force and shortening velocity of the pre-bed-rest fiber were 0.250 mN and 0.057 FL/s, respectively, for a power output of $14.2 \mu\text{N} \cdot \text{FL} \cdot \text{s}^{-1}$. At 19% of P_o , the force and shortening velocity of the post-bed-rest fiber were 0.155 mN and 0.081 FL/s, respectively, for a power output of $12.5 \mu\text{N} \cdot \text{FL} \cdot \text{s}^{-1}$, or 12% less than the power output observed for the pre-bed-rest fiber.

Fig. 4. Force-velocity-power relationships of a representative pre- and a post-bed-rest fiber. Both fibers were obtained from *subject 7*. Gel electrophoresis indicated that both fibers expressed type I myosin heavy chain exclusively. Solid lines, pre-bed rest; dotted lines, post-bed rest. See RESULTS for additional details

Because of the disproportionate effect that fibers from *subject 1* had on the interpretation of our data, we performed an analysis in which fibers from this individual were eliminated. For the 170 pre- and 118 post-bed-rest type I fibers obtained from *subjects 2–8*, P_o declined by 15% and $V_{\max}$ rose by 25% after bed rest (Table 1). The a/P_o ratio showed a small decline from 0.051 ± 0.002 to 0.044 ± 0.003 such that the force at peak power (% P_o) significantly dropped from a pre-bed-rest value of 17.3 ± 0.3 to $16.3 \pm 0.3\%$ post-bed rest ($P < 0.05$). Because of the decline in P_o and the reduced a/P_o ratio, the submaximal force associated with peak power fell by

21% ($P < 0.05$; Table 1). In contrast, the shortening velocity at peak power rose ($P < 0.05$) by 13% (Table 1). Consequently, average peak power declined by 0.79×1.13 , or by 11%, with bed rest (Table 1). A composite force-power relationship of the pre- and post-bed-rest fibers is illustrated in Fig. 5. After bed rest, peak power was substantially reduced at virtually all absolute and relative external loads. Figure 5 shows both the observed and predicted post-bed-rest force-power relationship. The latter shows the relationship that would have occurred if fiber velocity had not increased.

Table 1. Effect of 17 days of bed rest on slow type I fiber force, shortening velocity, and power

Variable	Pre-Bed Rest	Post-Bed Rest
<i>n</i>	170	118
Peak force, mN	1.04 ± 0.03	$0.88 \pm 0.02^*$
Maximal shortening velocity, FL/s	0.50 ± 0.02	$0.62 \pm 0.04^*$
Peak power, $\mu\text{N} \cdot \text{FL} \cdot \text{s}^{-1}$	14.0 ± 0.5	$12.4 \pm 0.5^*$
Force at peak power, mN	0.182 ± 0.006	$0.144 \pm 0.004^*$
Shortening velocity at peak power, FL/s	0.080 ± 0.002	$0.090 \pm 0.004^*$

Values are means $\pm$ SE for 7 subjects (*subjects 2–8*); *n*, no. of fibers. FL, fiber lengths. Significant difference between pre- and post-bed-rest means, $P < 0.05$.

Fig. 5. Composite force-power relationships for type I soleus fibers. Curves were constructed from mean force-velocity parameters of 7 subjects. *Inset*, same data plotted on a relative-force axis. Predicted curves were constructed assuming no change in fiber velocity, and a comparison of observed and predicted post-bed-rest (BR) relationships demonstrates the extent of protection induced by the increased fiber velocity.

Fiber Ca^{2+} sensitivity was determined from Hill-plot analysis (120 pre- and 88 post-bed-rest fibers). Again, the mean response of fibers obtained from *subject 1* (19 pre- and 11 post-bed-rest fibers) was opposite to the responses of fibers from the other seven subjects. In Fig. 3B, the average Ca^{2+} concentrations eliciting one-half maximal activation before and after bed rest have been plotted for each subject. *Subjects 2–8* displayed an average decline in Ca^{2+} sensitivity because substantially more free Ca^{2+} was required to attain half-maximal activation after bed rest. In contrast, fibers from *subject 1* became more sensitive to Ca^{2+} over the course of this study. In a similar manner, the Ca^{2+} activation threshold shifted to lower and higher levels of free Ca^{2+} for *subject 1* and *subjects 2–8*, respectively (Table 2).

Table 2. Effect of 17 days of bed rest on force-pCa relationships

Variable	Pre-Bed Rest	Post-Bed Rest
n	101	77
Activation threshold, $\mu\text{mol Ca}^{2+}$	0.078 ± 0.003	$0.117 \pm 0.005^*$
Half-maximal activation, $\mu\text{mol Ca}^{2+}$	1.353 ± 0.053	$1.602 \pm 0.064^*$
Slope of Hill plot for $P_r < 0.5$ (n_2)	2.05 ± 0.04	$2.27 \pm 0.05^*$
Slope of Hill plot for $P_r > 0.5$ (n_1)	1.40 ± 0.03	$1.62 \pm 0.04^*$

Values are means $\pm$ SE for 7 subjects (*subjects 2–8*); pCa, $-\log$ of Ca^{2+} concentration; P_r , ratio of submaximal force to maximal force as defined in Fig. 2; n_1 and n_2 , slope of Hill plot for values greater than and less than half-maximal activation, respectively.

*Significant difference between pre- and post-bed-rest means, $P < 0.05$.

When we eliminated *subject 1* from our analysis, the average force-pCa relationship of the remaining seven subjects was shifted to the right (Fig. 6). Thus the Ca^{2+} required for activation threshold increased significantly with bed rest as did the Ca^{2+} concentration eliciting one-half maximal activation (Table 2). In addition, the slope of the force-pCa curve below (n_2) and above (n_1) half-maximal activation increased with bed rest (Table 2).

Fig. 6. Mean pCa-force relationships for type I soleus fibers. Values are means $\pm$ SE of relative force (P_r) for 7 subjects. ●, Pre-bed rest; ○, post-bed rest.

DISCUSSION

The primary goal of this study was to investigate changes in the isotonic contractile properties of single muscle fibers after a 17-day period of non-weight bearing. We were particularly interested in peak power because this variable is an important characteristic of animal locomotor systems (26), and declines in power would likely have important physiological and functional implications both in space and on return to Earth. In this study, we focused on fibers expressing type I MHC because 1) in animal models, type I fibers are preferentially affected by non-weight bearing (11) and 2) type I fibers make up the majority of fibers present in the human soleus before and after 17 days of bed rest (28).

The average peak power of the 201 type I soleus fibers obtained before bed rest was $13.8 \pm 0.4 \mu\text{N} \cdot \text{FL} \cdot \text{s}^{-1}$. In previous studies from this laboratory, we reported mean peak power values of 8.2 ± 0.3 and $9.8 \pm 0.3 \mu\text{N} \cdot \text{FL} \cdot$

s⁻¹ for human type I gastrocnemius and type I soleus fibers, respectively (31, 32). The greater power values in the present study cannot be attributed to variations in shortening velocity as there was little difference in fiber $V_{\max}$ across these three studies. However, the fibers in the present study were larger in diameter and consequently produced more force than did fibers from our previous work. This appears to be the major reason for the greater peak power values in the present study. Additionally, the force-velocity relationships presented here were less curved (higher a/P_o ratio), such that higher velocities were obtained at a given percent of P_o . Bottinelli et al. (2) studied the isotonic contractile properties of human muscle fibers, and their reported peak power value for the slow type I fiber was considerably lower than ours. The primary reason for the discrepancy was fiber force because the mean value (corrected for the different temperature employed) was <70% of the fiber force reported here.

The average peak power for fibers obtained from seven of the eight subjects in this study declined by 11% with bed rest. A decline in peak power with non-weight bearing is consistent with observations made on type I soleus fibers obtained from adult rats subjected to hindlimb suspension (1, 20). However, reductions in peak power after 14–21 days of rat hindlimb suspension exceed the decline observed in the present study by a factor of three to four. Taken together, these data indicate that type I fibers from the human soleus experience less atrophy and loss of peak power than do fibers obtained from rats exposed to hindlimb suspension. This is in agreement with proposed interspecies differences in the net rate of protein loss during non-weight bearing (8).

The decline in peak power of the human fibers can be attributed primarily to a reduction in the ability of the post-bed-rest fibers to produce force, with a secondary contribution from changes in the curvature of the force-velocity relation. The loss of force after 17 days of bed-rest results primarily from a reduction in fiber CSA coupled with a relatively small reduction in force per CSA (30). The force-velocity relationship had a slightly greater curvature after bed rest so that peak power was attained at a lower percentage of fiber P_o . Both of these changes meant that the post-bed-rest fibers produced less force when they shortened at their optimal velocity for power production. Figure 5 shows a pre-, a post-, and a predicted post-bed-rest force-power relationship. The predicted curve shows the force-power relationship that would have occurred if fiber velocity had remained unchanged and peak power had declined in direct proportion to the average 21% loss in fiber force (at peak power). It is clear from Fig. 5 that the observed reduction in peak power was only about one-half of the decline that could be explained solely by a change in force production. The reason is that when the post-bed-rest fibers were producing optimal force for power output, they were shortening at a 13% greater velocity than were the pre-bed-rest fibers. The observed peak power was therefore greater than one would predict based solely on a change in fiber force. We conclude that the elevated shortening velocity of the post-bed-rest fibers played a compensatory role by reducing the reduction in peak power that occurred during non-weight bearing.

In the present study, we included only those fibers that expressed type I MHC. In previous work, we concluded that variations in myosin light chain composition could not explain the elevated shortening velocities of post-bed-rest fibers expressing type I MHC (30). The mechanism underlying the increase in their shortening velocity is therefore not clear. One possibility is that an unidentified MHC isoform comigrates with the adult β -MHC isoform observed on our gels (7). A second possibility pertains to morphological alterations that occur during non-weight bearing. It has recently been reported that post-bed-rest fibers obtained from the subjects in the present study had a reduction in thin filament density after the 17-day period of bed rest (25a). Shortening velocity is known to increase when the myofilament lattice spacing is expanded experimentally, whereas force changes very little under these conditions (23). Thus one would predict that power would increase as the myofilament lattice spacing increased.

It is important to note that the magnitude of the change in shortening velocity was insufficient to completely offset the lower forces of the post-bed-rest fibers. Thus countermeasures to human non-weight bearing must address the issue of muscle fiber atrophy. Also, it will be important to determine whether the increase in fiber

shortening velocity after bed rest affects other functional properties of human muscle fibers. McDonald and Fitts (21) found that myosin ATPase activity was increased in soleus fibers from hindlimb-suspended rats. If similar changes occur in human muscle after bed rest, one might predict a decrease in the efficiency of contraction and an increase in muscle fatigability.

The plantar flexors and knee extensors produce significantly less torque at any given angular velocity after ≥ 30 days of non-weight bearing (5, 15). In this study, the voluntary plantar flexor torque-velocity relationship did not change over the 17-day period of bed rest (25, 28). In a study of the same subjects, Narici et al. (25) studied the effects of direct electrical stimulation (50 Hz) of the plantar flexors. They found that peak torque declined by 10 and 18% from the pre-bed-rest baseline on the 13th day of bed rest and on the 2nd day of recovery from bed rest, respectively. These data compare favorably with our observation of a 12 (8 subjects) to 15% (7 subjects) reduction in peak Ca^{2+} -activated force of single muscle fibers obtained immediately at the end of the bed-rest period. Thus, when neural mechanisms (motivation, coactivation of antagonist) modulating muscle performance were bypassed, the relative change in the strength of the triceps surae was in good agreement with the relative change in the maximal Ca^{2+} -activated force of single soleus fibers.

Post-bed-rest fibers also experienced a significant decrease in Ca^{2+} sensitivity, as indicated by a 50% increase in the free- Ca^{2+} concentration at the activation threshold and an 18% increase in the Ca^{2+} required to attain half-maximal activation. These results are qualitatively similar to those observed for rat type I soleus fibers after hindlimb suspension (1, 11, 22). However, changes in the Ca^{2+} sensitivity of rat fibers are of a greater magnitude than that observed for the human fibers in the present study. Functionally, a decrease in Ca^{2+} sensitivity could have implications during the development of fatigue when the intracellular free- Ca^{2+} concentration may not completely saturate the Ca^{2+} -binding sites of the thin filament (9). Westerblad and Allen (29) have concluded that the intracellular Ca^{2+} concentration of mammalian muscle fibers may drop to as low as pCa 6.0 during fatiguing stimulation. At this intracellular Ca^{2+} concentration, force of the pre- and post-bed-rest fibers was 41 and 33%, respectively, of the maximal force (i.e., at pCa 4.5). The greater decline in the post-bed-rest fibers can be attributed to the right shift (reduced Ca^{2+} sensitivity) in the force-pCa relationship (Fig. 6).

We also found that the slopes of the force-pCa relationship (n_1 and n_2) increased after bed rest (Table 2). An increased n_1 and n_2 is generally interpreted as an indication of a greater degree of molecular cooperativity during activation of the cross bridge. The degree of cooperativity (indicated by the value of n_1 and n_2) can be influenced by alterations in thin-filament regulation such as an increased affinity of troponin C for Ca^{2+} and/or by thick-filament regulation (24).

A number of mechanisms are believed to confer Ca^{2+} sensitivity to skeletal muscle fibers [for recent review, see Moss et al. (24)]. In fast- and intermediate-velocity mammalian fibers, expression of the fast isoforms of troponin I, troponin T, and/or tropomyosin is associated with a decrease in Ca^{2+} sensitivity and an increase in cooperative activation of the thin filament (14, 27). In this regard, de novo expression of the fast isoforms of troponin T and troponin I have been reported in rat soleus muscles after hindlimb suspension (4). However, it is not known what effect, if any, these isoform shifts have on the force-pCa characteristics of slow fibers. Second, analysis of the present fibers on 12% polyacrylamide gels revealed the presence of only the slow isoform of troponin T in pre- and post-bed-rest fibers (data not shown). We were unable to reliably quantify troponin I isoforms because, for human fibers, this protein tends to comigrate with myosin light chain₂ on our gel system. An alternate hypothesis is that the rightward shift in the force-pCa relationship was a result of the same mechanism proposed for the increase in shortening velocity, i.e., a change in filament lattice geometry. A rightward shift in the force-pCa relationship occurs when the filament lattice expands by $\sim 20\%$ (13).

It is difficult to explain why fibers obtained from *subject 1* showed increases in peak power and Ca^{2+} sensitivity with bed rest. Contractile properties of pre-bed-rest fibers obtained from this individual were within normal

limits (for example, see Fig. 3). What is surprising is that the post-bed-rest fibers obtained from this individual were, on average, larger in diameter than his pre-bed-rest fibers. As a result, the post-bed-rest fibers produced more force and power. This cannot be attributed to a problem with compliance because subjects were monitored 24 h/day by the staff at the NASA Ames Research Center. It also seems unlikely that these post-bed-rest results were due to methodological factors. Post-bed-rest fibers from this subject were studied at the same time, and with the same equipment and solutions, as were the post-bed-rest fibers from the other seven subjects. It is important to note that *subject 1* displayed a reduction in electrically elicited plantar flexor torque that was similar to the responses of the other seven subjects (M. V. Narici, personal communication). Thus, at the whole-muscle level, *subject 1* appeared no different from the other subjects. The most plausible explanation for the disparate results of this subject at the single-fiber level is that his post-bed-rest biopsy sample was simply not representative of the soleus muscle as a whole. Recently, Harridge et al. (16) proposed a similar explanation for discrepancies between contractile properties measured at the whole-muscle and single-fiber levels in human subjects.

In summary, the peak power output of type I soleus fibers obtained from seven of eight subjects declined by an average of 11% after 17 days of bed rest. This decline in peak power would have doubled if not for a bed-rest-induced rise in fiber shortening velocity. Fibers from these same subjects showed a small, but statistically significant, reduction in Ca^{2+} sensitivity and an increase in cooperative activation of the cross bridge. These results provide a cellular basis for a number of the functional changes observed at the whole muscle level after bed rest, particularly in experimental designs where neural mechanisms regulating muscle activation are bypassed.

The authors especially thank all of the subjects for their commitment throughout the study and the staff of the Human Research Facility at the National Aeronautics and Space Administration (NASA) Ames Research Center, especially Dee O'Hara and Dr. Sara Arnaud.

FOOTNOTES

This research was supported by NASA Grant NAS9-18768 (to R. H. Fitts).

AUTHOR NOTES

Address for reprint requests: R. H. Fitts, Marquette Univ., Dept. of Biology, Wehr Life Sciences Bldg., P.O. Box 1881, Milwaukee, WI 53201-1881.

REFERENCES

- 1 Bangart J. J., Widrick J. J., Fitts R. H. Effect of intermittent weight bearing on soleus fiber force-velocity-power and force-pCa relationships. *J. Appl. Physiol.* 82:1997-1905, 1990
- 2 Bottinelli R., Canepari M., Pellegrino M. A., Reggiani C. Force-velocity properties of human skeletal muscle fibres: myosin heavy chain isoform and temperature dependence. *J. Physiol. (Lond.)* 495:199-657, 1996
- 3 Caiozzo V. J., Haddad F., Baker M. J., Herrick R. E., Prietto N., Baldwin K. M. Microgravity-induced transformations of myosin isoforms and contractile properties of skeletal muscle. *J. Appl. Physiol.* 81:1996-123, 1996
- 4 Campione M., Ausoni S., Guezennec C. Y., Schiaffino S. Myosin and troponin changes in rat soleus muscle after hindlimb suspension. *J. Appl. Physiol.* 74:1993-115, 1993
- 5 Dudley G. A., Duvoisin M. R., Convertino V. A., Buchanan P. Alterations of the in vivo torque-velocity relationship of human skeletal muscle following 30 days exposure to simulated microgravity. *Aviat. Space Environ. Med.* 60:1989-659, 1989

- 6 Fabiato A., Fabiato F. Calculator programs for computing the composition of the solutions containing multiple metals and ligands used for experiments in skinned muscle cells. *J. Physiol. Paris* 75:1979463505
- 7 Fauteck S. P., Kandarian S. C. Sensitive detection of myosin heavy chain composition in skeletal muscle under different loading conditions. *Am. J. Physiol.* 268:Cell Physiol. 371995C419C424
- 8 Ferrando A. A., Lane H. W., Stuart C. A., Davis-Street J., Wolfe R. R. Prolonged bed rest decreases skeletal muscle and whole body protein synthesis. *Am. J. Physiol.* 270:Endocrinol. Metab. 331996E627E633
- 9 Fitts R. H. Cellular mechanisms of muscle fatigue. *Physiol. Rev.* 74:19944994
- 10 Fitts R. H., Metzger J. M., Riley D. A., Unsworth B. R. Models of disuse: a comparison of hindlimb suspension and immobilization. *J. Appl. Physiol.* 60:198619471953
- 11 Gardetto P. R., Schluter J. M., Fitts R. H. Contractile function of single muscle fibers after hindlimb suspension. *J. Appl. Physiol.* 66:198927392749
- 12 Godt R. E., Lindley B. D. Influence of temperature upon contractile activation and isometric force production in mechanically skinned muscle fibers of the frog. *J. Gen. Physiol.* 80:1982279297
- 13 Godt R. E., Maughan D. W. Influence of osmotic compression on calcium activation and tension in skinned muscle fibers of the rabbit. *Pflügers Arch.* 391:1981334337
- 14 Greaser M. L., Moss R. L., Reiser P. J. Variations in contractile properties of rabbit single muscle fibres in relation to troponin T isoforms and myosin light chains. *J. Physiol. (Lond.)* 406:19888598
- 15 Grigoryeva L. S., Kozlovskaya I. B. Effect of weightlessness and hypokinesia on velocity and strength properties of human muscles. *Kosm. Biol. Aviakosm. Med.* 21:19872730
- 16 Harridge S. D. R., Bottinelli R., Canepari M., Pellegrino M. A., Reggiani C., Esbjörnsson M., Saltin B. Whole-muscle and single-fibre contractile properties and myosin heavy chain isoforms in humans. *Pflügers Arch.* 432:1996913920
- 17 Hill A. V. The heat of shortening and the dynamic constants of muscle. *Proc. R. Soc. Lond. B Biol. Sci.* 126:1938136195
- 18 Julian F. J., Moss R. L. Effects of calcium and ionic strength on shortening velocity and tension development in frog skinned muscle fibres. *J. Physiol. (Lond.)* 311:1981179199
- 19 Larsson L., Li X., Berg W. E., Frontera W. R. Effects of removal of weight-bearing function on contractility and myosin isoform composition in single human skeletal muscle cells. *Pflügers Arch.* 432:1996320328
- 20 McDonald K. S., Blaser C. A., Fitts R. H. Force-velocity and power characteristics of rat soleus muscle fibers after hindlimb suspension. *J. Appl. Physiol.* 77:199416091616
- 21 McDonald K. S., Fitts R. H. Effect of hindlimb unweighting on single soleus fiber maximal shortening velocity and ATPase activity. *J. Appl. Physiol.* 74:199329492957
- 22 McDonald K. S., Fitts R. H. Effect of hindlimb unloading on rat soleus fiber force, stiffness, and calcium sensitivity. *J. Appl. Physiol.* 79:199517961802
- 23 Metzger J. M., Moss R. L. Shortening velocity in skinned single muscle fibers. Influence of filament lattice spacing. *Biophys. J.* 52:1987127131
- 24 Moss R. L., Diffie G. M., Greaser M. L. Contractile properties of skeletal muscle fibers in relation to myofibrillar protein isoforms. *Rev. Physiol. Biochem. Pharmacol.* 126:1995163
- 25 Narici M. V., Kayser B., Barattini P., Cerretelli P. Changes in electrically evoked skeletal muscle contractions during 17-day spaceflight and bed rest. *Int. J. Sports Med.* 18:1997S290S292
- 25a Riley D. A., Bain J. L., Thompson J. L., Fitts R. H., Widrick J. J., Trappe S. W., Trappe T. A., Costill D. L. Disproportionate loss of thin filaments in human soleus muscle after 17-day bedrest. *Muscle Nerve* 21:199812801289
- 26 Rome L. C., Funke R. P., Alexander R. M., Lutz G., Aldridge H., Scott F., Freadman M. Why animals have different muscle fibre types. *Nature* 335:1988824827
- 27 Schachat F. H., Diamond M. S., Brandt P. W. Effect of different troponin T-tropomyosin combinations on thin filament activation. *J. Mol. Biol.* 198:1987551554

- 28 Trappe S. W., Trappe T. A., Costill D. L., Fitts R. H. Human calf muscle function in response to 17-days of bed rest (Abstract). **Med. Sci. Sports Exerc.** 28:1996S146
- 29 Westerblad H., Allen D. G. Changes of myoplasmic calcium concentration during fatigue in single mouse muscle fibers. **J. Gen. Physiol.** 98:1991615635
- 30 Widrick J. J., Romatowski J. G., Bain J. L. W., Trappe S. W., Trappe T. A., Thompson J. L., Costill D. L., Riley D. A., Fitts R. H. Effect of 17 days of bed rest on peak isometric force and unloaded shortening velocity of human soleus fibers. **Am. J. Physiol.** 273:Cell Physiol. 42:1997C1690C1699
- 31 Widrick J. J., Romatowski J. G., Karhanek M., Fitts R. H. Contractile properties of rat, rhesus monkey, and human type I fibers. **Am. J. Physiol.** 272:Regulatory Integrative Comp. Physiol. 41:1997R34R42
- 32 Widrick J. J., Trappe S. W., Costill D. L., Fitts R. H. Force-velocity and force-power properties of single muscle fibers from elite master runners and sedentary men. **Am. J. Physiol.** 271:Cell Physiol. 40:1996C676C683