

Marquette University

e-Publications@Marquette

Chemistry Faculty Research and Publications

Chemistry, Department of

4-2012

Synthesis and Structural Characterization of Iron(II) Complexes with Tris(imidazolyl)phosphane Ligands: A Platform for Modeling the 3-Histidine Facial Triad of Nonheme Iron Dioxygenases

Michael M. Bittner

Marquette University, michael.bittner@marquette.edu

Jacob Baus

Marquette University, jacob.baus@marquette.edu

Sergey V. Lindeman

Marquette University, sergey.lindeman@marquette.edu

Adam T. Fiedler

Marquette University, adam.fiedler@marquette.edu

Follow this and additional works at: https://epublications.marquette.edu/chem_fac

 Part of the [Chemistry Commons](#)

Recommended Citation

Bittner, Michael M.; Baus, Jacob; Lindeman, Sergey V.; and Fiedler, Adam T., "Synthesis and Structural Characterization of Iron(II) Complexes with Tris(imidazolyl)phosphane Ligands: A Platform for Modeling the 3-Histidine Facial Triad of Nonheme Iron Dioxygenases" (2012). *Chemistry Faculty Research and Publications*. 574.

https://epublications.marquette.edu/chem_fac/574

Marquette University

e-Publications@Marquette

Chemistry Faculty Research and Publications/College of Arts and Sciences

This paper is NOT THE PUBLISHED VERSION.

Access the published version at the link in the citation below.

European Journal of Inorganic Chemistry, Vol. 2012, No. 11 (April 2012): 1848-1856. [DOI](#). This article is © Wiley and permission has been granted for this version to appear in [e-Publications@Marquette](#). Wiley does not grant permission for this article to be further copied/distributed or hosted elsewhere without the express permission from Wiley.

Synthesis and Structural Characterization of Iron(II) Complexes with Tris(imidazolyl)phosphane Ligands: A Platform for Modeling the 3-Histidine Facial Triad of Nonheme Iron Dioxygenases

Michael M. Bittner

Department of Chemistry, Marquette University, Milwaukee, WI

Jacob S. Baus

Department of Chemistry, Marquette University, Milwaukee, WI

Sergey V. Lindeman

Department of Chemistry, Marquette University, Milwaukee, WI

Adam T. Fiedler


Department of Chemistry, Marquette University, Milwaukee, WI

Abstract

Several monoiron(II) complexes containing tris(imidazolyl)phosphane (TIP) ligands have been prepared and structurally characterized by using X-ray crystallography and NMR spectroscopy. Two TIP ligands were employed: tris(2-phenylimidazol-4-yl)phosphane (4-TIP^{Ph}) and tris(4,5-diphenyl-1-methylimidazol-2-yl)phosphane (2-TIP^{Ph²}). These tridentate ligands resemble the 3-histidine (3His) facial triad found recently in the active sites of certain nonheme iron dioxygenases. Three of the reported complexes are designed to serve as convenient precursors to species that model the enzyme–substrate intermediates of 3His dioxygenases; thus, each contains an [Fe(κ^3 -TIP)]²⁺ unit in which the remaining coordination sites are occupied by easily displaced ligands, such as solvent molecules and/or carboxylate groups. The viability of these complexes as precursors was demonstrated through the synthesis of TIP-based complexes with β -diketonate and salicylate ligands that represent faithful models of β -diketone dioxygenase and salicylate 1,2-dioxygenase, respectively.

Abstract

A series of monoiron(II) complexes with tris(imidazolyl)phosphane ligands have been generated and characterized by using X-ray crystallography. These complexes model the resting and substrate-bound states of certain nonheme iron enzymes, such as β -diketone dioxygenase and salicylate 1,2-dioxygenase, which employ three facially coordinating histidine ligands in their active sites.




Introduction


Mononuclear nonheme iron dioxygenases play a central role in the oxidative catabolism of a wide range of biomolecules and pollutants.¹ Members of this enzyme family include the extradiol catechol dioxygenases,² Rieske dioxygenases,³ homogentisate dioxygenase,⁴ and (chloro)hydroquinone dioxygenases.⁵ These enzymes feature a common active-site motif in which the ferrous center is facially ligated by one aspartate (or glutamate) and two histidine residues [the so-called 2-His-1-carboxylate (2H1C) facial triad].⁶ However, recent structural studies have shown that the Asp/Glu ligand in some monoiron dioxygenases is replaced with His, resulting in the 3His facial triad.⁷ Members of this “3His family” catalyze novel transformations that have expanded the known boundaries of Fe dioxygenase chemistry. For example, cysteine dioxygenase (CDO)⁸ – the first 3His enzyme to be structurally characterized – catalyzes the initial step in L-cysteine catabolism by converting the thiol into a sulfinic acid (Scheme 1), while β -diketone dioxygenase (Dke1) oxidizes acetylacetone to acetic acid and 2-oxopropanal.⁹ Other 3His Fe dioxygenases include gentisate 1,2-dioxygenase (GDO)¹⁰ and salicylate 1,2-dioxygenase (SDO),¹¹ both of which oxidatively cleave aromatic C–C bonds (Scheme 1). Each of these microbial enzymes participates in the degradation pathways of polycyclic aromatic hydrocarbons. While the reaction catalyzed by GDO is very similar to those catalyzed by the extradiol

catechol dioxygenases and likely follows a similar mechanism, SDO is unique in performing the oxidative cleavage of an aromatic ring with only one electron-donating group.


Cysteine Dioxygenase (CDO):




β -Diketone Dioxygenase (Dke1):



Gentisate 1,2-Dioxygenase (GDO):




Salicylate 1,2-Dioxygenase (SDO):



Scheme 1

Our knowledge of nonheme Fe dioxygenases has greatly benefitted from the development of small-molecule analogues that replicate important structural, spectroscopic, and/or functional properties of the enzyme active sites.¹² The 2H1C triad has been suitably modeled with anionic, tridentate supporting ligands such as tris(pyrazol-1-yl)borates (Tp),¹³ bis(pyrazolyl)acetates,¹⁴ and bis(1-alkylimidazol-2-yl)propionates.¹⁵ The last two ligand sets replicate the mixed N₂O donor set of the 2H1C triad by the inclusion of carboxylate arms. Given the unique and significant reactions catalyzed by the 3His family of Fe dioxygenases, it is important to develop supporting ligands with specific relevance to the 3His facial triad. To this end, we have sought to exploit the tris(imidazol-2-yl)phosphane (2-TIP^{R2}) and tris(imidazol-4-yl)phosphane (4-TIP^R) frameworks shown in Scheme 2, which accurately mimic the charge and donor strength of the 3His coordination environment. These ligands were initially generated to model the 3His ligand sets found in the active sites of carbonic anhydrase (Zn²⁺) and cytochrome c oxidase (Cu²⁺).¹⁶ To date, the application of the TIP framework to Fe systems has been limited to homoleptic [Fe(TIP)₂]^{2+/3+} complexes¹⁷ and carboxylate-bridged diiron(III) species.^{17c,18}




Scheme 2

A key advantage of the 2-TIP^{R2} and 4-TIP^R ligands is that their steric properties can be easily modified by altering the R substituent(s). Thus far, we have primarily employed the 2-TIP^{Ph2} and 4-TIP^{Ph} ligands, as the steric bulk of the phenyl rings discourages both dimerization and formation of the homoleptic [Fe(TIP)₂]²⁺ complexes. Recently, we described the synthesis and structural characterization of a series of [Fe²⁺(4-TIP^{Ph})(acac^X)]OTf complexes (acac^X = substituted β-diketonate; OTf = triflate) that serve as models of the Dke1 enzyme–substrate complex.¹⁹ These models were prepared by directly mixing one equivalent of the sodium salt of the appropriate β-diketone, Na(acac^X), with equimolar amounts of Fe(OTf)₂ and 4-TIP^{Ph} in MeOH. This “one-pot” approach, however, is not successful for various combinations of supporting and “substrate” ligands. Thus, as described in this article, we have generated several Fe²⁺ complexes with κ³-TIP ligands that also contain displaceable ligands (such as solvent, triflate, benzoate, and acetate) bound to the opposite face of the octahedron. These complexes resemble the resting states of 3His Fe dioxygenases, which feature two or three *cis*-labile H₂O molecules.²⁰ In addition, it is shown that these TIP-based complexes serve as excellent precursors for the formation of monoiron complexes with three facial imidazole donors and various bound substrates, including β-diketonates and salicylates (mimics of Dke1 and SDO, respectively). Thus, the chemistry described here establishes a valuable platform for future synthetic modeling studies of nonheme Fe dioxygenases with the 3His facial triad.

Results and Discussion

Fe²⁺ Complexes Containing 2-TIP^{Ph2}

The novel 2-TIP^{Ph2} ligand was synthesized by means of lithiation of 4,5-diphenyl-1-methylimidazole at the 2-position at –78 °C, followed by addition of PCl₃ (0.33 equiv.). Reaction of 2-TIP^{Ph2} with Fe(OTf)₂ in MeCN provided the complex [1](OTf)₂ in 60 % yield (Scheme 3). Crystals suitable for X-ray diffraction (XRD) analysis were obtained by layering a concentrated MeCN solution with diethyl ether. The structure features two symmetrically independent [1]²⁺ units with nearly identical metric parameters (Table 1; details concerning the data collection and analysis of all X-ray structures are summarized in Table 4). As shown in Figure 1, the six-coordinate (6C) Fe²⁺ center is ligated by 2-TIP^{Ph2} and three MeCN ligands in a distorted octahedral geometry. As expected, the 2-TIP^{Ph2} ligand coordinates in a facial manner. The average Fe–N distance of 2.19 Å is indicative of a high-spin Fe²⁺ center (*S* = 2), consistent with the measured magnetic moment of 5.2 μ_B. The triflate counteranions are not bound to the metal centers, and the asymmetric unit also contains one equivalent of noncoordinated Et₂O.



Scheme 3

Table 1. Selected metric parameters for $[\text{Fe}^{2+}(\text{L}_{\text{N}3})(\text{MeCN})_3]^{2+}$ complexes. Bond lengths in Å and angles in degrees.

	$[\mathbf{1}](\text{OTf})_2 \cdot 0.5\text{Et}_2\text{O}$ ^[a]	$[\text{Fe}(\text{trisoxt}^{\text{tBu}})(\text{MeCN})_3]^{2+}$ (ref.22) ^[b]	$[\text{Fe}(\text{tpm}^{\text{Ph}2})(\text{MeCN})_3]^{2+}$ (ref.21) ^[c]
Fe1–N2	2.186(1)	2.257(2)	2.199(2)
Fe1–N4	2.177(1)	2.205(2)	2.196(2)
Fe1–N6	2.182(1)	2.215(2)	2.205(3)
Fe1–N7	2.196(1)	2.163(2)	2.131(3)
Fe1–N8	2.179(1)	2.131(2)	2.166(2)
Fe1–N9	2.205(1)	2.171(3)	2.156(3)
Fe–N _{TIP} (av.)	2.181	2.226	2.200
Fe–N _{solv} (av.)	2.193	2.155	2.151
N2–Fe1–N4	88.87(5)	84.62(6)	84.40(9)
N2–Fe1–N6	91.43(5)	82.12(6)	85.84(9)
N4–Fe1–N6	88.32(5)	86.89(7)	83.27(9)
N7–Fe1–N8	85.48(5)	90.25(8)	87.6(1)
N7–Fe1–N9	82.06(5)	91.33(8)	86.4(1)
N8–Fe1–N9	83.23(5)	86.00(8)	90.3(1)

[a] Average values for the two independent, but chemically equivalent $[\mathbf{1}]^+$ cations.

[b] $\text{trisoxt}^{\text{tBu}} = 1,1,1\text{-tris}(4\text{-tert-butylloxazolin-2-yl})\text{ethane}$.

[c] $\text{tpm}^{\text{Ph}2} = \text{tris}(3,5\text{-diphenylpyrazol-1-yl})\text{methane}$.




Figure 1

Thermal ellipsoid plot (50 % probability) of $[\mathbf{1}](\text{OTf})_2 \cdot 0.5\text{Et}_2\text{O}$. Only one of the symmetrically inequivalent $[\mathbf{1}](\text{OTf})_2$ units is shown. Hydrogen atoms, counteranions, and noncoordinating solvent molecules have been omitted for clarity.

Two related high-spin Fe^{2+} structures with $[\text{Fe}(\text{L}_{\text{N}3})(\text{MeCN})_3]^{2+}$ compositions have been reported in the literature, and their metric parameters are also provided in Table 1. The average Fe–N_{TIP} distance of 2.18 Å in $[\mathbf{1}]^{2+}$ is significantly shorter than the distances observed for the analogous tris(3,5-diphenylpyrazol-1-yl)methane ($\text{tpm}^{\text{Ph}2}$)²¹ and 1,1,1-tris(4-*tert*-butylloxazolin-2-yl)ethane ($\text{trisoxt}^{\text{tBu}}$)²² complexes, which display average Fe–N distances of 2.20 and 2.23 Å, respectively. Conversely, the average Fe–N_{MeCN} distance in $[\mathbf{1}]^{2+}$ is approximately 0.04 Å longer than those reported

for the tpm^{Ph_2} and $\text{trisox}^{\text{tBu}}$ complexes. Both facts suggest that $2\text{-TIP}^{\text{Ph}_2}$ is a somewhat stronger donor than other neutral N3 ligands that have appeared in the literature.

Elemental analysis performed with ground and dried crystals of $[\mathbf{1}](\text{OTf})_2$ indicate that at least two MeCN ligands are removed under vacuum. In addition, evidence for Fe–OTf bonding in non-coordinating solvents was obtained by using ^{19}F NMR spectroscopy. For $[\mathbf{1}](\text{OTf})_2$ in CD_3CN , the triflate counteranion gives rise to a sharp peak at $\delta = -79.2$ ppm, which is identical to the chemical shift observed for $[\text{NBu}_4]\text{OTf}$ under the same conditions. The lengthy longitudinal relaxation time (T_1 value) of 128 ms measured for this feature suggests that the triflate counteranion is only weakly associated with the $[\text{Fe}(2\text{-TIP}^{\text{Ph}_2})]^{2+}$ unit in MeCN. In contrast, the ^{19}F NMR spectrum of $[\mathbf{1}](\text{OTf})_2$ in CD_2Cl_2 exhibits a broad feature at $\delta = -60.9$ ppm with a short T_1 value of 14 ms (Figure S1 in the Supporting Information), which indicates that the triflate ion is directly bound to the Fe center.

Reaction of equimolar amounts of $\text{Fe}(\text{OTf})_2$, $2\text{-TIP}^{\text{Ph}_2}$, and sodium benzoate (NaOBz) in MeOH provided the colorless complex $[\text{Fe}(2\text{-TIP}^{\text{Ph}_2})(\text{OBz})(\text{MeOH})]\text{OTf}$ ($[\mathbf{2}]\text{OTf}$), as shown in Scheme 3. X-ray-quality crystals were obtained from a solution of $[\mathbf{2}]\text{OTf}$ in MeOH layered with pentane. The resulting structure reveals a pentacoordinate (5C) iron(II) center with a $\kappa^3\text{-}2\text{-TIP}^{\text{Ph}_2}$ ligand, monodentate benzoate ligand, and bound solvent (Figure 2). In addition to the second-sphere triflate anion, the asymmetric unit also contains four MeOH molecules that do not directly interact with the $[\mathbf{2}]^+$ cation. The complex adopts a distorted square-pyramidal geometry ($\tau = 0.25$ ²³) with an O_2N_2 pseudobasal plane. Two phenyl rings of the $2\text{-TIP}^{\text{Ph}_2}$ ligand lie across the vacant coordination site (i.e., parallel to the plane of the benzoate ligand), which prevents further solvent binding. The Fe– N_{TIP} and Fe–O distances are typical for high-spin Fe^{2+} centers (Table 2). The H atom of the coordinated MeOH molecule was found objectively and refined. The resulting $\text{O}2\cdots\text{O}3$ distance of 2.610(2) Å and $\text{H}3\cdots\text{O}2$ distance of 1.81(1) Å are indicative of an intramolecular hydrogen bond that closes a six-membered ring.




Figure 2

Thermal ellipsoid plot (50 % probability) derived from [2]OTf·4MeOH. Non-coordinating solvent molecules, counteranions, and most H atoms have been omitted for clarity. The dotted line indicates the hydrogen-bonding interaction between H3 of the MeOH ligand and O2 of the benzoate anion.

Table 2. Selected metric parameters for ferrous carboxylate complexes [2]OTf·4MeOH, 3·2CH₂Cl₂, and [5]BPh₄·3MeOH. Bond lengths in Å and angles in degrees.

	[2]OTf·4MeOH	3·2CH ₂ Cl ₂ [a]	[5]BPh ₄ ·3MeOH
Fe1–N2	2.124(2)	2.129(3)	2.193(4)
Fe1–N4	2.127(2)	2.111(3)	2.195(4)
Fe1–N6	2.226(2)	2.210(3)	2.186(4)
Fe–N _{LN3} (av.)	2.158	2.150	2.191
Fe1–O1	2.011(1)	1.977(3)	2.245(4)
Fe1–O2			2.256(4)
Fe1–O3(N7)	2.105(1)	2.144(3)	2.077(4)
O1–C _{carboxyl}	1.273(2)	1.271(5)	1.267(6)
O2–C _{carboxyl}	1.254(2)	1.244(4)	1.268(6)
N2–Fe1–N4	93.94(6)	94.3(1)	90.2(2)
N2–Fe1–N6	85.36(6)	80.7(1)	90.3(2)
N4–Fe1–N6	91.69(6)	87.4(1)	90.6(2)
O1–Fe1–N2	153.44(6)	152.4(1)	105.9(2)
O1–Fe1–N4	112.06(6)	110.9(1)	91.1(2)
O1–Fe1–N6	88.38(6)	89.2(1)	163.7(2)
O1–Fe1–O3(N7)	87.64(6)	98.7(1)	84.6(2)
N2–Fe1–O3(N7)	93.34(6)	89.0(1)	93.9(2)
N4–Fe1–O3(N7)	100.01(6)	96.6(1)	174.8(2)
N6–Fe1–O3(N7)	168.29(6)	169.2(1)	92.6(2)
τ value	0.25	0.28	N/A

[a] Data from the literature.²⁴ The N and O atoms in the 3·2CH₂Cl₂ structure were renumbered to correspond to the numbering scheme used for the other complexes.

Complex [2]⁺ resembles the structure of [Fe(^{Ph,Me}Tp)(OBz)(^{Ph,Me}pyz)] (**3**; in which ^{Ph,Me}pyz = 3-phenyl-5-methylpyrazole) published by Fujisawa and co-workers.²⁴ Both complexes feature a distorted square-pyramidal geometry with a monodentate benzoate ion linked to a neutral ligand by means of an intramolecular hydrogen bond. As shown in Table 2, the metric parameters of [2]⁺ and **3** are quite similar; indeed, the average Fe–N_{TIP} distance of 2.16 Å found for [2]⁺ is only 0.01 Å longer than the average Fe–N_{Tp} distance in **3**. This result is consistent with our previous study of [Fe²⁺(L_{N3})(β-diketonato)]⁺⁰ complexes that found only slight differences (on average) between Fe–N_{TIP} and Fe–N_{Tp} bond lengths in 5C species, despite the different charges of the supporting ligands.¹⁹

Starting from either of these two Fe(2-TIP^{Ph2}) precursors – [1](OTf)₂ or [2]OTf – we were able to generate the complex [Fe(2-TIP^{Ph2})(acac^{PhF3})]OTf ([4]OTf; Scheme 3; in which acac^{PhF3} = anion of 4,4,4-trifluoro-1-phenyl-1,3-butanedione). The acac^{PhF3} ligand was selected for two reasons: (i) it is a viable

Dke1 substrate,[25](#) and (ii) previous studies in our laboratory found that $[\text{Fe}(\text{L}_{\text{N}3})(\text{acac}^{\text{PhF}3})]^{+0}$ complexes exhibit intense $\text{Fe}^{2+} \rightarrow \text{acac}^{\text{PhF}3}$ MLCT bands that serve as useful spectroscopic markers.[19](#) For both $\text{Fe}(2\text{-TIP}^{\text{Ph}2})$ precursors, reaction with $\text{Na}(\text{acac}^{\text{PhF}3})$ provides a deep purple solution that displays an absorption manifold centered at 502 nm ($\epsilon = 700 \text{ M}^{-1} \text{ cm}^{-1}$; see Figure S2 in the Supporting Information). Not surprisingly, the $[\mathbf{4}]\text{OTf}$ spectrum closely resembles the one published for $[\text{Fe}(4\text{-TIP}^{\text{Ph}})(\text{acac}^{\text{PhF}3})]\text{OTf}$, although the absorption features are blueshifted in the former by approximately 400 cm^{-1} .[19](#)

Crystals of $[\mathbf{4}]\text{OTf}$ were obtained from the reaction of $[\mathbf{1}](\text{OTf})_2$ and $\text{Na}(\text{acac}^{\text{PhF}3})$ in CH_2Cl_2 , followed by crystallization in $\text{CH}_2\text{Cl}_2/\text{pentane}$. The asymmetric unit contains two independent units with virtually identical structures. As shown in Figure [3](#), the 5C Fe^{2+} center is coordinated to the 2-TIP^{Ph} and $\text{acac}^{\text{PhF}3}$ ligands in a distorted trigonal-bipyramidal geometry ($\tau = 0.51$) with the O atom proximal to the CF_3 group (O1) in the axial position. The metric parameters of $[\mathbf{4}]\text{OTf}$ are not significantly different from those reported previously for $[\text{Fe}^{2+}(\text{Ph}2\text{Tp})(\text{acac}^{\text{PhF}3})]$ and $[\text{Fe}^{2+}(4\text{-TIP}^{\text{Ph}})(\text{acac}^{\text{PhF}3})]\text{OTf}$.[19](#)




Figure 3

Thermal ellipsoid plot (50 % probability) derived from $[\mathbf{4}]\text{OTf} \cdot 2\text{CH}_2\text{Cl}_2$. Non-coordinating solvent molecules, counteranions, and most H atoms have been omitted for clarity. Only one of the two independent $[\mathbf{5}]^+$ units is shown. Selected bond lengths [Å] and angles [°] for this unit: Fe1–O1 2.089(3), Fe1–O2 1.973(3), Fe1–N2 2.118(4), Fe1–N4 2.190(4), Fe1–N6 2.118(4); O1–Fe1–O2 87.2(1), O1–Fe1–N2 91.0(2), O1–Fe1–N4 176.4(2), O1–Fe1–N6 89.5(2), O2–Fe1–N2 120.4(2), O2–Fe1–N4 96.5(2), O2–Fe1–N6 146.3(2).

The solution structures of $[\mathbf{2}]\text{OTf}$ and $[\mathbf{4}]\text{OTf}$ in CD_2Cl_2 were probed using ^1H NMR spectroscopy, and the observed chemical shifts, peak integrations, and T_1 values are summarized in Table [3](#). The three imidazole ligands are spectroscopically equivalent in solution due to dynamic averaging of the ligand positions on the NMR spectroscopic timescale. The 2-TIP^{Ph2}-derived resonances were assigned with the help of peak integrations and by making two assumptions: (i) T_1 values follow the order *ortho* < *meta* < *para* for each phenyl ring,[13c,19](#) and (ii) T_1 values of the 4-Ph protons are shorter


than the corresponding protons on the 5-Ph ring. Thus, the fast-relaxing peaks ($T_1 \approx 1$ ms) near -20 ppm were attributed to the *ortho* protons of the 4-phenyl 2-TIP^{Ph2} substituents, which are positioned near the Fe²⁺ center. The peaks with the largest integration at (21 ± 1) ppm were assigned to the 1-*N*-Me protons. The remaining resonances were then identified as the benzoate and acac^{PhF3} groups of [2]OTf and [4]OTf, respectively, by using the relative T_1 values to assign the phenyl resonances of both ligands.

Table 3. Summary of ¹H NMR spectroscopic parameters for [2]OTf and [4]OTf in CD₂Cl₂.

[2]OTf			[4]OTf		
Resonance	δ [ppm]	T_1 [ms]	Resonance	δ [ppm]	T_1 [ms]
<i>o</i> -4-Ph	-21.0	1.1	<i>o</i> -4-Ph	-16.0	0.4
<i>m</i> -4-Ph	6.7	12.0	<i>m</i> -4-Ph	5.2	4.7
<i>p</i> -4-Ph	9.0	31.6	<i>p</i> -4-Ph	9.3	13.3
<i>o</i> -5-Ph	2.6	31.5	<i>o</i> -5-Ph	2.4	13.7
<i>m</i> -5-Ph	6.3	159	<i>m</i> -5-Ph	6.1	88.2
<i>p</i> -5-Ph	5.2	238	<i>p</i> -5-Ph	5.0	120
<i>N</i> -1-Me	21.9	15.7	<i>N</i> -1-Me	20.2	5.9
<i>o</i> -OBz	34.8	3.2	acac <i>o</i> -Ph	22.6	1.7
<i>m</i> -OBz	19.0	37.0	acac <i>m</i> -Ph	9.5	20.0
<i>p</i> -OBz	10.6	67.6	acac <i>p</i> -Ph	17.3	45.5
			acac H	39.4	0.8

Fe²⁺ Complexes Containing 4-TIP^{Ph}

The complex [Fe(4-TIP^{Ph})(OAc)(MeOH)]BPh₄ ([5]BPh₄) was generated by addition of NaBPh₄ to a solution of Fe(OAc)₂ and 4-TIP^{Ph} in MeOH, which resulted in the immediate formation of a white precipitate (Scheme 4). The IR spectrum of the isolated solid revealed a peak at 3259 cm⁻¹ from the ν (N-H) stretch of the 4-TIP^{Ph} ligands, along with acetate-derived features at 1562 and 1402 cm⁻¹. The 4-TIP^{Ph}-derived resonances in the ¹H NMR spectrum largely followed the pattern reported previously for [Fe²⁺(4-TIP^{Ph})(acac^X)]⁺ complexes.¹⁹ The acetate ligand of [5]BPh₄ exhibited a downfield signal at $\delta = +105$ ppm.



Scheme 4

X-ray-quality crystals of [5]BPh₄ were prepared by slowly cooling a solution of [5]BPh₄ in MeOH; the [5]⁺ cation is shown in Figure 4 and the corresponding bond lengths and angles are provided in Table 2.

The high-spin Fe²⁺ center is hexacoordinate with a facially coordinating 4-TIP^{Ph} ligand. The Fe–N_{TIP} distances in [5]⁺ are quite similar to those found for [1]²⁺ and [2]⁺, which suggests that the 4-TIP^{Ph} and 2-TIP^{Ph2} ligands possess comparable donor properties. The κ²-acetate ligand coordinates in a symmetric manner with nearly identical Fe–O_{acetate} distances of 2.251(6) Å. The remaining site is occupied by a solvent molecule *trans* to N4 with a relatively short Fe–O_{MeOH} distance of 2.077(4) Å. The crystal structure of [5]BPh₄·3MeOH also features an extensive hydrogen-bonding network. As shown in Figure 4, the coordinated acetate and MeOH moieties participate in hydrogen-bonding interactions with three MeOH “chaperones” that comprise a second-sphere shell surrounding one face of the [5]⁺ octahedron. In addition, the MeOH molecules that serve as hydrogen-bond donors to the acetate ligand also act as hydrogen-bond acceptors for two H–N_{imidazole} groups on adjacent [5]⁺ cations.




Figure 4

Thermal ellipsoid plot (50 % probability) derived from [5]BPh₄·3MeOH. The BPh₄ counteranion and most H atoms have been omitted for clarity. The dotted lines signify the hydrogen-bonding interactions between the coordinated acetate and MeOH ligands and three second-sphere solvent molecules. Note: Ellipsoids are not shown for the proximal 2-Ph substituent due to disorder.

Significantly, we found that [5]BPh₄ provides access to iron(II) salicylate (sal) species that mimic the enzyme–substrate complex of SDO. The complex [Fe(4-TIP^{Ph})(sal)] (6) was prepared by mixing [5]BPh₄ with salicylic acid (1 equiv.) in MeOH, followed by layering with MeCN (Scheme 4). As shown in Figure 5, the X-ray crystal structure of 6 reveals a neutral 5C Fe²⁺ complex with a geometry between square pyramidal and trigonal bipyramidal ($\tau = 0.35$). The dianionic salicylate ligand coordinates in a bidentate fashion with Fe–O bond lengths of 1.958(1) and 2.060(1) Å for the phenolate and carboxylate donors, respectively. To the best of our knowledge, 6 represents the first structurally characterized iron(II) salicylate complex in the chemical literature.²⁶





Figure 5

Thermal ellipsoid plot (50 % probability) derived from **6**·MeOH·MeCN. The noncoordinating MeCN and most H atoms have been omitted for clarity. The dotted line represents the hydrogen-bonding interaction between the salicylate ligand and MeOH. Selected bond lengths [Å] and angles [°]: Fe1–O1 2.060(1), Fe1–O3 1.958(1), Fe1–N2 2.135(1), Fe1–N4 2.150(1), Fe1–N6 2.183(1), O1–C28 1.257(2), O2–C28 1.273(2), O3–C30 1.327(2), O3···O1S 2.725(1); O1–Fe1–O3 86.29(4), O1–Fe1–N2 92.17, O1–Fe1–N4 96.90(4), O1–Fe1–N6 168.66(4), O3–Fe1–N2 147.75(4), O3–Fe1–N4 117.46(4), O3–Fe1–N6 91.71(4).

As with [5]BPh₄·3MeOH, the lattice of **6** exhibits numerous hydrogen-bonding interactions (see Scheme 5). The uncoordinated oxygen atom of the carboxylate (O2) forms hydrogen bonds with two H–N groups belonging to adjacent 4-TIP^{Ph} ligands. These interactions account for the fact that O2–C28 is unexpectedly longer than O1–C28 [1.273(2) vs. 1.257(2), respectively], which indicates that the negative charge is delocalized over the carboxylate moiety. The crystal also contains noncoordinating MeCN and MeOH molecules (one of each); the latter serves as a hydrogen-bond donor to the phenolate oxygen atom (O3) of the salicylate, while acting as a hydrogen-bond acceptor to an imidazole H–N group. Thus, in this structure, MeOH behaves in a manner similar to second-sphere residues in dioxygenase active sites, which often play a crucial role in stabilizing metal-bound substrates through noncovalent interactions.²⁷



Scheme 5

Hydrogen-bonding network in the solid-state structure of **6**.

Conclusion

This paper has described the synthesis and X-ray structural characterization of iron(II) complexes supported by tris(imidazolyl)phosphane ligands (2-TIP^{Ph2} and 4-TIP^{Ph}). Three of the complexes – [1](OTf)₂, [2]OTf, and [5]BPh₄ – feature easily displaced ligands, such as solvent molecules and/or carboxylates, in the coordination sites *trans* to the TIP chelate. These complexes exhibit variability in their coordination numbers (5C or 6C) and carboxylate binding modes (κ^1 or κ^2). Intra- and intermolecular hydrogen-bonding interactions between the ligands and solvent are evident in the solid-state structures of each complex {with the exception of [1](OTf)₂} In particular, the presence of unprotected imidazole groups in [5]BPh₄ gives rise to an extensive hydrogen-bonding network in which second-sphere MeOH molecules form bridges between acetate ligands and H–N_{imid} groups from neighboring [5]⁺ units.

Like the resting states of the enzymatic active sites, these “precursor” complexes are intended to serve as scaffolds that permit various substrate ligands to coordinate to the iron(II) center. The versatility of this approach was demonstrated by the formation of the Dke1 model [4]OTf from the reaction of Na(acac^{PhF3}) with the 2-TIP^{Ph2}-based complexes [1](OTf)₂ and [2]OTf. Similarly, the SDO model **6** was generated through the direct reaction of [5]BPh₄ with salicylic acid. The facile formation of [4]OTf and **6** indicates that the TIP ligands are resistant to displacement by strong, anionic ligands. This is significant because half-sandwich ferrous complexes with neutral L_{N3} ligands, such as trispyrazolylmethanes, have been shown to suffer from high lability and a tendency to decompose to the more stable bis-ligand species.²¹ The relatively short Fe–N_{TIP} bond lengths found in our series of complexes suggest that the TIP ligands bind tightly to the iron centers. Thus, the precursor complexes described here provide a robust platform for the development of synthetic models of dioxygenases with the 3His facial triad.

Experimental Section

General Procedures: All reagents and solvents were purchased from commercial sources and used as received unless otherwise noted. MeCN and CH₂Cl₂ were purified and dried using a Vacuum Atmospheres solvent purification system. The compounds 4,5-diphenyl-1-methylimidazole²⁸ and 4-TIP^{Ph}^[16h] were prepared according to literature procedures. The synthesis and handling of air-sensitive materials were carried out under an inert atmosphere using a Vacuum Atmospheres Omni-Lab glovebox equipped with a freezer set to -30 °C. Elemental analyses were performed at Midwest Microlab, LLC in Indianapolis, IN. Infrared (IR) spectra of solid samples were measured with a Thermo Scientific Nicolet iS5 FTIR spectrometer equipped with the iD3 attenuated total reflectance accessory. UV/Vis spectra were obtained with an Agilent 8453 diode array spectrometer. NMR spectra were recorded on a Varian 400 MHz spectrometer. ¹⁹F NMR spectra were referenced using the benzotrifluoride peak at -63.7 ppm. ³¹P NMR spectra were referenced to external H₃PO₄ (δ = 0 ppm). Magnetic susceptibility measurements were carried out using the Evans NMR method.

2-TIP^{Ph2}: 4,5-Diphenyl-1-methylimidazole (6.81 g, 29.1 mmol) was dissolved in THF (175 mL) and the solution was purged with argon for 25 min. The flask was cooled to -78 °C and *n*BuLi (32.0 mmol) was added dropwise. The solution was stirred for 30 min at -78 °C and then for 30 min at room temperature. The reaction was cooled again to -78 °C and PCl₃ (0.850 mL, 9.74 mmol) was added slowly. The mixture was allowed to slowly warm to room temp. over the course of several hours, and then 30 % NH₄OH (75 mL) was added and stirred for 1 h. The layers were separated and the aqueous layer was extracted with THF (2 × 35 mL). The combined THF layers were washed with H₂O and brine (50 mL each), dried with MgSO₄, and the solvent was removed under vacuum. The orange residue was triturated with pentane and washed with methanol, thereby providing a fine white powder (1.66 g); yield 24 %. C₄₈H₃₉N₆P (730.8): calcd. C 78.88, H 5.38, N 11.50; found C 78.05, H 5.83, N 11.03. The disagreement indicates that small amounts of impurities are present. ¹H NMR (400 MHz, CDCl₃): δ = 7.48 (m, 12 H, Ar-H), 7.40 (m, 6 H, Ar-H), 7.17 (m, 12 H, Ar-H), 3.64 (s, 9 H, CH₃) ppm. ¹³C NMR (100 MHz, CDCl₃): δ = 140.5, 140.0, 139.9, 134.8, 133.6, 131.1, 131.0, 129.2, 129.0, 128.2, 126.9, 126.5, 33.4 ppm. ³¹P NMR (162 MHz, CDCl₃): δ = -56.6 ppm. IR (neat): $\bar{\nu}$ = 3053, 2940, 2863, 1601, 1503, 1442, 1363, 1071, 1024, 961 cm⁻¹.

[Fe(2-TIP^{Ph2})(MeCN)₃](OTf)₂ {[1](OTf)₂]: 2-TIP^{Ph2} (1.32 g, 1.81 mmol) and Fe(OTf)₂ (670 mg, 1.90 mmol) were mixed in CH₃CN (20 mL) and stirred until the solution had become clear (about 3 h). The solution was filtered and layered with excess Et₂O; X-ray-quality crystals formed after one day. The white crystals were collected and dried under vacuum to provide 1.31 g of material; yield 60 %. Elemental analysis showed that at least two of the coordinated CH₃CN ligands are removed upon drying. C₅₀H₃₉F₆FeN₆O₆PS₂·CH₃CN (1125.9): calcd. C 55.47, H 3.76, N 8.71; found C 55.02, H 3.90, N 8.68. IR (neat): $\bar{\nu}$ = 3048, 2932, 2283 [v(C≡N)], 1466, 1444, 1257, 1222, 1145, 1028, 983 cm⁻¹.

[Fe(2-TIP^{Ph2})(OBz)(MeOH)]OTf ([2]OTf): 2-TIP^{Ph2} (779 mg, 1.07 mmol), NaOBz (155 mg, 1.07 mmol), and Fe(OTf)₂ (378 mg, 1.07 mmol) were combined in MeOH (12 mL). After stirring for several hours the precipitate was removed by filtration and the filtrate was reduced to about 5 mL in volume. Layering with pentane afforded the desired product as a white crystalline material (116 mg); yield 11 %. X-ray diffraction analysis revealed four uncoordinated MeOH molecules in the resulting structure, and elemental analysis indicated that two solvent molecules remain after drying under vacuum.

C₅₇H₄₇F₃FeN₆O₆PS·2CH₃OH (1152.0): calcd. C 61.51, H 4.81, N 7.30; found C 61.35, H 4.48, N 7.07. IR (neat): $\bar{\nu}$ = 3043, 2953, 1598, 1551, 1443, 1370, 1258, 1153, 1029, 981 cm⁻¹.

[Fe(2-TIP^{Ph2})(acac^{PhF3})]OTf ([4]OTf): A solution of 4,4,4-trifluoro-1-phenyl-1,3-butanedione (126 mg, 0.584 mmol) and NaOCH₃ (32 mg, 0.59 mmol) in THF was stirred for 30 min, after which the solvent was removed under vacuum to give white Na(acac^{PhF3}). Na(acac^{PhF3}) was then dissolved in CH₃CN (5 mL) and slowly added to a solution of [1](OTf)₂ (704 mg, 0.583 mmol) in CH₂Cl₂ (5 mL). The purple solution was stirred overnight and the solvent was removed under vacuum. The residue was dissolved in CH₂Cl₂ (5 mL), filtered, and layered with pentane to yield deep red crystals suitable for X-ray crystallography (457 mg); yield 68 %. The X-ray structure revealed uncoordinated CH₂Cl₂ molecules in the asymmetric units, and elemental analysis suggests that a small amount of solvent (≈0.7 equiv.) remains after vacuum drying. C₅₉H₄₅F₆FeN₆O₅PS·0.7CH₂Cl₂ (1210.36): calcd. C 59.24, H 3.86, N 6.94; found C 59.25, H 3.99, N 6.75. UV/Vis (MeCN): λ_{\max} (ϵ , M⁻¹ cm⁻¹) = 519 (720), 494 (730) nm. IR (neat): $\bar{\nu}$ = 3058, 2955, 1602 [ν (C=O)], 1572, 1462, 1443, 1253, 1141, 1029, 981 cm⁻¹. ¹⁹F NMR (376 MHz, CD₂Cl₂): δ = -44.9 (acac^{PhF3}), -77.7 (OTf) ppm.

[Fe(4-TIP^{Ph})(OAc)(MeOH)]BPh₄ ([5]BPh₄): Fe(OAc)₂ (488 mg, 2.81 mmol) and 4-TIP^{Ph} (1.28 g, 2.79 mmol) were stirred in MeOH (10 mL) for 10 min while the solution became clear. A solution of NaBPh₄ (956 mg, 2.79 mmol) in MeOH was then added dropwise and the mixture was stirred for 5 h. During this time, a white precipitate developed. The white solid was collected and recrystallized from MeOH at -30 °C; yield 48 %. Elemental analysis indicates that the bound MeOH ligand only partially (50 %) occupied the complex in the ground, vacuum-dried solid. C₅₃H₄₄BF₄FeN₆O₂P·0.5MeOH (910.6): calcd. C 70.57, H 4.98, N 9.23; found C 70.69, H 5.08, N 8.95. IR (neat): $\bar{\nu}$ = 3304, 3259 [ν (N-H)], 3054, 2999, 2993, 2928, 1562 [ν_{as} (OCO)], 1478, 1402 [ν_{s} (OCO)], 1341 cm⁻¹.

[Fe(4-TIP^{Ph})(sal)] (6): A suspension of [5]BPh₄ (142 mg, 0.159 mmol) and sodium salicylate (28.0 mg, 0.175 mmol) was stirred overnight in MeOH (5 mL). The resulting yellow solution was layered with MeCN to provide X-ray-quality crystals of **6**; yield 32 %. C₃₄H₂₅FeN₆O₃P (652.4): calcd. C 62.59, H 3.86, N 12.88; found C 62.19, H 3.98, N 12.52. UV/Vis (MeOH): λ_{\max} (ϵ , M⁻¹ cm⁻¹) = 440 (150) nm. IR (neat): $\bar{\nu}$ = 3133, 3052, 2900, 1598, 1563, 1521, 1476, 1458, 1439, 1386, 1314 cm⁻¹.

X-ray Structure Determination: XRD data were collected at 100 K with an Oxford Diffraction SuperNova kappa-diffractometer (Agilent Technologies) equipped with dual microfocus Cu/Mo X-ray sources, X-ray mirror optics, Atlas CCD detector, and low-temperature Cryojet device. Crystallographic data for particular compounds are summarized in Table 4. The data were analyzed with the CrysAlis Pro program package (Agilent Technologies, 2011) typically using a numerical Gaussian absorption correction (based on the real shape of the crystal), followed by an empirical multiscan correction using the SCALE3 ABSPACK routine. The structures were solved using the SHELXS program and refined with the SHELXL program²⁹ within the Olex2 crystallographic package.³⁰ B-, H-, and C-bonded hydrogen atoms were positioned geometrically and refined using appropriate geometric restrictions on the corresponding bond lengths and bond angles within a riding/rotating model (torsion angles of methyl hydrogen atoms were rotationally optimized to better fit the residual electron density). The positions of the methanolic hydrogen atoms (H3) in [2]OTf·4MeOH and [5]BPh₄·3MeOH were refined freely. The remaining OH groups were refined using geometrical restrictions and rotationally optimized to better fit the residual electron density. Crystals of [4]OTf·2CH₂Cl₂ represent pseudo-orthorhombic quasi-

merihedral twins ($\beta \approx 90^\circ$). Crystals of [5]BPh₄·3MeOH are systematic twins grown together along a common *bc* plane. The chiral space group (*P*2₁) of [5]BPh₄·3MeOH does not result from the molecular chirality of the cation, but rather from crystal packing. The cation itself has an approximate local mirror symmetry (in the direction perpendicular to crystallographic *z* axis). The apparent result of this local symmetry is the observed twinning. The twinning, however, annihilates the chirality on the macroscopic level since the components of the twin are of opposite chirality.

Table 4. Summary of the X-ray crystallographic data collection and structure refinement.

	[1](OTf) ₂ ·0.5Et 2O	[2]OTf·4MeO H	[4]OTf·2CH ₂ Cl ₂ [a]	[5]BPh ₄ ·3M eOH	6·MeCN·M eOH
Empirical formula	C ₅₈ H ₅₃ F ₆ FeN ₉ O 6.5PS ₂	C ₆₁ H ₆₄ F ₃ FeN ₆ O ₁₀ PS	C ₆₁ H ₄₉ Cl ₄ F ₆ FeN 6O ₅ PS	C ₅₇ H ₆₀ BFeN ₆ O ₆ P	C ₃₇ H ₃₂ FeN ₇ O ₄ P
Formula weight	1245.03	1217.06	1320.77	1022.74	725.52
Crystal system	triclinic	triclinic	monoclinic	monoclinic	monoclinic
Space group	<i>P</i> $\bar{1}$	<i>P</i> $\bar{1}$	<i>Pc</i>	<i>P</i> 2 ₁	<i>P</i> 2 ₁ / <i>n</i>
<i>a</i> [Å]	13.3432(3)	14.9470(5)	16.0689(5)	13.8829(3)	13.6187(7)
<i>b</i> [Å]	15.8007(3)	15.1921(5)	20.6668(5)	11.6385(4)	14.9164(9)
<i>c</i> [Å]	27.8621(6)	16.4349(6)	19.6239(4)	16.5130(4)	17.5278(8)
α [°]	76.994(2)	90.642(3)	90	90	90
β [°]	88.757(2)	113.784(3)	90.088(2)	91.591(2)	102.190(5)
γ [°]	87.690(2)	115.873(4)	90	90	90
<i>V</i> [Å ³]	5718.4(2)	2992.3(3)	6516.9	2667.1(2)	3480.4(3)
<i>Z</i>	4	2	4	2	4
<i>D</i> _{calcd.} [g cm ³]	1.446	1.351	1.325	1.274	1.385
λ [Å]	1.5418	1.5418	0.7107	1.5418	1.5418
μ [mm ⁻¹]	3.749	3.205	0.489	2.996	4.328
ϑ range [°]	7 to 149	7 to 148	7 to 59	4 to 148	4 to 149
Reflections collected	64740	39217	56565	32914	26690
Independent reflections	22662	11891	28249	10151	6954
	(<i>R</i> _{int} = 0.0315)	(<i>R</i> _{int} = 0.0299)	(<i>R</i> _{int} = 0.0339)	(<i>R</i> _{int} = 0.1419)	(<i>R</i> _{int} = 0.0278)
Data/restraints/parameters	22662/19/159 1	11891/2/784	28249/12/155 0	10151/87/6 43	6954/0/45 4
GOF (on <i>F</i> ²)	1.025	1.028	1.061	1.025	1.037
<i>R</i> 1/ <i>wR</i> 2 [<i>I</i> > 2σ(<i>I</i>)] [b]	0.0332/0.0855	0.0413/0.109 4	0.0608/0.1541	0.0682/0.17 78	0.0273/0.0 710
<i>R</i> 1/ <i>wR</i> 2 (all data)	0.0374/0.0886	0.0449/0.112 8	0.0666/0.1613	0.0872/0.19 63	0.0306/0.0 731

[a] One of the solvates in [4]OTf·2CH₂Cl₂ is partially occupied by a pentane molecule.

[b] $R1 = \sum ||F_o| - |F_c|| / \sum |F_o|$; $wR2 = [\sum w(F_o^2 - F_c^2)^2 / \sum w(F_o^2)^2]^{1/2}$.

Acknowledgements

This research was supported by the National Science Foundation (NSF) (CAREER CHE-1056845) and Marquette University.

References

- 1 1a M. Costas , M. P. Mehn , M. P. Jensen , L. Que Jr , *Chem. Rev.* 2004 , **104** , 939 – 986. 1b E. I. Solomon , T. C. Brunold , M. I. Davis , J. N. Kemsley , S.-K. Lee , N. Lehnert , F. Neese , A. J. Skulan , Y.-S. Yang , J. Zhou , *Chem. Rev.* 2000 , **100** , 235 – 349. 1c D. T. Gibson , R. E. Parales , *Curr. Opin. Biotechnol.* 2000 , **11** , 236 – 243.
- 2 2a T. D. H. Bugg , *Curr. Opin. Chem. Biol.* 2001 , **5** , 550 – 555. 2b T. D. H. Bugg , G. Lin , *Chem. Commun.* 2001 , **11** , 941 – 953. 2c E. G. Kovaleva , J. D. Lipscomb , *Nature Chem. Biol.* 2008 , **4** , 186 – 193.
- 3 3a B. Kauppi , K. Lee , E. Carredano , R. E. Parales , D. T. Gibson , H. Eklund , S. Ramaswamy , *Structure* 1998 , **6** , 571 – 586. 3b A. Karlsson , J. V. Parales , R. E. Parales , D. T. Gibson , H. Eklund , S. Ramaswamy , *Science* 2003 , **299** , 1039 – 1042.
- 4 4a G. P. Titus , H. A. Mueller , J. Burgner , S. R. d. Córdoba , M. A. Penalva , D. E. Timm , *Nat. Struct. Biol.* 2000 , **7** , 542 – 546. 4b F. H. Vaillancourt , J. T. Bolin , L. D. Eltis , *Crit. Rev. Biochem. Mol. Biol.* 2006 , **41** , 241 – 267.
- 5 5a T. E. Machonkin , A. E. Doerner , *Biochemistry* 2011 , **50** , 8899 – 8913. 5b T. E. Machonkin , P. L. Holland , K. N. Smith , J. S. Liberman , A. Dinescu , T. R. Cundari , S. S. Rocks , *J. Biol. Inorg. Chem.* 2010 , **15** , 291 – 301. 5c Y. Yin , N. Y. Zhou , *Curr. Microbiol.* 2010 , **61** , 471 – 476. 5d L. Y. Xun , J. Bohuslavsek , M. A. Cai , *Biochem. Biophys. Res. Commun.* 1999 , **266** , 322 – 325.
- 6 6a K. D. Koehntop , J. P. Emerson , L. Que Jr , *J. Biol. Inorg. Chem.* 2005 , **10** , 87 – 93. 6b Arianna Bassan , Tomasz Borowski , P. E. M. Siegbahn , *Dalton Trans.* 2004 , 1353 – 1362.
- 7 7a G. D. Straganz , B. Nidetzky , *ChemBioChem* 2006 , **7** , 1536 – 1548. 7b S. Leitgeb , B. Nidetzky , *Biochem. Soc. Trans.* 2008 , **36** , 1180 – 1186.
- 8 8a C. R. Simmons , Q. Liu , Q. Huang , Q. Hao , T. P. Begley , P. A. Karplus , M. H. Stipanuk , *J. Biol. Chem.* 2006 , **281** , 18723 – 18733. 8b C. A. Joseph , M. J. Maroney , *Chem. Commun.* 2007 , 3338 – 3349.
- 9 9a G. Straganz , L. Brecker , H. J. Weber , W. Steiner , D. W. Ribbons , *Biochem. Biophys. Res. Commun.* 2002 , **297** , 232 – 236. 9b G. D. Straganz , A. Glieder , L. Brecker , D. W. Ribbons , W. Steiner , *Biochem. J.* 2003 , **369** , 573 – 581.
- 10 10a J. Chen , W. Li , M. Z. Wang , G. Y. Zhu , D. Q. Liu , F. Sun , N. Hao , X. M. Li , Z. H. Rao , X. C. Zhang , *Protein Sci.* 2008 , **17** , 1362 – 1373. 10b M. R. Harpel , J. D. Lipscomb , *J. Biol. Chem.* 1990 , **265** , 22187 – 22196.
- 11 11a I. Matera , M. Ferraroni , S. Burger , A. Scozzafava , A. Stolz , F. Briganti , *J. Mol. Biol.* 2008 , **380** , 856 – 868. 11b J. P. Hintner , T. Remtsma , A. Stolz , *J. Biol. Chem.* 2004 , **279** , 37250 – 37260. 11c J. P. Hintner , C. Lechner , U. Riegert , A. E. Kuhm , T. Storm , T. Reemtsma , A. Stolz , *J. Bacteriol.* 2001 , **183** , 6936 – 6942.
- 12 P. C. A. Bruijninx , G. van Koten , R. J. M. K. Gebbink , *Chem. Soc. Rev.* 2008 , **37** , 2716 – 2744.
- 13 13a N. Kitajima , N. Tamura , H. Amagai , H. Fukui , Y. Moro-oka , Y. Mizutani , T. Kitagawa , R. Mathur , K. Heerwegh , C. A. Reed , C. R. Randall , L. Que Jr , K. Tatsumi , *J. Am. Chem. Soc.* 1994 , **116** , 9071 – 9085. 13b E. H. Ha , R. Y. N. Ho , J. F. Kesiel , J. S. Valentine , *Inorg. Chem.* 1995 , **34** , 2265 – 2266. 13c M. P. Mehn , K. Fujisawa , E. L. Hegg , L. Que Jr , *J. Am. Chem. Soc.* 2003 , **125** , 7828 – 7842. 13d A. Mukherjee , M. A. Cranswick , M. Chakrabarti , T. K. Paine , K. Fujisawa , E. Munck , L. Que , *Inorg. Chem.* 2010 , **49** , 3618 – 3628. 13e T. Ogihara , S. Hikichi , M. Akita , Y. Moro-oka , *Inorg. Chem.* 1998 , **37** , 2614 – 2615.
- 14 14a N. Burzlaff , in *Advances in Inorganic Chemistry*, Vol. 60 (Ed.: R. van Eldik) , 2008 , pp. 101–165. 14b A. Beck , B. Weibert , N. I. Burzlaff , *Eur. J. Inorg. Chem.* 2001 , 521 – 527. 14c A. Beck , A. Barth , E. Hulbner , N. Burzlaff , *Inorg. Chem.* 2003 , **42** , 7182 – 7188.

- 15 P. C. A. Bruijninx, M. Lutz, A. L. Spek, W. R. Hagen, B. M. Weckhuysen, G. van Koten, R. J. M. K. Gebbink, *J. Am. Chem. Soc.* 2007, **129**, 2275 – 2286.
- 16 16a R. S. Brown, J. Huguet, *Can. J. Chem.* 1980, **58**, 889 – 901. 16b R. Breslow, J. T. Hunt, R. Smiley, T. Tarnowski, *J. Am. Chem. Soc.* 1983, **105**, 5337 – 5342. 16c H. Slebocka-Tilk, J. L. Cocho, Z. Frakman, R. S. Brown, *J. Am. Chem. Soc.* 1984, **106**, 2421 – 2431. 16d W. E. Allen, T. N. Sorrell, *Inorg. Chem.* 1997, **36**, 1732 – 1734. 16e C. Kimblin, W. E. Allen, G. Parkin, *J. Chem. Soc., Chem. Commun.* 1995, 1813 – 1815. 16f W. E. Lynch, J. D. M. Kurtz, S. Wang, R. A. Scott, *J. Am. Chem. Soc.* 1994, **116**, 11030 – 11038. 16g P. C. Kunz, G. J. Reiss, W. Frank, W. Klaui, *Eur. J. Inorg. Chem.* 2003, 3945 – 3951. 16h P. C. Kunz, W. Klaui, *Collect. Czech. Chem. Commun.* 2007, **72**, 492 – 502.
- 17 17a A. Y. S. Malkhasian, B. Nikolovski, B. E. Kucera, R. Loloee, F. A. Chavez, *Z. Anorg. Allg. Chem.* 2007, **633**, 1000 – 1005. 17b M. P. Batten, A. J. Canty, K. J. Cavell, T. Ruther, B. W. Skelton, A. H. White, *Acta Crystallogr., Sect. C* 2004, **60**, M311–M313. 17c F.-J. Wu, D. M. Kurtz Jr, *J. Am. Chem. Soc.* 1989, **111**, 6563 – 6572.
- 18 18a F. J. Wu, D. M. Kurtz, K. S. Hagen, P. D. Nyman, P. G. Debrunner, V. A. Vankai, *Inorg. Chem.* 1990, **29**, 5174 – 5183. 18b V. A. Vankai, M. G. Newton, D. Kurtz Jr, *Inorg. Chem.* 1992, **31**, 342 – 343.
- 19 H. Park, J. S. Baus, S. V. Lindeman, A. T. Fiedler, *Inorg. Chem.* 2011, **50**, 11978 – 11989.
- 20 20a G. D. Straganz, A. R. Diebold, S. Egger, B. Nidetzky, E. I. Solomon, *Biochemistry* 2010, **49**, 996 – 1004. 20b S. Leitgeb, G. D. Straganz, B. Nidetzky, *Biochem. J.* 2009, **418**, 403 – 411.
- 21 P. G. Edwards, A. Harrison, P. D. Newman, W. J. Zhang, *Inorg. Chim. Acta* 2006, **359**, 3549 – 3556.
- 22 L. H. Gade, G. Marconi, C. Dro, B. D. Ward, M. Poyatos, S. Bellemin-Lapponnaz, H. Wadepohl, L. Sorace, G. Poneti, *Chem. Eur. J.* 2007, **13**, 3058 – 3075.
- 23 A. W. Addison, T. N. Rao, J. Reedijk, J. Vanriijn, G. C. Verschoor, *J. Chem. Soc., Dalton Trans.* 1984, 1349 – 1356.
- 24 K. Fujisawa, N. Tada, Y. Nishida, Y. Miyashita, K. Okamoto, *Inorg. Chem. Commun.* 2008, **11**, 381 – 384.
- 25 25a G. D. Straganz, H. Hofer, W. Steiner, B. Nidetzky, *J. Am. Chem. Soc.* 2004, **126**, 12202 – 12203. 25b G. D. Straganz, B. Nidetzky, *J. Am. Chem. Soc.* 2005, **127**, 12306 – 12314.
- 26 Ferric salicylate complexes have been reported in the following papers, see:
- 26a M. R. McDevitt, A. W. Addison, E. Sinn, L. K. Thompson, *Inorg. Chem.* 1990, **29**, 3425 – 3433. 26b N. Y. Oh, M. S. Seo, M. H. Lim, M. B. Consugar, M. J. Park, J.-U. Rohde, J. Han, K. M. Kim, J. Kim, L. Que Jr, W. Nam, *Chem. Commun.* 2005, 5644 – 5646. 26c S. Taktak, M. Flook, B. M. Foxman, L. Que Jr, E. V. Rybak-Akimova, *Chem. Commun.* 2005, 5301 – 5303. 26d O. V. Makhlynets, P. Das, S. Taktak, M. Flook, R. Mas-Balleste, E. V. Rybak-Akimova, L. Que, *Chem. Eur. J.* 2009, **15**, 13171 – 13180.
- 27 27a S. L. Groce, J. D. Lipscomb, *Biochemistry* 2005, **44**, 7175 – 7188. 27b J. D. Lipscomb, *Curr. Opin. Struct. Biol.* 2008, **18**, 644 – 649.
- 28 F. J. LaRonde, M. A. Brook, *Inorg. Chim. Acta* 1999, **296**, 208 – 221.
- 29 G. M. Sheldrick, *Acta Crystallogr., Sect. A* 2008, **64**, 112 – 122.
- 30 O. V. Dolomanov, L. J. Bourhis, R. J. Gildea, J. A. K. Howard, H. Puschmann, *J. Appl. Crystallogr.* 2009, **42**, 339 – 341.