Marquette University

e-Publications@Marquette

Biological Sciences Faculty Research and
Publications

Biological Sciences, Department of

1-28-2010

A New Species of Gekko (Squamata: Gekkonidae) from Tà Kóu Nature Reserve, Binh Thuan Province, Southern Vietnam

Van Tri Ngo

Vietnamese Academy of Sciences and Technology

Tony Gamble

Marquette University, anthony.gamble@marquette.edu

Follow this and additional works at: https://epublications.marquette.edu/bio_fac

Part of the [Biology Commons](#)

Recommended Citation

Ngo, Van Tri and Gamble, Tony, "A New Species of Gekko (Squamata: Gekkonidae) from Tà Kóu Nature Reserve, Binh Thuan Province, Southern Vietnam" (2010). *Biological Sciences Faculty Research and Publications*. 749.

https://epublications.marquette.edu/bio_fac/749

Published in final edited form as:

Zootaxa. 2010 January 28; 2346: 17–28.

A new species of *Gekko* (Squamata: Gekkonidae) from Tà Kóu Nature Reserve, Binh Thuan Province, Southern Vietnam

Ngo Van Tri¹ and Tony Gamble²

¹Department of Environmental Management and Technology, Institute of Tropical Biology, Vietnamese Academy of Sciences and Technology, 85 Tran Quoc Toan Street, District 3, Hochiminh City, Vietnam. trigeckonid@hotmail.com

²Department of Genetics, Cell Biology and Development, University of Minnesota, 6-160 Jackson Hall, 321 Church St SE, Minneapolis MN 55455. USA, gamb007@umn.edu

Abstract

A new species of *Gekko* Laurenti is described from Ta Kou Mountain, an isolated granitic peak in Ta Kou Nature Reserve, Ham Thuan Nam district, Binh Thuan province, southern Vietnam. The species is distinguished from its congeners by its moderate size, with snout to vent length (SVL) reaching a maximum 107.0 mm; dorsal pattern of 5–8 white vertebral blotches between the nape and sacrum and 6–8 pairs of short white bars on the flanks; 11–14 precloacal pores in males; 14–17 longitudinal rows of smooth dorsal tubercles; and 18–20 broad lamellae beneath the fourth toe. *Gekko takouensis* sp. nov. is the second endemic gekkonid discovered in the Ta Kou Nature Reserve, *Cyrtodactylus takouensis* Ngo & Bauer being the first.

Keywords

Conservation; *Gekko takouensis* sp. nov.; granitic outcrop; Ta Kou Mountain; Vietnam

Introduction

The last decade has seen a dramatic increase in the number of gekkotan species described from Vietnam and 21 of the 22 new species described since 2000 are endemic (Geissler *et al.* 2009; Grismer & Ngo 2007; Heidrich *et al.* 2007; Hoang *et al.* 2007; Nazarov *et al.* 2008; Ngo 2008; Ngo & Bauer 2008; Ngo & Ziegler 2009; Nguyen *et al.* 2006; Orlov *et al.* 2007; Orlov *et al.* 2008; Rösler *et al.* 2005; Ziegler *et al.* 2002; Ziegler *et al.* 2008). The majority of these newly described geckos have limited distributions occurring on offshore islands or in isolated, rocky terrain such as karst outcrops and limestone caves. Most have been within the genera *Cyrtodactylus* Gray and *Cnemaspis* Strauch but two new *Goniurosaurus* Barbour, a new *Dixonius* Bauer, Good & Branch and two new *Gekko* Laurenti have also been described (Ngo *et al.* 2009; Ngo & Ziegler 2009; Rösler *et al.* 2005; Ziegler *et al.* 2008).

Correspondence to: Ngo Van Tri.

Note added in proof:

An additional *Gekko* species, *Gekko canhi*, was recently described from Vietnam. *Gekko takouensis* sp. nov. differs from *G. canhi* by the following characters: more precloacal pores (11–14 vs. 5); fewer interorbital (27–34 vs. 47–50); and fewer ventral scales across the belly between the ventrolateral folds (30–34 vs. 46–51).

Rösler, H., Nguyen, T. Q., Doan, K. V., Ho, C. T., Nguyen, T. T. & Ziegler, T. (2010) A new species of the genus *Gekko* Laurenti (Squamata: Sauria: Gekkonidae) from Vietnam with remarks on *G. japonicus* (Schlegel). *Zootaxa*, 2329, 56–68.

There are 13 species of *Gekko* in Indochina (Bauer *et al.* 2008; Bobrov 1995; Darevsky & Orlov 1994; Grossmann & Ulber 1990; Günther 1994; Ngo *et al.* 2009; Ota *et al.* 1995; Rösler *et al.* 2005; Szczerbak & Nekrasova 1993; Taylor 1962). Seven occur in Vietnam: *Gekko gecko* (Linnaeus), *G. palmatus* Boulenger, *G. ulikovskii* Darevsky & Orlov; *G. badenii* Szczerbak & Nekrasova, *G. grossmanni* Günther, *Gekko scientiadventura* Rösler, Ziegler, Thanh, Herrmann & Böhme and *G. russelltraini* Ngo, Bauer, Wood & Grismer (Bobrov & Semenov 2008; Ngo *et al.* 2009). *Gekko chinensis* (Gray) and *G. japonicus* (Schlegel) have historically been listed as occurring in Vietnam but their presence is doubtful (Ota *et al.* 1995).

Recent surveys conducted on Ta Kou Mountain, an isolated peak in Binh Thuan Province, southern Vietnam, have uncovered two new gecko species. *Cyrtodactylus takouensis* was described by Ngo & Bauer (2008) and here we describe another of the genus *Gekko* from the same region.

Materials and Methods

Field surveys were conducted in November 2005 and June – August 2009. Voucher specimens were collected by hand during the day and at night. Geographic coordinates and elevation were recorded using a Garmin III GPS. Photographs were taken using a Panasonic DMC-FZ30 digital camera and Sony DCR-TRV22. Specimens were first euthanized with tricaine methanesulfonate (MS-222) following Conroy *et al.* (2009) and subsequently fixed in 10% neutral buffered formalin, rinsed in water and finally stored in 75% ethanol. Liver samples, for use in subsequent DNA analyses, were stored in 95% ethanol and kept in cool conditions. All type specimens were deposited in the Zoological Collection of the University of Natural Sciences (UNS) in Hochiminh City, Vietnam.

The following measurements were taken with a digital caliper following the methods of Bauer *et al.* (2002; 2003): crus length (CrusL); ear length (EarL); eye to ear distance (EyeEar); forearm length (ForeaL); head depth (HeadD); head length (HeadL); head width (HeadW); internarial distance (Internar); interorbital distance (shortest distance between left and right superciliary scale rows) (Interorb); orbital diameter (OrbD); snout to eye distance (SnEye); nare to eye distance (NarEye); snout–vent length (SVL); tail length (TailL); trunk length (TrunkL). Measurements are reported to the nearest 0.1 mm, but mensural ratios calculated from raw measurements are reported to the nearest 0.01 mm. Measurements and scale counts were taken from the right side of each gecko unless otherwise noted. Scale counts and external observations on morphology were made using an Olympus SZ binocular dissecting microscope. The following meristic values were recorded: number of dorsal scale rows at midbody (MidBod); number of longitudinal rows of tubercles at midbody (TubRow); number of ventral scales between the ventrolateral folds at midbody (VentSc); number of chin scales bordering posterior margin of the first postmental scales (ChinSc); number of supralabials (SupraL); number of infralabials (InfraL); number scales across narrowest point of frontal bone (FrontSc); number of interorbital scale rows (IntOrb); number of subdigital lamellae under digit I of pes (LamPesI); number of subdigital lamellae under digit IV of pes (LamPesIV); number of postanal tubercles at tail base (PAT); number of preloacal pores (PrecP); and the number of middorsal scales rows in the third caudal whorl (Whorl3).

Comparisons were made with specimens in the Zoological Museum of the National University of Hanoi and the Zoological Collection of University of Natural Sciences, as well as original species descriptions and other published faunal and taxonomic treatments (e.g., Bauer *et al.* 2008; Darevsky & Orlov 1994; Grossmann & Ulber 1990; Günther 1994; Ngo

et al. 2009; Ota *et al.* 1995; Rösler *et al.* 2005; Szczerbak & Nekrasova 1993; Taylor 1962; Zhou *et al.* 1982).

Systematics

Gekko takouensis sp. nov.

Figs 1–2.

Holotype

Zoological Collection of University of Natural Sciences in Hochiminh City UNS 0491, adult male (Fig. 1A); Ta Kou Mountain, Ham Thuan Nam district, Binh Thuan Province, Vietnam, approximately 425 m elevation (10°48.815'N, 107°53.718'E), collected by Ngo Van Tri, 30 June 2009.

Paratypes

UNS 0489–90 and 0492–0497 (Figs 1B, C), adult males except for UNS 0495, subadult male, were also collected by Ngo Van Tri with holotype. UNS 0206 and UNS 0208 were collected on 2 November 2005.

Etymology

The specific epithet *takouensis* denotes the name of the mountain and nature reserve where the type series was collected. We suggest the following common names: English — Ta Kou Marbled Gecko; Vietnamese — Thằn lằn đá Tà Kôu.

Diagnosis

A medium sized *Gekko*, maximum SVL 107.0 mm, distinguished from its congeners by the following combination of characteristics: dorsum with 14–17 longitudinal rows of enlarged, smooth tubercles; 83–93 scale rows at midbody; 30–34 ventral scale rows between ventrolateral folds at midbody; 11–14 precloacal pores in males in an angular, continuous series; 18–20 subdigital lamellae on fourth toe; dorsal pattern of 5–7 whitish vertebral blotches between nape and sacrum, and 6–8 pairs of short, sometimes irregular spots or white bars on flanks between limb insertions.

Description of holotype

(Fig. 1A), UNS 0491, adult male. SVL of 95.5 mm. Head relatively long (HeadL/SVL: 0.26), wide (HeadW/HeadL: 0.66), somewhat depressed (HeadD/HeadL: 0.40); head distinct from neck, snout tapering, rounded at tip. Loreal and interorbital regions weakly inflated, frontonasal region strongly concave, snout elongate (SnEye/HeadL: 0.42), pointed, longer than eye diameter (OrbD/SnEye: 0.55); scales on snout and forehead small, granular, homogeneous; scales on snout larger than those on occipital region except for scattered smooth tubercles (approximately 2–3 times size of adjacent scales); 29 interorbital scale rows. Eye large (OrbD/HeadL: 0.23); pupil vertical with crenulated margins when closed and round when opened to maximum; superciliaries smooth, short, bearing several minute conical spines posteriorly. Ear opening oval, obliquely oriented, small (EarL/HeadL: 0.11); eye to ear distance longer than diameter of the eye (EyeEar/OrbD: 1.56). Rostral quadrangular, much wider (3.1 mm) than high (1.3 mm), with a median shallow groove. Supranasals are in broad contact and a small scale lies in the shallow cleft between the dorsal portions of the supranasals (Fig. 2A); rostral in contact with supralabial I and supranasals; nostrils round, each surrounded by supranasal, rostral, first supralabial and two enlarged postnasals; 3–4 rows of small scales separate the orbit from supralabials. Mental

triangular, wider (2.4 mm) than deep (2.2 mm); anterior pair of postmentals elongated (2.8 mm long, 0.8 mm wide), each bordered anteromedially by mental, medially in broad contact with other postmental, bordered anterolaterally by first infralabial, laterally by second postmental, posteriorly by three enlarged chin scales (Fig. 2B); 13 (right) to 15 (left) supralabials, 12 sublabials on both sides; 19 scale rows on the frontal bone and 29 interorbital scales in the closest distance between two eye edges.

Body robust, relatively short (TrunkL/SVL : 0.46) with weak ventrolateral folds. Dorsal scales smooth, round, granular and juxtaposed; 83 dorsal scale rows at midbody, intermixed with enlarged, smooth tubercles (2–5 times the size of adjacent scales, smaller on flanks, and smallest in occipital region) extending from occipital region to tail base; tubercles in 16 rows at midbody (Fig. 2C). Ventral scales much larger than dorsals, smooth, relatively hexagonal, and imbricate, largest posteriorly; 31 scale rows across venter between ventrolateral folds (Fig. 2D); gular region with relatively homogeneous, smooth scales. Thirteen precloacal pores arranged in angular series; somewhat enlarged scale rows extending posteriorly from pore-bearing scales to the anterior of cloacal lip (plate 2F); no enlarged femoral scales. Scales on palms and soles smooth, flattened, round, subimbricate without enlarged tubercles; smooth, flattened, subimbricate scales on venter of fore and hind limbs.

Limbs long and relatively robust (ForeaL/SVL = 0.13; CrusL/SVL = 0.17). Digits moderately dilated, all bearing curved claws except the first finger and first toe; the following number of broad lamellae occur beneath each digit (17–13–15–18–14 manus; 15–15–19–18–18 pes); two to five narrow lamellar rows between base and digits; weakly developed interdigital webbing. Length of digits (measurement in mm in parentheses): manus: IV(7.8) > III(7.5) > V(6.3) > II(6.2) > I(4.3); (pes): V(9.6) > IV(9.5) > III(9.0) > II(7.8) > I(5.2).

Original tail relatively robust with shallow groove on its dorsum (Fig. 2G), tapering towards the tip; tail longer than snout vent length (TailL/SVL : 1.25), two smooth postanal tubercles on the right side and three on the left side (Fig. 2E). Tail segmented; each tail segment approximately 10 dorsal scales rows (Fig. 2G) and 3 transversely enlarged subcaudal scales in length; scales on the tail dorsum heterogeneous – rectangular to hexagonal, juxtaposed. Subcaudal region with 80 median enlarged transverse plates (Fig. H).

Coloration

(in preservative) Dorsal color is gray with a series of 5–7 whitish irregular vertebral blotches between the nape and sacrum. The anterior most dorsal blotch is occasionally in contact with a light-colored nuchal loop, although the nuchal loop is not always present. Dorsal blotches extend along the tail, becoming ring-like although not fully encircling the tail and occasionally chevron-shaped. There are 6–8 pairs of light-colored, small, irregular bars or spots along the flanks between limb insertions. The dorsum has scattered dark brown or black flecking. Limbs are colored as dorsum with irregular white or light gray blotches and dark brown to black flecking. Multiple white or light gray, irregular blotches on the head interspersed with dark brown or black flecks and vermiculations. Two whitish, parallel stripes extend from the rostrum to the eye on each side of the head. One is an unbroken stripe from nare to eye while the other is broken and spotty and runs from the nasals to above the orbit. A light-colored nuchal loop is present in some individuals although it is often broken and irregular. Ventral coloration is whitish with occasional, small, dark brown or black speckles under the chin and belly.

In life, dorsal color is light to medium brown, with a green or olive overtone. Limbs are colored as the dorsum with irregular light blotches and dark speckling. Iris color is variable from chestnut brown to slate gray.

Variation

Variation in meristic and mensural characters among the type series is shown in Table 1. All individuals have a single small scale in the shallow cleft between the dorsal portions of the supranasals, except UNS 0492, which has 4 small scales. There is considerable variation in the shape and size of the dorsal blotches (Fig. 1) with some individuals having round or oval shaped blotches (Figs. 1A and C) while another has irregular shaped blotches with jagged edges (Fig. 1B). There is also diel variation in color with some geckos becoming lighter in color at night, a phenomenon that is common among many gecko species (Beebe 1944; Chan *et al.* 2006; Vitt *et al.* 2008).

Natural History

Gekko takouensis sp. nov. were active mostly at night although some individuals could be observed hiding in rock crevices during the day. The holotype and paratypes UNS 0491–UNS 0497 were collected on the rocky outcropping around the entrance of Hang To cave on Ta Kou Nature Reserve after a heavy rain. UNS 0206 and UNS 0208 were collected during the day while collecting *Cyrtodactylus takouensis* in November 2005. Another individual was observed at dawn and photographed living in the hole of a fig tree with *Gekko gecko*. Eggs, two to a cluster, were found glued on the sides and under large rocks. Several individuals were observed feeding on insects at night under the lamps and neon light on the wall and ceiling of the Linh Son Pagoda close to Hang To cave. *Gekko takouensis* sp. nov. is sympatric with the recently described *Cyrtodactylus takouensis* (Ngo & Bauer, 2008). The two species occur in close proximity with each other although *G. takouensis* sp. nov. typically occupies the cave entrance while *C. takouensis* lives in deeper parts of the cave, emerging to the mouth of the cave only at night.

Comparisons

Among its Vietnamese congeners, *Gekko takouensis* sp. nov. may be distinguished from *G. scientiadventura* by the presence of dorsal tubercles; from *G. palmatus* by the lack of broad webbing between the toes; from *G. gecko* by smaller adult SVL (107 mm maximum SVL vs. 173 mm maximum SVL – from Rösler *et al.* (2005)) and lack of large, mucronate dorsal tubercles; from *G. ulikovskii* and *G. badenii* by lack of narrow bands on dorsum. *Gekko takouensis* sp. nov. appears closely related to *G. grossmanni* and *G. russelltraini*, but differs from *G. grossmanni* in the following characters (in adults): larger size (89.7–107.0 mm SVL vs. 71.4–89.4 mm SVL); the number of interorbital scale rows (27–34 vs. 38–48); the number of scales across the belly between the ventrolateral folds (30–34 vs. 27–30); and the number of dorsal scales at midbody (83–90 vs. 94–115). *Gekko takouensis* sp. nov. differs from *G. russelltraini* in the following characters (in adults): larger size (89.7–107.0 mm SVL vs. 70.3–82.9 mm SVL); the number of scales across the belly between the ventrolateral folds (30–34 vs. 28–30); the number of dorsal scales at midbody (83–90 vs. 90–107); the number of precloacal pores (11–14 vs. 8–11); and the number of chin scales that border the first postmental (3–4 vs. 5–7).

Discussion

We have described a new species in the genus *Gekko* from Ta Kou Mountain, a granitic peak isolated from the Truong Son Mountain range. This new species is sympatric with the recently described *Cyrtodactylus takouensis* (Ngo & Bauer 2008). Ta Kou Mountain is about 50 – 60km from another isolated peak, Chua Chan Mountain in Dong Nai province,

where another pair of cave dwelling *Cyrtodactylus* and *Gekko* were discovered, *Cyrtodactylus huynhi* (Ngo & Bauer 2008) and *Gekko russelltraini* (Ngo *et al.* 2009). Isolated mountains have harbored other recently described geckos (eg: *Gekko badenii* Szczerbak & Nekrasova 1993; *Cyrtodactylus badenensis* Nguyen *et al.* 2006; *Cyrtodactylus nigriocularis* Nguyen *et al.* 2006). Mountain regions have long been known to host endemic species and elevational environmental gradients have been suspected as a contributing factor in the speciation of montane forms (Kozak and Wiens 2007). This phenomenon is thought to be even more important in the tropics where climatic zonation is more extreme than in temperate regions (Janzen 1967). While the exact speciation mechanisms that occur in montane regions may vary (Moritz *et al.* 2000) the importance of mountains to generating and fostering biodiversity is undeniable. Geckos may be particularly subject to montane endemism and diversification due to substrate specificity associated with a highly evolved digital adhesive mechanism. “Substrate islands” have been implicated in the high levels of species diversity of geckos in Southern Africa (Bauer 1999; Bauer *et al.* 2007) and Southeast Asian *Cyrtodactylus* (Bauer 2003; Ngo and Bauer 2008) and could also be important to other Southeast Asian gekkotans.

Conservation Status

Habitat surrounding the nature reserve has been degraded by the removal of rocks at the foot of Ta Kou Mountain for use as building material and the conversion of forest to banana plantations. *Gekko takouensis* sp. nov. is also threatened by collecting. Most geckos are hunted for food although some are also taken for the pet trade. Conversations with gecko trappers indicate that the price paid for geckos is 70.000–100.000 VND/kg or \$4–6 USD/kg. Preliminary monitoring of the wild population of *Gekko takouensis* sp. nov. since 2005 suggests these activities, particularly hunting, appear to be reducing the size of the wild population. We recommend continued monitoring of the *Gekko takouensis* sp. nov. population and a preliminary assessment of its conservation status that considers the potential impact of further habitat loss/alteration and continued commercial exploitation.

Acknowledgments

We thank Dr. Hoang Nghia Son, Director of the Institute of Tropical Biology, Vietnamese Academy of Sciences and Technology for permission to conduct this research. A.M. Bauer, L. L. Grismer and an anonymous reviewer provided helpful comments. We thank Dr. Hoang Duc Huy (Faculty of Biology of the University of Natural Sciences in Hochiminh City). We would also like to thank L. L. Grismer for providing MS-222. N.V.T was supported by Small Grant Number RD78 from WWF, USA. T. G. was supported by Grant Number T32DE007288 from the National Institute of Dental & Craniofacial Research. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute of Dental & Craniofacial Research or the National Institutes of Health.

References

- Bauer AM. Evolutionary scenarios in the *Pachydactylus* group geckos of southern Africa: new hypotheses. *African Journal of Herpetology*. 1999; 48:53–62.
- Bauer AM. Descriptions of seven new *Cyrtodactylus* (Squamata: Gekkonidae) with a key to the species of Myanmar (Burma). *Proceedings of the California Academy of Sciences*. 2003; 54:463–498.
- Bauer AM, Lamb T, Branch WR. A revision of the *Pachydactylus serval* and *P. weberi* groups (Reptilia: Gekkota: Gekkonidae) of Southern Africa, with the description of eight new species. *Proceedings of the California Academy of Sciences*. 2006; 57:595–709.
- Bauer AM, Sumontha M, Pauwels OSG. A new red-eyed *Gekko* (Reptilia: Gekkonidae) from Kanchanaburi province, Thailand. *Zootaxa*. 2008; 1750:32–42.
- Beebe W. Field notes on the lizards of Kartabo, British Guiana, and Caripito, Venezuela. Part 1. Gekkonidae. *Zoologica (NY)*. 1944; 29:145–160.

- Bobrov VV. Checklist and bibliography of the lizards of Vietnam. Smithsonian Herpetological Information Service. 1995; 105:1–28.
- Bobrov VV, Semenov DV. Lizards of Vietnam [in Russian]. *Tovarishstvo Nauchnih Izdanii KMK*, Moscow. 2008:226. XVIII pls.
- Bourret R. Notes hépetologiques sur l'Indochine française. XII. Les lézards de la collection du Laboratoire des Sciences Naturelles de l'Université. Description de cinq espèces nouvelles. *Bulletin Général de l'Instruction Publique*. 1937; 16(9):3–22. annexe.
- Bourret R. Notes hépetologiques sur l'Indochine française. XX. Liste des reptiles et batraciens actuellement connus en Indochine française. *Bulletin Général de l'Instruction Publique*. 1939; 19(4): 49–60. annexe.
- Chan SKF, Cheung K, Ho C, Lam F, Tang W. The geckos of Hong Kong. *Hong Kong Biodiversity*. 2006; 13:1–9.
- Conroy CJ, Papenfuss T, Parker J, Hahn NE. Use of tricaine methanesulfonate (MS222) for euthanasia of reptiles. *Journal of the American Association for Laboratory Animal Science*. 2009; 48:28–32. [PubMed: 19245747]
- Darevsky IS, Orlov NL. Eine bemerkenswerte neue, großwüchsige Art der Gattung *Gekko*: *Gekko ulikovskii* sp. nov. aus Zentralvietnam. *Salamandra*. 1994; 30:71–75.
- Geissler P, Nazarov R, Orlov NL, Böhme W, Phung MT, Nguyen QT, Ziegler T. A new species of the *Cyrtodactylus irregularis* complex (Squamata: Gekkonidae) from southern Vietnam. *Zootaxa*. 2009; 2161:20–32.
- Grismer LL, Ngo VT. Four new species of the gekkonid genus *Cnemaspis* Strauch, 1887 (Reptilia: Squamata) from Southern Vietnam. *Herpetologica*. 2007; 63:482–500.
- Grossmann W, Ulber T. Ein neuer Gecko aus Zentral-Thailand: *Gekko siamensis* sp. nov. (Reptilia: Sauria: Gekkonidae). *Sauria*. 1990; 12(3):9–18.
- Günther R. A new species of the genus *Gekko* (Reptila, Squamata, Gekkonidae) from southern Vietnam. *Zoologische Anzeiger*. 1994; 233:57–67.
- Heidrich A, Rösler H, Vu NT, Böhme W, Ziegler T. Another new *Cyrtodactylus* (Squamata: Gekkonidae) from Phong Nha – Ke Bang National Park, Central Truong Son, Vietnam. *Zootaxa*. 2007; 1445:35–48.
- Hoang XQ, Orlov NL, Ananjeva NB, Johns AG, Hoang NT, Dau QV. Description of a new species of the genus *Cyrtodactylus* Gray, 1827 (Squamata: Sauria: Gekkonidae) from the karst of North central Vietnam. *Russian Journal of Herpetology*. 2007; 14(2):98–106.
- Janzen DH. Why mountain passes are higher in the tropics. *The American Naturalist*. 1967; 101:233–249.
- Kozak KH, Wiens JJ. Climatic zonation drives latitudinal variation in speciation mechanisms. *Proceedings of the Royal Society B-Biological Sciences*. 2007; 274:2995–3003.
- Moritz C, Patton JL, Schneider CJ, Smith TB. Diversification of rainforest faunas: An integrated molecular approach. *Annual Review of Ecology and Systematics*. 2000; 31:533–563.
- Ngo VT. Two new cave-dwelling species of *Cyrtodactylus* Gray, 1827 (Squamata: Gekkonidae) from Southwestern Vietnam. *Zootaxa*. 2008; 1909:37–51.
- Ngo VT, Bauer AM. Descriptions of two new species of *Cyrtodactylus* Gray, 1827 (Squamata: Gekkonidae) endemic to southern Vietnam. *Zootaxa*. 2008; 1715:27–42.
- Ngo VT, Grismer LL, Grismer JL. A new endemic cave dwelling species of *Cyrtodactylus* Gray, 1827 (Squamata: Gekkonidae) in Kien Giang Biosphere Reserve, Southwestern Vietnam. *Zootaxa*. 2008; 1967:63–72.
- Ngo VT, Bauer AM, Wood PL Jr, Grismer JL. A new species of *Gekko* Laurenti, 1768 (Squamata: Gekkonidae) from Dong Nai Province, Southeastern Vietnam. *Zootaxa*. 2009; 2238:33–42.
- Ngo VT, Ziegler T. A new species of *Dixonius* from Nui Chua National Park, Ninh Thuan Province, southern Vietnam (Squamata, Gekkonidae). *Zoosystematics and Evolution*. 2009; 85:117–125.
- Nguyen VS, Ho TC, Nguyen QT. Danh lục ech nhai va bo sat Viet Nam. A Checklist of Amphibians and Reptiles of Vietnam [in Vietnamese]. Agricultural Publishing, Ha noi. 2005:180.
- Nguyen NS, Orlov NL, Darevsky IS. Descriptions of two new *Cyrtodactylus* Gray, 1827 (Squamata: Sauria: Gekkonidae) from Southern Vietnam. *Russian Journal of Herpetology*. 2006; 13:215–226.

- Orlov NL, Nguyen QT, Nazarov RA, Ananjeva NB, Nguyen NS. A new species of the genus *Cyrtodactylus* Gray, 1827 and redescription of *C. paradoxus* Darevsky et Szczerbak, 1997 [Squamata: Sauria: Gekkonidae] in Southern Vietnam. Russian Journal of Herpetology. 2007; 14(2):145–152.
- Orlov NL, Nguyen QT, Nguyen TT, Ryabov SA, Ho TC. A new species of *Goniurosaurus* (Squamata: Sauria: Eublepharidae) in North Vietnam. Russian Journal of Herpetology. 2008; 15(3):229–244.
- Ota H, Lau MW, Weidenhöfer T, Yasukawa Y, Bogadek A. Taxonomic review of the geckos allied to *Gekko chinensis* Gray, 1842 (Gekkonidae: Reptilia) from China and Vietnam. Tropical Zoology. 1995; 8:181–196.
- Rösler H, Ziegler T, Vu NT, Herrmann HW, Böhme W. A new lizard of the Genus *Gekko* Laurenti, 1768 (Squamata: Sauria: Gekkonidae) from the Phong Nha – Ke Bang National Park, Quang Binh province, Vietnam. Bonner Zoologische Beiträge. 2005; 53:135–148.
- Szczerbak NN, Nekrasova OD. A contribution to the knowledge of gekkonid lizards of southern Vietnam with description of a new species (Reptilia, Gekkonidae) [in Russian]. Vestnik Zoologii. 1993; 1994:48–52.
- Taylor EH. New Oriental reptiles. University of Kansas Science Bulletin. 1962; 43:209–263.
- Vitt L, Magnusson EE, Pires TCA, Lima AP. Guia de Lagartos da Reserva Adolpho Ducke, Amazônia Central. Áttema Design Editorial, Manaus. 2008:176.
- Zhou KY, Liu YZ, Li DJ. Three new species of *Gekko* and remarks on *Gekko hokouensis* (Lacertiformes, Gekkonidae). Acta Zootaxonomica Sinica. 1982; 7:438–446. pls. 1–2. [English translation by Ota, H., Hikida, T. & Lue, K. (1989) *Smithsonian Herpetological Information Service*, 77, 1–10, 2 pls.].
- Ziegler T, Rösler H, Herrmann HW, Vu NT. *Cyrtodactylus phongnhakebangensis* sp. n., ein neuer Bogenfingergecko aus dem annamitischen Karstwaldmassiv, Vietnam. Herpetofauna. 2002; 24(141):11–25.
- Ziegler T, Nguyen QT, Schmitz A, Stenke R, Rösler H. A new species of *Goniurosaurus* from Cat Ba Island, Hai Phong, northern Vietnam (Squamata: Eublepharidae). Zootaxa. 2008; 1771:16–30.

Figure 1.
Gekko takouensis sp. nov. (A) Holotype UNS 0491 in hand; B: Paratype UNS 0492 photographed *in situ*; C: Paratype UNS 0489 showing the lighter coloration often observed at night. Photographed *in situ* under a neon light on the dormitory wall of the Linh Son Pagada on Ta Kou Mountain.

Figure 2.

Detailed photographs of the holotype of *Gekko takouensis* sp. nov., UNS 0491 (A) Detail of the rostrum, note the small scale in the dorsal cleft of the supralabials; B: Ventral view of the chin illustrating three enlarged postmental scales; C: Dorsal scalation at midbody, note smooth, enlarged rows of tubercles; D: Ventral view at midbody illustrating subimbricate ventral scales; E: Dorsal scalation of the base of the tail showing two smooth, postanal tubercles; F: Ventral view of the colocal area showing 13 precloacal pores; G: Dorsal view of the tail, note segmented scales and lack of tubercles; H: Ventral view of the tail illustrating the enlarged median subcaudal scales.

Figure 3.
(A) Habitat of *Gekko takouensis* sp. nov. on Ta Kou Mountain; (B) Map showing the type locality of *Gekko takouensis* sp. nov. in Binh Thuan province, Vietnam.

Table 1

Mensural and meristic data for the type series of *Gekko takouensis* sp. nov.

	Holotype UNS 0491	Paratype UNS 0489	Paratype UNS 0490	Paratype UNS 0492	Paratype UNS 0493	Paratype UNS 0494	Paratype UNS 0495	Paratype UNS 0496	Paratype UNS 0497	Paratype UNS 0206	Paratype UNS 0208	Min–Max $\bar{x} \pm S.D.$
Sex	M	M	M	M	M	M	M	M	M	M	M	11M
SVL	95.5	100.1	97.7	99.2	93.3	107.0	81.3	89.7	97.3	106.5	99.6	81.3 – 107.0 97.0 $\pm$ 7.3
TailL	119.0	118.8	113.2	111.0	91.4 (Reg)	115.5	100.5	102.0	92.7	128.6 (broken)	73.8(R)	100.5 – 128.6 113.6 $\pm$ 3.3
HeadL	25.0	24.9	24.7	25.5	24.3	28.4	21.1	24.0	26.2	27.5	25.6	21.1 – 28.4 25.2 $\pm$ 0.6
HeadW	16.4	17.9	17.4	16.8	16.5	20.1	13.6	16.5	17.8	19.5	16.7	13.6 – 20.1 17.2 $\pm$ 1.7
HeadD	10.1	10.4	10.3	10.5	10.0	12.7	8.5	9.4	10.3	10.8	10.2	8.5 – 12.7 10.3 $\pm$ 1.0
SnEye	10.4	10.1	10.5	10.5	9.7	11.7	8.8	9.9	10.8	11.7	10.9	8.8 – 11.7 10.5 $\pm$ 0.8
NarEye	8.2	8.1	8.2	8.6	7.7	9.1	6.7	7.8	8.8	9.3	8.8	6.7 – 9.3 8.3 $\pm$ 0.7
OrbD	5.7	6.0	5.8	6.1	5.5	6.2	4.7	5.1	5.7	6.2	5.9	4.7 – 6.2 5.7 $\pm$ 0.5
EarL	2.7	2.7	2.6	2.8	2.7	3.4	2.2	3.0	2.6	3.3	2.7	2.2 – 3.4 2.8 $\pm$ 0.3
EyeEar	8.9	9.3	8.8	9.6	9.0	11.1	7.6	8.7	8.9	10.9	8.7	7.6 – 11.1 9.2 $\pm$ 1.0
Interorb	8.2	8.0	7.9	8.8	8.3	9.8	6.9	8.0	8.5	10.2	8.5	6.9 – 10.2 8.5 $\pm$ 0.9
Internar	2.4	2.9	2.6	2.7	2.5	3.0	2.1	2.2	2.5	2.8	2.6	2.1 – 3.0 2.6 $\pm$ 0.2
ForeaL	12.7	12.9	12.3	12.0	12.5	13.8	10.5	12.4	13.4	13.7	12.5	10.5 – 13.8 12.6 $\pm$ 0.9
CrusL	16.3	16.4	15.1	14.5	16.5	17.1	13.4	15.4	16.4	18.2	16.2	13.4 – 18.2 16.0 $\pm$ 1.3
TrunkL	43.8	44.6	44.3	44.5	43.7	49.7	35.2	42.8	44.0	49.1	43.7	35.2 – 49.7 41.1 $\pm$ 3.7
Midbod	83	93	90	90	85	85	83	87	90	92	85	83 – 93
TubRow	16	17	16	16	16	16	14	16	16	16	17	14 – 17

	Holotype UNS 0491	Paratype UNS 0489	Paratype UNS 0490	Paratype UNS 0492	Paratype UNS 0493	Paratype UNS 0494	Paratype UNS 0495	Paratype UNS 0496	Paratype UNS 0497	Paratype UNS 0206	Paratype UNS 0208	Min – Max $\bar{x} \pm \text{SD}$.
VentSc	31	31	34	30	32	33	30	32	32	31	30	30 – 34
ChinSc	3	4	4	3	3	4	3	3	2	4	3	3 – 4
SupraL	13/15	13/14	14/13	13/13	15/14	14/14	14/14	14/15	14/16	14/14	13/14	13–15/13–16
InfraL	12/12	12/11	12/12	12/12	13/12	12/12	12/11	12/12	12/12	11/12	10/12	10–13/11–12
FrontSc	19	22	20	18	18	24	19	20	18	20	19	18 – 24
IntOrb	29	29	31	34	27	27	27	28	27	30	27	27 – 34
LamPesI	15	14?	16	14	15	16	16	16	16	15	15	14 – 16
LamPesIV	18	18	18	18	19	18	18	18	19	20	19	18 – 20
PrecP	13	14	12	12	12	12	11	12	11	13	13	11 – 14
Whorl3	10	10	10	10	11	10	11	9	9	11	10	9 – 11